



.

.

·

,

25331

Digitized by the Internet Archive in 2016

https://archive.org/details/b2203898x_0001

BOTANIST'S GUIDE

THROUGH

ENGLAND AND WALES.

BY

DAWSON TURNER, F.R.S. A.S. & L.S.

Member of the Imp. Acad. Naturæ Curiosorum, And Honorary Member of the Dublin Society, &c. &c.

 $\mathbf{A} \mathbf{N} \mathbf{D}$

LEWIS WESTON DILLWYN, F.R.S. & L.S.

VOL. I.

LONDON: PRINTED AND SOLD BY PHILLIPS AND FARDON, CEORGE YARD, LOMBARD STREET.

1805.

A DATE OF A DATE OF A DATE OF



NISTORICAL MEDICAL

PREFACE.

IF utility has generally been allowed to those works in various branches of science, which, without aiming at critical investigation, tend to direct the Traveller in the path of his researches, it will, we trust, scarely be denied to a similar attempt in that particular department, where, more than in all others, the time and labor of the student are employed in seeking after the means of prosecuting his studies, and where it not unfrequently happens that the merely meeting with the object in view is in itself the ultimate end and reward proposed. Under this impression it is hoped, that, at a period when pursuits in Natural History become almost daily more prevalent. the present work will not be deemed an unacceptable offering to the Public; and if, by removing some

VOL. I.

PREFACE.

of those difficulties which discourage the beginner, or by affording to the experienced Botanist the means of more successfully prosecuting his inquiries, it tend to promote the study of that interesting and amusing science, we fhall have attained to the extent of our wishes; for, though we have found by experience that *in tenui est labor*, we are well aware that the remainder of this sentence cannot be applicable to a work, the very nature of which precludes all display of knowledge or investigation, and forbids us to look for credit beyond that of careful and industrious compilers.

HENCE, however, arises the advantage that our responsibility, as authors, is confined to the mere circumstance of having faithfully copied and properly arranged, the materials before us, a task we have endeavoured to fulfil with as much accuracy as can fairly be expected in a work of this nature. There are indeed some errors, which must be regarded as inseparable from our undertaking, and for which it is particularly necessary to deprecate the severity of critical censure. We have found it impossible to

avoid mistakes respecting the names of places, the orthography of which is in many instances so little settled, that they have frequently been differently written by our different correspondents; nor have we known where to apply for an infallible standard, the most perfect Index Villaris having afforded us comparatively but trifling assistance. We have therefore in this, as well as in almost every other instance, endeavoured strictly to confine ourselves to the task of copyists, and have scarcely ever ventured intentionally to deviate from the examples we have received, even though they have involved the necessity of spelling in more than one manner names which we had reason to believe were intended to be the same. A second, and considerable source of error, lies in the places of growth being sometimes so loosely defined, that it has not been in our power to determine with certainty to what division of our work a plant really belonged : this has been particularly the case, when, as often occurs, it is simply stated that the plant in question grows in the neighbourhood of some Town, which is situated perhaps on the borders of two or more Counties. From a somewhat similar cause we fear

b 2

we may have been led to introduce *habitats* as distinct, which, though they are really the same, have been described to us in words somewhat different. In such circumstances, even when we apprehended this to be the case, we have still considered it a lesser evil to subject ourselves to the charge of useless repetition, than to incur any hazard of depriving our readers of information, which might possibly prove serviceable; and it must be sufficiently evident that, although we might often have reason to suspect this error, it could but seldom be in our power to ascertain it. These, and other inaccuracies, we flatter ourselves with the hopes of amending, should the approbation of the Public ever call upon us for a second edition : in the mean time we shall lose no opportunity of correcting whatever we may ourselves discover to be erroneous, and we earnestly request the assistance of our Botanical friends for the same purpose.

With regard to the plants selected as most uncommon, we dare not indulge the hope that our list will meet with universal concurrence; for the rarity of vegetables is evidently a term of comparative import, and there are few British species, which, however abundant in some districts of the kingdom, are not scarce in others.

Our object has been to avoid, as far as possible, introducing those, which, though confined to peculiar situations, are in these situations almost always to be found; and, indeed, had we not thus restricted ourselves, it would have been necessary to have embraced in our catalogue the whole British Flora. At the same time it will be remarked, that by this plan many plants of rather unfrequent occurrence, are necessarily excluded: thus Cistus Helianthemum, Hippocrepis comosa, Asperula cynanchica, and others, which appear almost constantly with chalk, Beta maritima, Salsola Kali, with a long list of maritime vegetables, and those that are equally limited to marshes or sands, cannot but be rarities to the Naturalist, whose neighbourhood does not furnish similar stations; not to mention two very striking instances in Cotyledon Umbilicus and Digitalis purpurea, neither of which is to be found in this part of the kingdom, though the former of them in the Northern, and the

PREFACE.

latter in the Southern and Western Counties, are even more common than any Nettle or Thistle in the Eastern. We have made a point of noticing several remarkable varieties, under the conviction that they deserve more attention than has been paid to them by most Botanists; and in some few instances we have deviated from our general character of compilers, by subjoining, as notes, the observations of our friends, or our own, upon these and other subjects that did not appear unworthy of remark, to which, lest from their scattered situation they should be wholly useless, we have added a separate Index.

THE general Index of our work we have from its great length felt some reluctance in publishing; but our scruples on that head have given way to the consideration that by means of this alone an estimate can be formed of the comparative rarity of different plants, and that, were it omitted, it would be necessary to search through the whole work to ascertain the known stations of those individuals, which have hitherto been detected in only two or three places in the kingdom.

THAT part, which includes the class Cryptogamia, we regret to find even more faulty and imperfect than we had expected. I feel it, therefore, but justice to Mr. Dillwyn, to acknowledge, that the whole of this branch has rested exclusively with myself; and I must be allowed to add that the principal causes of my failure have been the present imperfect state of our knowledge respecting those plants, which made it difficult to determine what species are really most scarce; the small number of Botanists, from whom it was possible to procure assistance; and above all, the confusion of the fynonymy, arising in part from the inaccurate observations of the more early authors, but, in much greater degree, from the changes, which have been for some time past, and even now continue to be made among the names as well of the genera, as of the species, the unsettled state of which must long continue a material obstacle to the improvement we may hereafter expect in this department. This confusion of the synonymy has led me in a few instances to introduce the * same

* The instances, in which I have already detected this, are Neckera beteromalla called Fontinalis secunda; Lichen Letè-virens, pellitus, and cinereo-fuscus, called L. herbaceus, polyerbizos, and ferrugineus; and Fuscus rotundus called F. radiatus.

PREFACE.

plant in different places under different names, an error, for my acquittal from which I appeal with confidence to the students of Cryptogamic Botany, who will feel with me how difficult, I might indeed say how impossible, it is to avoid similar mistakes in compiling from authors, where the several terms are indiscriminately employed. The Fungi it was necessary entirely to omit, for the difficulties and obscurities which prevail in this tribe, are even greater than those among the other orders of the same class, and we have besides to lament the total want of that knowledge, which might enable us in any degree to unravel their perplexities. It might indeed have been possible to have rendered our Cryptogamic list somewhat more interesting and complete, by entering upon the discussion of doubtful or contested points; but, besides that this would have been in direct opposition to the principles of the work, it would not be practicable to treat these matters in a manner satisfactory either to our readers or ourselves within the limits of our book.

THE names here given to plants are almost uniformly those employed in the Flora Britannica or English Botany, except where the contrary is expressly mentioned, and except in the cases of individuals, to which those publications have not vet extended. We have, however, ventured to insert some phanogamous species which Dr. Smith has omitted, but in these instances we have always stated the reasons by which we were influenced, that other Botanists might judge of the propriety of allowing them hereafter to remain in the British Flora. Upon this head I must beg leave to offer it as my opinion that much less injury can be done to science by admitting twenty, the rights of which are questionable, than by being too scrupulous, and thereby incurring the hazard of excluding one which properly belongs to us. For, after all our care, it is impossible to ascertain how far any species is really indigenous or merely naturalized, nor, were this practicable, would it be worth the trouble; as the use of every Flora must always be either to enable the Botanist to ascertain the plants he has already found, or to point out those which he may expect to meet with, and whatever tends to circumscribe these effects cannot but be injurious to the utility of the work.

PEEFACE.

WE will not indeed deny that, independently of the consideration just mentioned, the reflection that we may possibly be instrumental in extending the limits of the Flora of our Country has afforded us no trifling pleasure; and it has also been a gratification to us that in some instances it has been in our power remove the uncertainty attending upon species, the existence of which in England was regarded as doubtful; nor ought we to omit mentioning it as a source of satisfaction, that in one or two instances we have been enabled to shew that the admission of particular plants originated only in mistake; for, whatever be the tendency of error, whether to favor or oppose our own wishes, the dispelling of it must ever be gratifying to an ingenuous mind.

IT is necessary to apologize to our Correspondents and our Readers that the distribution of materials adopted in our work has obliged us to omit some information, which we could not satisfactorily refer to its proper County. But this has been the case only in a few instances, except indeed among the older authors, nor did any other arrangement appear likely to be attended with equal convenience and with less objection.

A considerable difficulty arose in our minds whether habitats, which have been long known, should in general be referred to those who originally discovered them, or whether, where it was in our power, it would not be proper to quote the testimony of more recent observers; the latter plan may appear to rest upon better authority, and would evidently have the advantage of shewing that the plants continue to grow in the places assigned to them: but a sentiment of justice towards those of whose labors we now reap the fruits, and a feeling that we are acting in the manner in which we should ourselves wish others to act towards us, have induced us to forego these advantages, except in the cases of those stations which have been called in question, and except where the communications of our friends defined the spot more accurately than had been done before.

In addition to what has already been said, it is right to observe, that the present work is indebted for its

PREFACE:

origin exclusively to Mr. Dillwyn, who, like many other Botanists, accustoming himself as often as he left home to select the places of growth of those plants which were likely to occur to his notice, was induced to extract whatever referred to the most uncommon species from * Ray, Withering, &c, and reduce it to the following arrangement. The plan thus begun merely for his own use and amusement proved a more extensive task than he had conceived, and was regarded by his friends as likely to be of general utility; so that by their persuasion he was led to entertain the idea of preparing it for the public in the manner which is here effected by our united endeavours. Far, however, from confining ourselves to the selecting of that information which had been previously published, we have ventured to call upon many of the Botanists of the Kingdom for their assistance, and we are proud to own that our requests have on all sides been complied with in the most flattering manner.

* Whatever habitats are taken from Ray, Blackstone, Waring, Deering, Jacobs, and most of the more early writers, rest entirely upon the authority of Mr. Dillwyn.

XII

For the interest and trouble which these Gentlemen have been so kind as to take on our behalf, we feel ourselves extremely indebted, and, as the most effectual means of acknowledging our obligations, we have uniformly subjoined their names to the information they refpectively afforded : indeed, for the sake of our readers, we thought it right so entirely to conform to this principle, as even to place our own initials after those stations which depend upon our personal observation.

WE are not aware that a work similar to the present has ever yet appeared, either in England or any other Country, excepting Professor Martyn's *Herbationes* annexed to his *Plantæ Cantabrigienses*, an undertaking, commenced under precisely the same impression, but extremely limited in point of extent, and of which it is much to be regretted, that, though he admits the greater part to be a compilation, he has only in few instances quoted his authorities; so that he leaves his reader wholly ignorant as to what is derived from his own remarks, and how far the sources whence he has drawn his information are those on

PREFACE.

which safe reliance may be placed. We must own that we have more than once had reason to suspect the accuracy of his Catalogue; as when, for instance, we found Euphorbia Portlandica placed in an inland situation, Primula farinosa in so southern a County as Kent, Lycopodium annotimum on Iver heath, with some other instances, which we have noticed in their places, from a desire of inducing other Botanists to examine the fact, and ascertain the plants really intended.

BESIDES the general obligations which we owe to our different friends for the information they have afforded, particular thanks are due to Mr. Joseph Woods, jun. who took upon himself the trouble of selecting a considerable quantity of materials from Merrett's Pinax, and various topographical authors to which our situations, remote from the splendid libraries of the Metropolis, denied us the power of obtaining access. That much more information lies scattered in other similar publications cannot be doubted, and we fhould have been happy to have availed ourselves of it, but the difficulty of procuring these works must plead our excuse for not having quoted them.

xiv

IT has been suggested that the Botanist's Guide, by pointing out the flations of plants, may possibly lead to the extirpation of some which are least common. Could we allow force to this argument we would never permit the Manuscript to see the light; but we flatter ourselves that the Botanists of England do not deserve such a reproach, and we really believe that there is too much honor among the Votaries of Science to destroy, wantonly, for their own gratification, that which may hereafter prove a source of pleasure to many others : indeed, we are convinced that, far from wishing to eradicate, there is not a Naturalist in the Kingdom who would not feel gratified at any addition to the British Flora, and who would not, on the other hand, experience a sensible regret at the erasure of a single individual from it.

THERE are two other objects, which, though less immediately in our view, the present work may be calculated to answer. It will point out those parts of the kingdom hitherto least explored; and, by giving a comparative view of the productions of different Counties, may be useful in denoting their soils, and how far they may be adapted to improvements in agriculture or manufactures. It will also tend to prove to the Naturalists of other parts how zealously the study of indigenous Botany is here cultivated, and will shew them that, while the vegetables trodden under their feet are in many places neglected for the more gaudy productions of the hotter zones, the Botanists of England place their greatest pride and pleasure in the cultivation of English Botany. D. T.

Yarmouth, 10th August, 1805.

xvi

ANGLESEA, though divided only by a narrow channel from the vast ridge of Caernarvonshire mountains, contains in itself no remarkable elevation or diversity of surface. Its antiquities and mineral treasures have long since rendered it an object of curiosity to the traveller; but, with respect to its vegetable productions, little has either been discovered in it, or expected from it. If, therefore, the list we are here enabled to furnish be larger than might be expected from so small a County, we are happy to acknowledge ourselves wholly indebted for this circumstance to the kindness of the Rev. Hugh Davies, who has for many years, with equal industry and success, investigated the Botany of this Island, and favoured us with the communications, followed by his name, from a manuscript Flora, which will probably be one day presented to the public.

VGL. I.

flexilis.Rills and ditches, not rare.Id.VERONICA montana.Woods near Plâfnewydd; and Baron Hill.Id.UTRICULARIA minor.Peat-pits, frequent.Id.SALVIA pratensis.In dry meadows near Llanidan.Bingley.IRIS fætidissima.Priestholm Island.Rev. H. Davies.SCHENUS mariscus.Llanddyfnan, &c.Id
and Baron Hill. Id. UTRICULARIA minor. Peat-pits, frequent. Id. SALVIA pratensis. In dry meadows near Llanidan. Bingley. IRIS fatidissima. Priestholm Island. Rev. H. Davies. SCHŒNUS mariscus. Llanddyfnan, &c. Id. — rufus. Near Aberffraw; and Bodowen. Id. SCIRPUS multicaulis. Holyhead Mountain. Id. — pauciflorus. Near Bodorgan; and Aber- ffraw. Id. — acicularis. Abundant by the sides of shal- low pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
UTRICULARIA minor. Peat-pits, frequent. Id. SALVIA pratensis. In dry meadows near Llanidan. Bingley. IRIS fatidissima. Priestholm Island. Rev. H. Davies. SCHENUS mariscus. Llanddyfnan, &c. Id. rufus. Near Aberffraw; and Bodowen. Id. SCIRFUS multicaulis. Holyhead Mountain. Id. pauciflorus. Near Bodorgan; and Aber- ffraw. Id. ffraw. Id. ffraw. Id. ffraw. Id. ffraw. Id. ffraw. Id. ffraw. Id. PHLEUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
 SALVIA pratensis. In dry meadows near Llanidan. Bingley. IRIS fatidissima. Priestholm Island. Rev. H. Davies. SCHENUS mariscus. Llanddyfnan, &c. Id. rufus. Near Aberffraw; and Bodowen. Id. SCIRPUS multicaulis. Holyhead Mountain. Id. pauciflorus. Near Bodorgan; and Aberffraw. Id. acicularis. Abundant by the sides of shallow pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
 IRIS fatidissima. Priestholm Island. Rev. H. Davies. SCHENUS mariscus. Llanddyfnan, &c. Id. rufus. Near Aberffraw; and Bodowen. Id. SCIRPUS multicaulis. Holyhead Mountain. Id. pauciflorus. Near Bodorgan; and Aberffraw. Id. acicularis. Abundant by the sides of shallow pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
 IRIS fatidissima. Priestholm Island. Rev. H. Davies. SCHENUS mariscus. Llanddyfnan, &c. Id. rufus. Near Aberffraw; and Bodowen. Id. SCIRPUS multicaulis. Holyhead Mountain. Id. pauciflorus. Near Bodorgan; and Aberffraw. Id. acicularis. Abundant by the sides of shallow pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
 SCHŒNUS mariscus. Llanddyfnan, &c. Id. rufus. Near Aberffraw; and Bodowen. Id. SCIRPUS multicaulis. Holyhead Mountain. Id. pauciflorus. Near Bodorgan; and Aberffraw. Id. acicularis. Abundant by the sides of shallow pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
 rufus. Near Aberffraw; and Bodowen. Id. SCIRPUS multicaulis. Holyhead Mountain. Id. pauciflorus. Near Bodorgan; and Aberffraw. Id. acicularis. Abundant by the sides of shallow pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. IV. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
 SCIRPUS multicaulis. Holyhead Mountain. Id. pauciflorus. Near Bodorgan; and Aber- ffraw. Id. acicularis. Abundant by the sides of shal- low pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. IV. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
— pauciflorus. Near Bodorgan; and Aber- ffraw.Id.Ifraw.Id.— acicularis. Abundant by the sides of shal- low pools, reduced by the summer heat.Id.ERIOPHORUM polystachion. Spongy meadows near Beaumaris.Id.PHLEUM pratense. var. vivip. Found in this county byRev. W. Bingley.KNAPPIA agrostidea.Sandy pastures on the south west coast.Rev. H. Davies.Pox humilis.Common, about Beaumaris.Id.
ffraw. Id. — acicularis. Abundant by the sides of shal- low pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
low pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
low pools, reduced by the summer heat. Id. ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
ERIOPHORUM polystachion. Spongy meadows near Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
Beaumaris. Id. PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
 PHLEUM pratense. var. vivip. Found in this county by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
County by Rev. W. Bingley. KNAPPIA agrostidea. Sandy pastures on the south west coast. Rev. H. Davies. POA humilis. Common, about Beaumaris. Id.
KNAPPIA agrostidea.Sandy pastures on the south west coast.West coast.Rev. H. Davies.Poa humilis.Common, about Beaumaris.Id.
west coast. Rev. H. Davies. Poa humilis. Common, about Beaumaris. Id.
POA humilis. Common, about Beaumaris. Id.
FETTICA hromoldes. Millocks, not late.
uniglumis. Among Arundo arenaria, on
the south west coast Id.
loliacea. Meadows near Beaumaris. Id.
AVENA fatua. Among Potatoes, &c. very common. Id.
pubescens. Llanfihangel: Dinsilwy. Id.
LOLIUM arvense. Corn fields near Beaumaris, &c. Id.

TRITICOM loliaceum. The Green near Beaumaris,
and other sandy pastures. Rev. H. Davies.
RUBIA peregrina. Sea-rocks, south west of Beau-
maris. Id.
CENTUNCULUS minimus. Side of a small Lake,
near Bangor Ferry, by the road side. Id.
RUPPIA maritima. Between Beaumaris & Pen-
mon, &c. Rev. H. Davies. near Llan-
dduum
PULMONARIA maritima. North east coast. Rev. H.
Davies. Near Trefarthen, about half
way betwixt the ferry from Caernarvon
and Aurol - De
LYSIMACHIA thyrsiflora. Llyn Llechylched. Mr. Lbwyd.
VIOLA hirta. Penmon Park in D. J.
VIOLA hirta. Penmon Park, &c. Rev. H. Davies. GENTIANA preumonanthe Head
GENTIANA pneumonanthe. Heathy commons, Rhos- meirch.
l d.
OFNANTUR die U. H. Sieneyn. Id.
OENANTHE pimpinelloides. Salt marshes on the north east coast.
peucedanifolia Common : 1
DAMBUCUS Ebulus Noon Day
LINUM usitatissimum. Road sides and corn fields,
not uncommon.
DROSERA longifolia. Turf bogs. Id
GALANTHUS nivalis. Pentraeth. Id.
DRNITHOCALUM and all Id.
ORNITHOGALUM umbellatum. Maes y Porth woods,
near Newborough. Bingley,
CILLA verna. Cliffs on the sea shore. Dillenius.
B 2

5

Abundant in Anglesea, particularly along Rev. H. Davies. the sea coast. ASPARAGUS officinalis. Sands at Llanfaelog. Id. ----- var. β. Near the pool at Llanfaelog. Lhzvyd. RUMEX sanguineus. Woods above Baron Hill. Rev. H. Davies. ALISMA ranunculoides. Very common in Anglesea. Id. Aberffraw common; and POLYGONUM minus. Id. elsewhere in the county. SAPONARIA officinalis. Hedges at Llangadwalader, Id. and other places. CUCUBALUS baccifer. Gathered in hedges in Anglefea, and communicated to Dr. Richardson. Ray. Rev. H. Davies never found it. (This fpecies is not annual, * as Bingley afferts, as I have a plant now living, which has been in my poffession feventeen years. Mr. Griffith.) SILENE anglica. Not uncommon in this county. Rev. H. Davies. ARENARIA tenuifolia. On an Island in Llanddwyn, Bingley. near Newborough. EUPHORBIA portlandica. In feveral places in the Rev. H. Davies. south west coast. + Rosa tomentosa. Not rare in this county. Id. * It is marked perennial, by Withering, Smith, &c. † Another species of Rose, not yet, I believe, noticed by any

author, has been found in this county by the Rev. H. Davies, but he has not mentioned the exact habitat, and the specimens in my pos-

TORMENTILLA reptans. Near Llusdulas. Rev. H. Davies. GEUM rivale. Lligwy wood. Id.

CISTUS marifolius. Arthur's Round Table.

Amlwch. Rev. H. Davies. Sandy pastures on Mount Llech ddû, near Holyhead. Brown. (The habitat commonly assigned to this plant in the Isle of Man appears to be a mistake, arising from the mistranslating of the word Mona, by which Dillenius, and his friend, Samuel Brewer, always intended Anglesea, though it is still a question among the learned whether that Island or the Isle of Man was the Mona of the ancients. There are many similar errors among the British authors arifing from the fame circumstance.)

THALICTRUM minus.Near the sea at Llanfaelog,
and Red Wharf.Rev. H. Davies.RANUNCULUS lingua.Not rare in this county. At
Berw-marsh, &c. &c.Id.parviflorus.Towyn y Capel, near

Holyhead: Aberffraw; &c. Id.

session are not sufficiently perfect to enable me here to attempt a description. It comes nearest to *Rosa Hibernica*, a new species found abundantly in the North of Ireland by Mr. Templeton, and by him presented to the Dublin Society; but differs in having prickly capsules, and the upper surface of the leaves covered with an exceedingly fine minute pubescence. *D. T.*

Id.

MENTHA rotundifolia. Llanfair ynghorneoy; and
Bryn-Sieneyn. Rev. H. Davies.
LEPIDIUM latifolium. Aberffraw. Id.
COCHLEARIA officinalis. Priestholm Island. Id.
var. β . Beaumaris. Id.
anglica. Not uncommon in the county. Id.
danica. Not uncommon in the county.
Rev. H. Davies. Near Llanbadrick
church. Lhwyd.
IBERIS nudicaulis. Hedges at Pentraeth. Rev. H. Davies.
CRAMBE maritima. Sandy sea coast between
Rhuddgaer and Llanddwyn. Bingley.
SISYMBRIUM mononse. Abermenai. Lhwyd.
CHEIRANTHUS sinuatus. About Abermenai Ferry.
Ray. On Newborough sands. Bingley.
TURRITIS hirsuta. Old walls, and limestone rocks,
common. Rev. H. Davies.
The Discourse of Department Id
ERODIUM moschatum. Between Beaumaris and Penmon. Id.
ERODIUMmoschatum. Between Beaumaris and Pennon. Ia. maritimum. Near Beaumaris, plentifully. Id.
maritimum. Near Beaumaris, plentifully. Id.
GERANIUM pratense. Near Beaumaris, plentifully. Id. Id.
maritimum. Near Beaumaris, plentifully.Id.GERANIUM pratense.Near Beaumaris.Id.sanguineum.Very common along the
maritimum. Near Beaumaris, plentifully. Id. GERANIUM pratense. Near Beaumaris. Id. sanguineum. Very common along the sea coast. Rev. H. Davies. On cliffs
maritimum. Near Beaumaris, plentifully. Id. GERANIUM pratense. Near Beaumaris. Id. sanguineum. Very common along the sea coast. Rev. H. Davies. On cliffs between the Ferry from Caernaryon and
maritimum. Near Beaumaris, plentifully. Id. GERANIUM pratense. Near Beaumaris. Id. sanguineum. Very common along the sea coast. Rev. H. Davies. On cliffs between the Ferry from Caernaryon and Moel y Don. Bingley.
— maritimum. Near Beaumaris, plentifully.Id.GERANIUM pratense.Near Beaumaris.Id.— sanguineum.Very common along the sea coast.Rev. H. Dawies.On cliffs between the Ferry from Caernaryon and Moel y Don.Moel y Don.Bingley.LAVATERA arborea.Islets in the south west coast.

6

VICIA sylvatica. Old Park, and other places near	
Beaumaris. Rev. H.	Davies.
TRIFOLIUM ornithopodioides. On the green, by Beau-	
maris.	Id.
HYPERICUM androsæmum. Old park, and hedges	
near Beaumaris.	Id.
CARDUUS tenuiflorus. About Beaumaris, and many	
other places.	Id.
SANTOLINA maritima. Abermenai Ferry, plentifully	y. Ray.
GNAPHALIUM dioicum. Heathy Commons. Rev. H.	
GNAPHALIUM sylvaticum. Mill-dingle near Beau-	
maris.	Id.
INULA Helenium. Between the King's Bridge and	
Llangoed Mill, plentifully.	Id.
crithmoides. Sea rocks, particularly on the	
south west coast. Rev. H. Davies. Rocks	
at Llanddwyn.	Ray.
CINERARIA integrifolia. B. Declivities above the	C .
sea, near Holyhead. Rev. H. 1	Davies.
ANTHEMIS nobilis. Frequent on grassy Commons.	Id.
SATYRIUM viride. Meadows at Pentraeth: Llan-	
sadwrn, &c.	Id.
OPHRYS muscifera. Bodecho, and Llandyfnan bogs.	Id.
SPARGANIUM natans. Frequent in muddy ditches.	Id.
CAREX dioica. In boggy meadows, and peat	
grounds common.	Id.
teretiuscula. Bogs at Llansadwrn, and	
Llandegfan.	Id.

CAREX	limosa.	Parish of Llanbadrie	e. Rev.	H. Davies.
	fulva.	Frequent in wet mea	idows.	Id.
par line optimistic	extensa.	Near Aberffraw: H	[olyhead; a	und
	Penmo	11.		Id.
	* binervis	. Heathy banks, &c	. frequent.	- Id.
	distans.	Sea coast near Beau	maris, on	the
	way to Ba	ngor Ferry, in many	places.	Id.
	lævigata.	Dingle in Ty-fry d	demesne.	Id.
()	vesicaria	. Near Bodgylcha	d lake,	by
	Beaum	aris.		Id.
	ampullası filiformis.	Peat Bogs, very	y common	. Id.
LITTO	RELLA la	austris. Shallow lak	tes and in	un-
	dated c	ommons, frequent.		Id.
SALIX d	arenaria.	On the south west of	coast.	Id.
ATRIPLEX laciniata. Sea coast on the south west. Bingley.				
OSMUNDA lunaria. Not rare in old pastures. Rev. H. Davies.				
LYCOP	ODIUM Sa	laginoides. Aberffra	aw comm	on.
	Real 1	7 Danies On Town	n Trewar	. Thousd.

* Though this Carex is in general but little known, I have found it in every part of England, that I have visited, and I believe it to be one of the most common Britifh species. On the other hand, I consider the true *C. distans* to be still less understood, and extremely rare: never having seen it, except in very few places in the immediate vicinity of the sea. However they resemble each other when examined, they are at first sight to be distinguished, from the arilli and fquanae of the latter being always of a very pale green, while those of *C. binervis* are of a deep glossy brown colour. *D. T.*

ASPIDIUM thelypteris. Edge of Llwydiart lake, Pentraeth. Rev. H. Davies. ASPLENIUM marinum. Frequent on the fea rocks. Rev. H. Davies. Rocks about Priestholm Island; and at Llanddwyn. Ray. SCOLOPENDRIUM Ceterach. Holyhead mountain. Rev. H. Davies. PILULARIA globulifera. Spongy meadows and ditches. I.I. T. ANICTANGIUM ciliatum. Frequent on rocks and stones throughout the county. D. T. GRIMMIA Daviesii. 7 North east coast on limeverticillata. S rocks. Rev. H. Davies. ----- crifpula. TRICHOSTOMUM lineare. Common in the county. Id. ---- polyphyllum. Musc. Hib. DICRANUM rigidulum. Near Beaumaris. Id. --- cerviculatum. Banks of rivulets and ditches. Id. -* callistomum. Near the Point by Beaumaris. Id.

* Mr Davies has lately been so obliging as to furnish me with specimens of this plant, which I have carefully compared with Dr. Smith's Herbarium, and can fafely pronounce them all to be only *Dicranum varium*: but the *D. callistomum* of the Muscologia Hibernica, which was deferibed from a specimen in my own collection

DICRANUM viridissimum. common in the county.
Rev. H. Davies
POLYTRICHUM urnigerum. Ty-fry, near Beaumaris. Id.
HYPNUM myosuroides. Abundant on rocks about
Beaumaris. D. T.
LICHEN pilularis. Rocks in Bodowen Park.
the county. Id.
miniatus. On a dry rock at Llanfaethly.
Mr. S. Brewer.
omphalodes. Rocks, not unfrequent. Rev. H. Davies.
aquilus. Abundant on rocks, adjoining
the sea near Beaumaris. Id.
parietinus. A most beautiful variety of
this Lichen, not noticed, I believe, by
any author, with very narrow, linear, mul-
tifid fegments, of a deep orange colour
above, and very pale yellow below, is
common on rocks by the sea side. D. T.
tiliaceus. Rocks in this county. Rev. H. Davies.
sinualus. Stone hedges at Ty-fry, near
Beaumaris. D. T.

sent by Mr. Dickson, and which is the true plant of his fasciculi, I am now persuaded is nothing more than D. rigidulum, stripped of its barren elongated shoots, as maintained by Dr. Swartz. See Musc. Hib. p. 63.—Dicranum callistomum must therefore be expunged from the Muscologia. D. T.

LICHEN vulpinus. Branches of trees in this county.

Mr. S. Brewer.

Fucus sinuosus. y.	
——————————————————————————————————————	
tamariscifolius.	
membranifolius. S.	
esculentus. B.	
ceranoides.	
Laceratus. β.	
laciniatus.	On the shores of this
edulis.	county. Rev. H. Davies.
pinnatifidus. β.	
plumosus.	
flagelliformis.	
subfuscus.	
pedunculatus.	
kaliformis.	
<u></u> . β. ј	
Opuntia. Rocks was	hed by the sea oppo-
site Priestholm Isla	nd. Dillenius.
amphibius. On the s	shores of this county.
	Rev. H. Davies.
ULVA ligulata.]	
fistulosa.	
plumosa. > On the	shores of this county. Id.
filiformis.	
rubens.	
CONFERVA faniculacea. On	rocks in this Island,

and on Fuci among the rejectamenta of
the fea. Dillenius.
CONFERVA aruginofa. On the lesser Fuci on the
rocks by Llanfaethly. Id.
purpurascens. Near Llanfaethly. Id.
fluviatilis. Common in rocky rivulets. Rev. H. Davies.
• atra. In a well near the Parish of
Tregayan. Rev. H. Davies. In a fmall
fountain, which discharges itfelf into a
lake, called Gors Velen, near Llanfaethly;
and in a spring of the plain Gors-bach,
called Ffynnon bach y Lusg, half way
between the church and Trefaedog. Dillenius.
mutabilis. Springs in this county. Rev. H. Davies.
corallina.
setacea.
villosa.
pennata. On the sea shore. Id.
byssoides.
* echinulata. E. B. In a lake in this
county. Id.

* This little plant belongs to Dr. Roth's new genus, *Rivularia*, to which *Ulva incrassata*, *pruniformis*, and *pisiformis* also must be referred, and I must be allowed, on the authority of specimens from Mr. Davies, to express my doubts how far it is specifically distinct from the last of these plants. The magnified figure in English Botany is far from happy or accurate. D. T.

BEDFORDSHIRE.

THAT this work is to be regarded not so much as a specimen of what may be expected from the different Counties, as of the activity of those Naturalists who have hitherto explored them, is an observation that has repeatedly forced itself upon our minds in the course of our undertaking; and no where more strongly than in the instance of Bedfordshire, which, from its situation, its size, and the uniformity of its surface, appears to promise as little as any in the kingdom; and yet the unwearied activity of Dr. Abbot, almost wholly unassisted, has enabled him to publish a Flora containing no less than 1225 species. From this source we have necessarily extracted almost all our materials for the list of the County; and we beg leave to observe that the example of Dr. Abbot affords the most pleasing as well as convincing proof to every young Botanist, that, if he have but zeal and industry, there is no fear of his finding a sufficient field for discovery to repay his exertions.

CHARA hispida. B. Clapham Springs : Stevington	
bogs.	Abbot.
VERONICA montana. Woods and hedges at Ever-	
sholt.	Id.
UTRICULARIA minor. Bogs at Ampthill: Potton	
marshes.	Id.
SALVIA pratensis. In a pasture near Ford-end	
Farm. Rev. T. O.	Marsh.
IRIS fatidissima. Hedges at Bromham : Stevington :	
Pavingham.	Abbot.
SCHENUS albus. Ampthill moor: Bogs at Potton;	
and Aspley.	Id.
ERIOPHORUM polystachion. Bogs near Dunstable.	
Mr. L	Vickson.
PHLEUM paniculatum. Found in this county by	
Rev. Dr. Abbot, since the publication of	
the Flora Bedfordiensis, but we are un-	
acquainted with the habitat. (It is erro-	
neously observed in English Botany that	
I communicated this plant to Mr. Sower-	
by from Bedfordshire. $D. \mathcal{T}.$)	
AGROSTIS Spica venii. Corn fields at Ford-end	
farm : Luton downs.	Albot.
BROMUS arvensis. Pastures at Bromham-grange	
Farm.	Id.
pinnatus. Hedges at Clapham between	
Woods. Albot. In Houghton Conquest	
park. Mr. E. Forst	

AVENA fatua. Common in corn fields.	Abbot
pubescens. Common in pastures.	Id.
ARUNDO epigejos. Maulden wood.	Id.
Calamagrostis. Sheer-hatch wood.	Id.
ELYMUS europæus. Thurleigh, and Putnoe woods.	Id.
DIPSACUS pilosus. Wet hedges at Kemptston;	
Medbury; and Goldington. Abbot. About	
Toddington ; and Clophill. Mr. E. Forst.	er.iun.
	Abbot.
pusillum. Fields at Shefford.	Id.
CENTUNCULUS minimus. Boggy ground on Ampt-	
hill moor.	Id.
SANGUISORBA officinalis. Moist meadows at	
Bromham : Fenlake : Cow Meadows.	Id.
PULMONARIA officinalis. Between Thurleigh and	
Milton-Ernys.	Id.
ANAGALLIS arvensis. y. Oakley west field: Bar-	
ton Hill.	Id.
CAMPANULA latifolia. Woods and hedges at	A
Whipsnade, near Dunstable : Studham	
0012120.00	Id.
VIOLA hirta Barton LIII	Id.
RHAMNUS catharticus. Clapham ; and Ravensden,	
in woods and hedges. Abbot. Hedges	
near Elston. Mr. A.	Aikin.
VINCA major. Hedges at Ravensden, and Clapham.	Aubot.
CHENOPODIUM hybridum. Dunghill at Mill lane,	
Bedford	Id.

GENTIANA campestris. Barton Hill, very rare.	Abbot.
Cuscura europaa. On nettles, beans, &c. common	. Id.
BUPLEURUM rohundifolium. Corn fields at Barton	
Hill : Woburn.	Id.
CAUCALIS daucoides. Oakley west field, very rare.	
Mr. James	Payne.
latifolia. Corn fields at Oakley, and	
Thurleigh.	Abbot.
* HERACLEUM sphondylium. B. Hedges at Els-	
tow.	Id.
SIUM latifolium. Ditches, and banks of rivers, com-	
mon.	Id.
repens. Goldington Green : Stevington bogs.	Id.
SISON segetum. Road sides at Goldington, and	
Claphan.	Ib.
OENANTHE peucedanifolia. Potton marshes : Fen-	
lake meadow.	Id.
CICUTA virosa. Oakley fprings.	Id.
SCANDIX cerefolium. Goldington road side.	Id.
CARUM carui. Pastures at Thurleigh. Mr. John	Payne.
PIMPINELLA magna. Common in woods.	Abbot.

* Dr. Abbot gives it as his opinion that this plant is specifically different from the common *H. sphondylium*, and though I diffent with reluctance from so able a Naturalist, I do it with the more confidence, not only because of the opinion of many Botanists, but because I have myself more than once seen the leaves of both growing from the fame root. *D. T.*

17

SAMBUCUS Ebulus. Hedges, common,	Abbot.
LINUM usitatissimum. Corn fields at Barton Hill.	Id.
DROSERA longifolia. } Ampthill Bogs.	Id.
MYOSURUS minimus. Corn fields at Biddenham:	
Fenlake: Caldwell.	Id.
GALANTHUS nivalis. Sandy Hills.	Id.
ALLIUM carinatum. Keysoe Park. Rev. Dr.	Abbot.
FRITILLARIA Meleagris. Moist meadows at	
Bromham.	Abbot.
TULIPA sylvestris. Old chalk pits at Whipsnade, on	
the borders of Hertfordshire. Rev. D.	Jenks.
ORNITHOGALUM pyrenaicum. Between Eaton-So-	
con and Thurleigh. Abbot. Keyson	
Park Wood. Rev. T. O. Marsh. In the	
hedges and thickets, in the open fields,	
on the outside of the woods belonging to	
the late William Gery, Esq. of Bush-	
meed, near Eaton. Sir T. G. C	
umbellatum. Everton Heath.	Abbot.
nutans. A specimen was brought	
me by a merchant at St. Neot's, which he	
found growing wild in Eaton ford field.	
Sir T. G. Cullum. (From the different	
stations, in which this plant has been	
found, it appears to have a right to a place	
in any future edition of the Flora Bri-	
tannica.)	

VOL. I.

CONVALLARIA multiflora. Thurleigh Wood.	Abbot.
Acorus Calamus. Moat at the Hassocks farm.	Id.
RUMEX maritimus. Goldington green.	Id.
COLCHICUM autumnale. Wet meadows at Bar-	
ford; and Thurleigh.	Id.
ALISMA ranunculoides. Ampthill Bogs.	14.
EPILOBIUM angustifolium. Woods near Dunstable.	Id.
CHLORA perfoliata. Milton Hill: Bromham pas-	
tures.	Id.
VACCINIUM oxycoccus. Ampthill Bogs: Potton	
Marshes.	Id.
POLYGONUM minus. Wet places at Elstow; and	
Goldington.	Id.
Bistorta. Hyde Mill, Luton : Thur-	
leigh.	Id.
PARIS quadrifolia. Woods at Hawnes ; and Ren-	•
hold: Clapham Park wood.	Id.
MONOTROPA hypopithys. Frequent in this county	•
Hudson. Hostler's wood, near Marke	t
Street. Mr. Jo	s. Sibley.
PYROLA minor. Whipsnade wood : woods nea	r
Luton.	Abbot.
SAPONARIA officinalis. Hedges and walls at Brom	
ham; and Sharnbrook.	Id.
DIANTHUS deltoides. Everton : Pegnut Hill, nea	r
Potton.	Id.
SILENE noctiflora. Oakley West field.	Id.
quinquevulnera. Corn fields in this county	
Rev. L	r. Abbot

.

SILENE anglica. Corn fields at Woburn ; and Bar-	
ton.	Abbot.
STELLARIA glauca: Ampthill Bogs: Ford-End	
ditches.	Id.
ARENARIA tenuifolia. Newenham Wall.	Id.
SEDUM dasyphyllum. Walls at Market-Street.	Id.
album. Walls at Stevington ; and Sharnbrook	. Id.
SPERCULA pentandra. Sandy fields at Ampthill.	Iđ.
LYTHRUM hyssopifolium. Goldington green : Oak-	
ley west field.	Id.
EUPHORBIA platyphylla. In the corn near Bedford.	Id.
Cyparissias. Barton Leat Wood. Rev. Dr	Abbot.
Pyrus Aria. Hedges between Fenlake and Car-	
dington.	Abbot.
TORMENTILLA reptans. Common by road sides.	Id.
GEUM rivale. Putnoe wood.	Id.
PAPAVER hybridum. Corn fields, common.	Id.
DELPHINIUM Consolida. Corn fields at St. Leo-	
nard's farm near Bedford.	Id.
ANEMONE Pulsatilla. Barton Hill : Luton downs.	Id.
apennina. In a wood near Luton Hoe.	Ray.
RANUNCULUS Lingua. By rivers and ditches at	
Goldington and Oakley.	Abbot.
parviflorus. Common by road sides.	Id.
HELLEBORUS viridis. Pastures at Goldington.	Id.
fætidus. Thickets at Bromham:	
Stevington: Stagsden.	Id,
Ajuca Chamapitys. Dry pastures on Luton downs.	
Mr. 7.	Sibleva

TEUCRIUM Chamædrys. Road sides and walls at	
Warden.	Albet.
MENTHA sylvestris. In the orchard at Toddington	
parsonage. Mr. E. Forste	er, jun.
odorata. Undoubtedly wild in a ditch	
near Bedford. Rev. Dr.	Abbot.
STACHYS germanica. On a hill two miles from	
Bedford. Rev. John H	-
LEONURUS Cardiaca. Road fide at Ford-End.	
MELAMPYRUM cristatum. Frequent in woods.	
LATHRÆA squamaria. Hedges at Whipsnade ; and	
Studham.	Abbot.
ANTIRRHINUM Cymbalaria. Walls at Luton Hoe.	Id.
spurium. Corn fields at Clapham :	
Bromham: Stevington. Abbot. Near	
Milton Bryant. Mr. E. Fors	ter, jun.
minus. Oakley West Field.	
LIMOSELLA aquatica. Goldington Green.	Id.
DRABA muralis. On the Wardon Hills, near Bar-	
ton in the Clay. Rev. Di	r. Abbot.
IBERIS nudicaulis. Sandy banks : Ampthill : Clop	-
hill: Aspley.	Abbot.
SISYMERIUM sylvestre. Common in meadows.	Id.
ERODIUM moschatum. On Ampthill Warren.	
	r. Abbot.
GERANIUM phaum. Pastures and lanes at Even	-
sholt.	Abbot.
pratense. Meadows, common.	Id.

GERANIUM pyrenaicum. Rubbish at Luton Hoe:	
Ford-End.	Abbot.
LATHYRUS Nissolia. Wood near Ampthill. Coun-	
tess of Ossory. Hedges at Putnoe and	
Bromham.	Abbot.
sylvestris. Common in woods.	Id.
latifolius. Hawnes; and Bromham	
woods.	Id.
VICIA sylvatica. Sheerhatch; and Eversholt woods.	Id.
lathyroides. Grassy fpots at Ampthill :	
Maulden : Clophill.	Id.
ASTRACALUS glycyphyllos. Hedges at Bromham;	
and Oakley.	Id.
hypoglottis. Hills near Dunstable.	
Mr. Wood	ward.
TRIFOLIUM ochroleucum. Potton; Everton; Clap-	
ham.	.Id.
scabrum. Dry pastures at Potton:	
Sandy: Ampthill.	Id.
HYPERICUM dubium. Luton woods. Mr.	Vaux.
HYOSERIS minima. Corn fields, Ampthill: Maul-	
	Abbot.
HYPOCHÆRIS glabra. Ampthill; Sandy; Potton	
Heaths.	Id.
CARDUUS tenuiflorus. Borders of fields by the	
Alm's houses: Cox's pits.	Id.
eriophorus. Dry pastures, common.	Id.
GNAPHALIUM rectum. Clophill : Aspley : Southill.	Id.

INULA Helenium. Ravensden; and Stevington pas-	
tures.	Abbot.
ORCHIS ustulata. Barton Hill: Luton downs.	Id.
SATYRIUM viride. Stevington; Thurleigh; Blet-	
soe Meadows. Abbot. Near Luton	
downs. Mr.	Sibley.
OPHRYS Nidus-avis. Clapham Park Wood : Host-	
ler's wood: ChilternGreen, near Dunstable.	Abbot.
muscifera. Chalky pastures near Shefford.	
Mr. 1	Fowler.
apifera. Hawnes : Barton : Thurleigh:	
in dry spots.	Abbot.
MALAXIS paludosa. Potton Marshes.	Id.
SERAPIAS longifolia. Stevington bogs.	Id.
grandiflora. Woods in this county, fre-	/
quent.	Hudson.
TYPHA angustifolia. Knotting Green.	Abbot.
CAREX dioica. Turfy begs at Ampthill. Abbot.	
Near Toddington. Mr. E. Forst	
distans. (binervis?). Ditches at Stevington.	
strigosa. Putnoe wood: Renhold.	Id.
pendula. Woods, common.	Id.
ampullacea. Ampthill, and Potton moors.	•
Abbot. In a marshy meadow near Tod.	-
dington. Mr. E. Fors	ter, jun.
Buxus sempervirens. Hills near Dunstable.	
Mr. Wo	
MYRIOPHYLLUM verticillatum. Common in Ditche	es. Abbon

t

QUERCUS sessiliflora. Common in woods.	Abbot.
SALIX rubra. Near Bedford. Rev. Dr.	Abbot.
EQUISETUM sylvaticum. Hawnes wood : Anipt-	
hill park.	Abbot.
EQUISETUM fluviatile. Marshy places, common.	
Abbot. Near Toddington. Mr. E. Forst	er, jun.
bogs.	Albot.
OSMUNDA lunaria. Oakley West Field.	Id.
regalis. Aspley wood. Mr. Geo.	Dixon.
LYCOPODIUM inundatum. Ampthill bogs.	Abbot.
Aspidium thelypteris. Potton marshes.	Id.
PHASCUM alternifolium. Stevington bogs.	Id.
piliferum. 5 Clapham, on banks.	Id.
multicapsulare. Cart ruts in Clapham	
Park Wood.	Id.
crispum.	
curvisetum. } Near Bedford. Rev. Dr.	Abbot.
bryoides. At Clapham Springs.	Id.
GYMNOSTOMUM ovatum. Mud walls, common.	Abbot.
	Id.
* TRICHOSTOMUM trifarium. Stones at Clapham	
Springs.	Id.

• I have lately had occasion carefully to examine some specimens of this moss, which Dr. Abbot was so obliging as to send me three years ago, and they appear to mc distinct, as well from Hedwig's Swartzia trifaria, as from every other species, with which I am acquainted. D. T.

Тпосотомим fontinalioides. B. Musc. Hib. Oak-	
ley mill.	Abbot.
TORTULA rigida. Common on walls.	Id.
DICRANUM cerviculatum. Ampthill Heath.	Id.
HYPNUM ruscifolium. y. Common on mill wheels.	Id.
plumosum. Sheerhatch Wood.	Id.
JUNGERMANNIA pusilla. Hazelwood lane.	Id.
LICHEN cæruleo-nigricans. Barton Hill: Kempston.	Id.
bæomyces. Ampthill Heath, very rare.	Id.
cinereo-fuscus. (vernalis. Fl. Bedf.) Com-	
mon on rails and posts.	Id.
pulmonarius. Maulden Wood.	Id.
aphthosus. Ampthill and Maulden woods.	Id.
borizontalis. Heaths and woods at Clop-	
hill, and Sheerhatch.	Id.
alcicornis. Common on heaths.	Id.
jubatus. Ampthill Park pales.	Id.
ULVA crispa. Stones at Elstow.	Id.
pruniformis. On moss on the bath at Wrest	
Park.	Id.
CONFERVA reticulata. Ditches at Woburn and	
Aspley.	Id.

SCHOENUS albus. Bullmarsh Heath. Mr. Rudge. Bog in Windsor Great Park. Mr. Gotobed. SCIRPUS acicularis. Bullmarsh Heath. Mr. Rudge. _____ sylvaticus. Sonning. Id. AGROSTIS Spica-venti. Corn fields near Old Windsor. Mr. Gotobed. AVENA fatua. Common in fields about Windsor. 1d. pubescens. Pig's Green Lane near Reading. Mr. Fardon. Meadows about Windsor, common. Mr. Gotobed. DIPSACUS pilosus. Sonning Lane. Mr. Rudge. GALIUM verum. var. flore albo. (It differs from G. verum in its higher growth, as well as in the colour of the flowers, and their not being so abundant.) In Sonning Lane. Id. MENYANTHES nymphæoides. In the Thames at Ankerwick, near Windsor. Rev. Dr. Goodenough.

POLEMONIUM caruleum. On a ditch bank near Windfor, but may possibly be the outcaft of a garden. Mr. Gotobed. VIOLA hirta. In Basildon Wood. Mr. E. Forster, jun. Mr. Rudge. VERBASCUM Blattaria. Binfield. DATURA Stramonium. In a meadow near Reading. Mr. Fardon. RHAMNUS catharticus. Hedges near Reading. Id. THESIUM linophyllum. Chalk banks near Hurley. Mr. Gotobed. VINCA major. Hedge near Old Windsor. Id. CUSCUTA curopaa. On the west side of Bullmarsh heath. Mr. Rudge. Hedges near Eton Mr. Gotobed. in great plenty. CAUCALIS daucoides. Near Reading. Mr. Fardon. SIUM latifolium. Ditches about Radcot bridge, Id. near Faringdon. SISON segetum. Hedges near Eton, but rare. Mr. Gotobed. OENANTHE peucedanifolia. Moist meadows near Id. Eton. SCANDIX cerefolium. Hedge near Windsor, proba-Id. bly from a garden. DROSERA longifolia. Bullmarsh heath. Mr. Rudge. Bogs in Windsor Great Park with the beautiful Narthecium ossifragum. Mr. Gotobed. MYOSURUS minimus. Corn fields near Old Wind-Rev. H. Davies. sor. LEUCOJUM æstivum. Near Reading. Murray.

FRITIULARIA Meleagris. Meadows about Burghfield bridge, near Reading. Mr. Fardon. CONVALLARIA multiflora. Wood at Kintbury, near Hungerford. Mr. Gotobed. JUNCUS Forsteri. In the woods of Sir William East, Bart. at Hall place near Harley-ford. Mr. W. F. Drake. ALISMA Damasonium.On Wink field Plain, near Windsor. Rev. Dr. Goodenough. Ditches about Southcote, near Reading. Mr. Fardon. CHLORA perfoliata. Near Reading. Id. MONOTROPA bypopithys. Frequent in this county. Hudson. In a wood at Basildon house. Mr. E. Forster, jun. SAPONARIA officinalis. Hedge near Old Windsor. Mr. Gotobed. STELLARIA glauca. Moist meadows near Eton. 1d. ASARUM europæum. Between Maidenhead and Henley. Rev. Dr. Abbot. LYTHRUM byssopifolium. Once found in a wet waste piece of ground near Windsor. Mr. Gotobed. EUPHORBIA lathyris. Certainly wild, and perhaps indigenous in several places in, and near the parish of Ufton near Reading, springing up in dry stony thickets, periodically for a year or two after they have been cut, and till choked by briars, &c. Rev. Dr. Beeke. GEUM rivale. Banks of the Kennet, near Newbury. Mr. Gotobed.

MENTHA Pulegium. Tilehurst Common, near the workhouse. Mr. Fardon. Тнумия Nepeta. Road side near Wickham. Mr. Gotobed. ANTIRRHINUM Cymbalaria. On the walls of Rev. Dr. Goodenough. Windsor Castle. repens. Near Basildon. Mr.T.Forster, jun. minus. Corn fields near Eton. Mr. Gotobed. IBERIS amara. About Wallingford, plentiful, and undoubtedly wild. Dr. Smith. Chalky ground near Hurley. Mr. Gotobed. CARDAMINE amara. Banks of the Thames near Id. Windsor, rare. SISYMBRIUM sylvestre. Banks of the Thames near Id. Windsor, common. ERISYMUM cheiranthoides. Corn fields, and road Id. sides near Eton. TURRITIS glaora. By the road side near Maiden-Mr. E. Forster, jun. head. GERANJUM pratense. Meadows about Reading. Mr. Fardon. Common in meadows near the Thames by Windsor. Mr. Gotobed. LATHYRUS palustris. In a wood near Abingdon. Blackstone. ASTRAGALUS glycyphyllos. By the road side between Twyford and Reading. Mr.E. Forster, jun. LEONTODON palustre. Bog in Windsor Great Park, Mr. Gotobed. rare. CARDUUS pratensis. Boggy ground in Windsor Id. Great Park.

OPHRYS anthropophora. Near the highway from
Wallingford to Reading, on the Berk-
shire side of the river. Merrett.
muscifera. Hedge in a chalky soil near
Cookham. Mr. Gotobed.
SERAPIAS grandiflora. In the woods of Sir Wil-
liam East, Bart. at Hall place, near
Harley-ford. Dr. Smith.
* ARISTOLOCHIA clematitis. Hedge near Wind-
- sor. (A root brought into my garden pro-
duced three capfules last year.) Mr. Golobed.
CAREX curia. At Virginia Water in Windsor
Forest. Rev. Dr. Goodenough.
Pseudo-cyperus. Ditches near Eton, not
<i>Incommon.</i> Mr. Gotobed. <i>binervis.</i> Black park, and Windsor Great
Darl
Invitata Paris III' I C Da
ampullacea. At Virginia Water.
Rev. Dr. Goodenough.
MYRIOPHYLLUM verticillatum. Ponds and ditches
near Rhon
QUERCUS sessilifiora. About Newbury. Ray.
In a copfe at Kintbury, near Hunger-
foud
SALIX rubra. Between Maidenhead and Wind-
Por
Merret gives " Near Reading," as a habitat for this plant,
but Ray says it is a mistake

LYCOPODIUM inundatum. At Virginia Water. Mr. Got	oled.
POLYPODIUM vulgare, B. On the walls of Wind-	
sor castle. Mr. Manning	ham.
POLYTRICHUM urnigerum. On a steep wet bank	
near Virginia Water. Mr. Got	obed.
LICHEN aleurites. On Windsor Park pales. 1	d.
) Both these species bear fruit	
<i>physodes.</i> <i>prunastri.</i> Both these species bear fruit near Virginia Water, in Windsor Forest.	
prunastri.) Windsor Forest. 1	d.
cretaceus. On chalk about Eton.	d.
lynceus. Oaks in Windsor Park. D). T.

BRECKNOCKSHIRE.

THE very reverse of what we have observed on Bedfordshire holds good with regard to Brecknock, and the greater part of the Welsh counties. No where in our Island, south of the Tweed, does nature prefent more favorable opportunities for the cultivation, or more extensive fields for the discovery of her treasures, and yet, fingular as it may appear, no where has less attention been paid to them. This work, therefore, by shewing how little has hitherto been done, we truft will operate as a stimulus to those Botanists who have leisure and opportunity, to turn their minds to these parts of the kingdom; and we fhall certainly at all times consider ourselves obliged by communications, tending to enrich any future Edition, should the approbation of the Public ever require one. ERIOPHORUM vaginatum. Frequent in this county. L. W. D. CAMPANULA patula. Between Brecknock and Hay. Sir John Cullum. Frequent in the neighbourhood of Crickhowel. Rev. T. Butt. VIOLA lutea. On the Black Mountains. Merrett. SIUM verticillatum. Cader Arthur, near Brecon. * Evans. VACCINIUM Oxycoccus.) On the Black Mountains. SAXIFRAGA bypnoides. 5 Mr. 7. Woods, jun. SAPONARIA officinalis. In great quantities at Llangattock near Crickhowel. Mr. Fones. ARENARIA verna. Cader Arthur, three miles from Breeon. Evans. GEUM rivale. Usgoed Hendry. Mr. 7. Woods, jun.

* It would have been easy, by extracts from the works of Mr. Evans, not only to have swelled our list, as well in the present, as in every other Welsh county, but to have added several new species to the British Flora; and we should have been happy to have availed ourselves of the information he affords, especially as he professes to have undertaken his journies " for botanical purposes, and with a view to publishing at some future period a Flora Cambrica," had we not had too much reason to suspect his botanical accuracy, and indeed, had we not in some instances from the assurances of Botanists resident upon the spot, and from having ourselves followed his footsteps, been convinced that the plants are not always to be found in the stations he has assigned to them. This, therefore, we trust will plead our apology for not having quoted him more freely; and, as his Tours have been favorably received by the public, we hope, that in some future edition he will correct those inaccuracies, of which Naturalists have now cause to complain.

BRECKNOCKSHIRE.

TROLLIUS europaus. Vallies of the Black Moun-	
tains. Mr. J. Wo	ods, jun.
OROBANCHE minor. Clover fields near Crick	-
howel. Rev. Dr. Good	denough.
GERANIUM pratense. Plentiful about Istrad Veluid.	
L.	W. D.
OROBUS sylvaticus. Observed by Thomas Willifel	
below Brecknock Hills in the way to Car-	
diff. Ray. Frequent about Brecon. Mr	
HYPERICUM Androsæmum. About Brecon. L.	W. D.
INULA Helenium. Near the Iron Works, about	
a mile from Brecon. Dr.	Turton
SALIX herbacea. On a Mountain called the Beacon,	m ses post é
D	Jones.
EMPETRUM nigrum. On the Black Mountains.	J 01103 B
Mr. J. Wood	le inn
POLYPODIUM dryopteris. Woods in the deep	· · · · · · · · · · · · · · · · · · ·
Glens of this County.	Id.
phegopteris. At Ulgoed Hendry,	2 66 0
and Ufgoed Eynon Garn.	Id.
HYMENOPHYLLUM tunbrigense. Rocks in the	160
Vallies of the Black Mountains.	<i>T</i>)
and an and a structure of the structure	Id.

 \mathfrak{n}

BUCKINGHAMSHIRE.

VERONICA montana. Wood near Beaconsfield, Mr. Gotobed. rare. IRIS fatidissima. Chalk hill near Hedfor Wharf. Id. SCHOENUS albus. On Iver Heath. Mr.T.F.Forster, jun. SCIRPUS acicularis. Banks of the Thames, rare. Mr. Golobed. FESTUCA loliacea. Moist meadows near Eton. Id. AVENA fatua. Cornfields, common. Id. ----- pubescens. Meadows, common, Id. ELYMUS europaus. Marlow Wood. Id. DIPSACUS pilosus. Hedges near Ditton Park. Mr. G.tobed. At High Wycombe. Mr. 7. Rayer.*

* For all the information in this County, followed by the name of the late Mr. Jacob Rayer, we are indebted to Mr. Edward Forster, jnn.

Common near Bulstrode, in great plenty, with Radvola millegrana. Mr. Gotobed MENYANTHES nymphæoides. Datchet Common, plentiful. Id. VIOLA birta. Side of a wood near Hedgerly. Id. VERBASCUM Blattaria. Hedge near Eton-wick. Mr. Gotobed. Near Denham. Rohfon. DATURA Stramonium. On rubbish at Salt hill. Mr. Gotobed. RHAMNUS catharticus. Hedges near Eton. Id. THESIUM linophyllum. Chalk banks near Marlow. Id. VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greateft abundance. Mr. Gotobed. SIUM latifoliam. Ditches near Eton. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DATURA Ingifolia. On Iver Heath, and else- where. Mr. 7 Remut	CENTUNCULUS minimus. On Gerard's Cro	ſs
 with Radiola millegrana. Mr. Gotobed MENYANTHES nympheoides. Datchet Common, plentiful. Id. VIOLA birta. Side of a wood near Hedgerly. Id. VIOLA birta. Side of a wood near Hedgerly. Id. VERBASCUM Blattaria. Hedge near Eton-wick. Mr. Gotobed. Near Denham. Rohfon. DATURA Stramonium. On rubbish at Salt hill. Mr. Gotobed. RHAMNUS catharticus. Hedges near Eton. Id. THESIUM linophyllum. Chalk banks near Marlow. Id. VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton.wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- where 	Common near Bulstrode, in great plent	у,
plentiful.Id.VIOLA birta.Side of a wood near Hedgerly.Id.VERBASCUM Blattaria.Hedge near Eton-wick.Mr. Gotobed.Near Denham.Robfon.DATURA Stramonium.On rubbish at Salt hill.Mr. Gotobed.RHAMNUS catharticus.Hedges near Eton.Id.THESIUM linophyllum.Chalk banks near Marlow.Id.VINCA major.Hedge near Slough.Id.CHENOPODIUM olidum.On rubbish at Eton.Id.CUSCUTA curopæa.Hedges near Eton, in great plenty.Id.GENTIANA campestris.At the bottom of Short Hill near Ellesborough.Blackstone.TORDYLIUM maximum.Hedges near Eton.wick, in the greateft abundance.Mr. Gotobed.SIUM latifolium.Ditches near Eton.Id.CINN Latifolium.Ditches near Eton.Id.COENANTHE peucedanifulia.Moist meadows near Eton.Id.DROSERA longifolia.Cn Iver Heath, and else- urboreId.		
 VIOLA birta. Side of a wood near Hedgerly. Id. VERBASCUM Blattaria. Hedge near Eton-wick. Mr. Gotobed. Near Denham. Robfon. DATURA Stramonium. On rubbish at Salt hill. Mr. Gotobed. RHAMNUS catharticus. Hedges near Eton. Id. THESIUM linophyllum. Chalk banks near Marlow. Id. VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. CISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- uthere 		
 VERBASCUM Blattaria. Hedge near Eton-wick. Mr. Gotobed. Near Denham. Robfon. DATURA Stramonium. On rubbish at Salt hill. Mr. Gotobed. RHAMNUS catharticus. Hedges near Eton. Id. THESIUM linophyllum. Chalk banks near Marlow. Id. VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- uthare 	plentiful.	Id.
 VERBASCUM Blattaria. Hedge near Eton-wick. Mr. Gotobed. Near Denham. Robfon. DATURA Stramonium. On rubbish at Salt hill. Mr. Gotobed. RHAMNUS catharticus. Hedges near Eton. Id. THESIUM linophyllum. Chalk banks near Marlow. Id. VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- uthare 	VIOLA birta. Side of a wood near Hedgerly.	Id.
Mr. Gotobed.Near Denham.Robfon.DATURA Stramonium.On rubbish at Salt hill.Mr. Gotobed.RHAMNUS catharticus.Hedges near Eton.I.I.THESIUM linophyllum.Chalk banks near Marlow.Id.VINCA major.Hedge near Slough.Id.CHENOPODIUM olidum.CHENOPODIUM olidum.On rubbish at Eton.Id.CUSCUTA europæa.Hedges near Eton, in great plenty.Id.GENTIANA campestris.At the bottom of Short Hill near Ellesborough.TORDYLIUM maximum.Hedges near Eton-wick, in the greateft abundance.SIUM latifolium.Ditches near Eton.Mon.Id.SISON segetum.Hedges near Eton, but rare.Id.OENANTHE peucedanifolia.Moist meadows near Eton.Id.DROSERA longifolia.On Iver Heath, and else- whore	VERBASCUM Blattaria. Hedge near Eton-wick	
DATURA Stramonium. On rubbish at Salt hill. Mr. Gotobed. RHAMNUS catharticus. Hedges near Eton. I.I. THESIUM linophyllum. Chalk banks near Marlow. Id. VINCA major. Hedge near Slough. Id. VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- whore		
Mr. Gotobed. RHAMNUS catharticus. Hedges near Eton. I.I. THESIUM linophyllum. Chalk banks near Marlow. Id. VINCA major. Hedge near Slough. Id. VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- uthere		
RHAMNUS catharticus. Hedges near Eton. 1.1. THESIUM linophyllum. Chalk banks near Marlow. Id. VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton.wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- where Id.		
THESIUM linophyllum. Chalk banks near Marlow. Id.VINCA major. Hedge near Slough.Id.CHENOPODIUM olidum. On rubbish at Eton.Id.CUSCUTA europæa. Hedges near Eton, in great plenty.Id.GENTIANA campestris.At the bottom of Short Hill near Ellesborough.Blackstone.TORDYLIUM maximum. Hedges near Eton-wick, in the greatest abundance.Mr. Gotobed.SIUM latifolium.Ditches near Eton.Id.SION segetum.Hedges near Eton, but rare.Id.OENANTHE peucedanifolia.Moist meadows near Id.Id.DROSERA longifolia.On Iver Heath, and else- whereVince		
 VINCA major. Hedge near Slough. Id. CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- where 	THESIUM linophyllum. Chalk banks near Marlow	
 CHENOPODIUM olidum. On rubbish at Eton. Id. CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greatest abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SIUM latifolium. Ditches on Datchet Com- mon. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- uthore 	VINCA major. Hedge near Slough.	
CUSCUTA europæa. Hedges near Eton, in great plenty. Id. GENTIANA campestris. At the bottom of Short Hill near Ellesborough. Blackstone. TORDYLIUM maximum. Hedges near Eton-wick, in the greatest abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. SIUM latifolium. Ditches near Eton. Id. Mon. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- where		
plenty.Id.GENTIANA campestris.At the bottom of Short Hill near Ellesborough.Blackstone.TORDYLIUM maximum.Hedges near Eton-wick, in the greateft abundance.Mr. Gotobed.SIUM latifolium.Ditches near Eton.Id.SIUM latifolium.Ditches near Eton.Id.Mon.Id.Id.SISON segetum.Hedges near Eton, but rare.Id.OENANTHE peucedanifolia.Moist meadows near Id.Id.DROSERA longifolia.On Iver Heath, and else- uthereId.	Cuscura europæa. Hedges near Eton, in great	
Hill near Ellesborough.Blackstone.TORDYLIUM maximum. Hedges near Eton-wick, in the greatest abundance.Mr. Gotobed.SIUM latifolium.Ditches near Eton.Id		
Hill near Ellesborough.Blackstone.TORDYLIUM maximum. Hedges near Eton-wick, in the greatest abundance.Mr. Gotobed.SIUM latifolium.Ditches near Eton.Id	GENTIANA campestris. At the bottom of Short	
TORDYLIUM maximum. Hedges near Eton-wick, in the greateft abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. — repens. Sides of Ditches on Datchet Com- mon. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- where		
in the greatest abundance. Mr. Gotobed. SIUM latifolium. Ditches near Eton. Id. — repens. Sides of Ditches on Datchet Com- mon. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- where	TORDYLIUM maximum. Hedges near Eton-wick,	
SIUM latifolium. Ditches near Eton. Id. repens. Sides of Ditches on Datchet Common. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else- Vuboro	in the greatest abundance. Mr. G	Fotobed.
 repens. Sides of Ditches on Datchet Common. I.I. SISON segetum. Hedges near Eton, but rare. I.I. OENANTHE peucedanifolia. Moist meadows near Eton. I.I. DROSERA longifolia. On Iver Heath, and else- 	SIUM latifolium. Ditches near Eton.	LL
mon. Id. SISON segetum. Hedges near Eton, but rare. Id. OENANTHE peucedanifolia. Moist meadows near Eton. Id. DROSERA longifolia. On Iver Heath, and else-	repens. Sides of Ditches on Datchet Com-	
OENANTHE peucedanifolia. Moist meadows near Eton. DROSERA longifolia. On Iver Heath, and else- where	nion.	
Eton. Id. DROSERA longifolia. On Iver Heath, and else-	SISON segetum. Hedges near Eton, but rare.	Id.
DROSERA longifolia. On Iver Heath, and else-		
tthoro		Id.
where. Mr. 7 Power		
ATAL & J. A. D. VET.	where. Mr. J. 1	Rayer.

MYOSURUS minimus. Cornfields near Slough. Mr. Gotobed. LEUCOJUM astivum. In a moist meadow at Upton, remembered for 50 years by the prefent tenant of the Farm : alfo in a peatfield near Dorney. Id. FRITILLARIA Meleagris. Peat-field near Dorney. Id. ORNITHOGALUM umbellatum. Hedges between Datchet and Eton in abundance; and sparingly in the meadows near Eton. Id. CONVALLARIA multiflora. About High Wycombe. Withering. EPILOBIUM angustifolium. In a wood on Comb Hill near Ellesborough. Blackst ne. Hedge near Farnham, in abundance. Mr. Gotoled. CHLORA perfoliata. Gravel pit near Burnham; and at Cliffden. Mr. Gotobed. In the old chalk pit at Gerard's Crofs, plentiful. Blackstone. POLYGONUM Bistoria. Hedge in a wet meadow near Chippenham. Mr. Gotobed. PARIS quadrifolia. In a chalk pit in a wood near Little Missenden Church. Blackstone. MONOTROPA hypopithys. In Marlow Wood, in abundance. Mr. Gotobed. Common in Hudson. this County. ANDROMEDA polifolia. On Iver Heath. Mr. 7. Rayer.

PYROLA minor. In the woods at Loudwater near	
High Wycombe. Mr. Gotob	eď
DIANTHUS deltoides. On a wall at Langley near	
Iver. Blackstone. On Salt Hill. Mr. Gotob.	ed.
SILENE anglica. Cornfields near Burnham. Mr. J. Ray	
STELLARIA glauca. Moist meadows near Eton.	
Mr. Gotobe	d.
SPERGULA subulata. Dry Banks on Iver Heath. Id.	
THALICTRUM majus. By the river fide between	
Marlow and Templehouse. Rev. H. Davie	2.0
RANUNCULUS Lingua. Ditch at Wexham, rare.	3.0
Mr. Gotobec	,
parviflorus. Cornfield near Salt-hill, Id.	•
HELLEBORUS viridis. Woods near Denham.	
Blackstone. Woods about Great Marlow	
and High Wycombe. Mr. J. Rayer	
fatidus. Chalk-hill near Hedsor	
Wharf. Mr. Gotobed	,
MENTHA rotundifolia. Between High and West-	•
TT7 1	
<i>Tubra.</i> By the river side a mile below	9
Dapham	
Pulegium. Sides of ditches near Upton.	,
Mr. Gotobed.	
Гнумия Nepeta. Road side near Bulstrode. Mr.	
Gotobed. Between St. Peters and St.	
MELAMPYRUM cristatum. In a field that goes off	
an a nera that goes off	

BUCKINGHAMSHIRE.

Moreton Green in the road from Wend-
over to Ellesborough. Blackstone.
ANTIRRHINUM minus. Cornfields near Eton.
Mr. Gotobed.
THLASPI arvense. On the West side of Chalfont
St. Peters, abundantly. Blackstone. Corn-
fields near Marlow. Mr. Gotobed. About
Aylesbury. D.T.
IBERIS amara. Chalky ground near Marlow.
Mr. Gotobed.
nudicaulis. Salt-hill. Id.
DENTARIA bulbifera. In the woods at Loud-
water, between Beaconsfield and High
Wycombe, rare. Id.
CARDAMINE amara. In meadows subject to be
overflown by the river Coln near Coln-
brook, on the London side of the Town.
Sir J. Banks. Banks of the Thames,
rare. Mr. Gotobed.
SISYMBRIUM sylvestre. Banks of the Thames,
common. Id.
Irio. Road sides near Eton. Id.
ERYSIMUM cheiranthoides. Corn fields and road
sides near Eton and Datchet. Id.
TURRITIS glabra. Road sides, and old Gravel
pits near Burnham. Mr. Gotobed. About
Denham. Mr. J. Rayer.

ERODIUM moschatum. On the rubbish near Salt-	
hill. Mr. Go	tobed.
GERANIUM pratense. Meadows near the Thames,	
common.	Id.
LATHYRUS Aphaca. Among the Corn near Den-	
ham. Blackstone. Gravel pit near Burn-	
ham. Mr. Go	tobed.
sylvestris. In a hedge a little beyond	
Lathbury on the road to Bedford. Sir J. Cu	llum.
ASTRAGALUS glycyphyllos. At the bottom of	
Coomb Wood, near Ellesborough. Blacks	stone.
ANTHEMIS nobilis. Langley Heath, on the edges	
of the cart-ruts, common. Mr. Goto	obed.
ORCHIS militaris. Marlow Wood, in plenty. Mr.	
Gotobed. Woods between High Wy-	
combe and Great Marlow. Mr. J. R.	ayer.
SATYRIUM viride. Marlow wood, rare. Mr. Goto	obed.
OPHRYS Nidus avis. Marlow wood, frequent.	
Mr. Gotobed. Wood near Temple-	
house, plentifully. Rev. H. Da	vies.
monorchis. In a chalk pit by the road	
side at Gerard's Crofs. Black.	stones
muscifera. In a beech wood just below	
	đ.
apifera. On the chalk hill near Hedfor	
Wharf; and in a chalk pit near Bulftrode.	
Mr. Gotobed. At Gerard's Cross. Blacks	tone.
SERAPIAS grandiflora. Woods about Marlow, and	

BUCKINGHAMSHIRE.

other parts of the County. Hudson. Plenful in a beech wood just below the Duke of Bedford's seat at Cheyneis. Blackftone. CAREX Pseudo cyperus. Ditches near Eton, not uncommon. Mr. Gotobed. fulva. Wet meadow near Eton, called Chalvey mead. Mr. Gotobed. ampullacea. Chalvey-Ditch near Eton. Id. LITTORELLA lacustris. Langley Heath. Id. MYRIOPHYLLUM verticillatum. Ponds and Ditches near Eton. Id. OSMUNDA regalis. On Taplow Common. Mr. J. Rayer. LICHEN cretaceus. About Eton. Mr. Gotobed.

THAT our list of the more rare plants of Cambridgeshire should be so comparatively disproportionate to its extent will cease to appear remarkable, when it is considered how much its Botany has been cultivated, beyond that of any other part of England. Mr. Ray, who in 1660 published his Catalogue of Cambridgeshire plants, enumerated no less than 624 indigenous to the County, to which in 1685 he had made the addition of 102 more. Professor Martyn's Plantæ Cantabrigienses, edited in 1763, increased the number to 829, and that at a time when the whole Flora Anglica comprized only 1569; since which the indefatigable Mr. Relhan and his friends have made a farther addition of 525, raising the list of species contained in the last Edition of his excellent Flora Cantabrigiensis to 1334. This addition indeed is greatly to be accounted for from the furprizing increafe which our knowledge of Cryptogamic Botany

has of late years received; an increase still so rapidly continuing that the number of British plants is almost daily enlarged. The fuperiority of Cambridgeshire, however, though principally, is by no means wholly to be attributed to the able Botanists her University has produced : the soil is also peculiarly favorable: containing a variety of chalky hills, extensive heaths, rich pastures, cornfields, woods, sands, and bogs, so that there are few plants which may not here find a suitable situation. We have in this County in one respect deviated from our usual plan, by quoting Mr. Relhan, not only for the plants found by himself, but for those also of Ray, the two Profeffors, J. and T. Martyn, and other Botanists, a deviation to which we were led, not merely with the view of saving trouble, but because we understand that he has himself seen every species in the station he has assigned to it in his Flora, which gives the habitat so much greater authority.

 CHARA bispida. In the water-courfe by the side of Hinton Moor: Sawston Moor, &c. Relban.
 FRAXINUS excelsior. β. ramis pendulis. Gamlingay. Id.
 VERONICA spicata. On the wall of St. John's

College: In several places near Newmarket Heath, beyond Bottisham: in a close near the Beacon: among the Furze near Hare Park : about Horseheath Hall. Relban. VERONICA montana. In the woody part of the Devil's Ditch, on Newmarket Heath : Hall Wood; Linton: Gamlingay Park. Id. UTRICULARIA minor. Found by Mr. Dent on Teversham Moor. Ray. Hinton Moor, on the edge next the cornfields: Fulbourn, Sawston, and Chippenham Moors : Gamlingay quaking bogs. Relhan. CROCUS sativus. Cultivated formerly near Gogmagog Hills. Relban. I have often fought for it in vain in the neighbourhood. D.T. IRIS fatidissima. In a grove on the right hand side of the road leading from Cambridge to Hinton Church : Teversham : Fulborn : Triplow. Relhan. Schoenus mariscus. Fulborn, Hinton, Teversham, and Chippenham Moors : Isle of Ely. Relhan. So common on the Moors about Cambridge that it is often used in that Town for lighting fires. Mr. Crowe. compressus. Hinton Moor: between Little Shelford and Whittlesford : near Bottisham Load. Relban.

SCHOENUS albus. On boggy ground near Gamlin	I-
gay Park.	Relhan.
Scirpus acicularis. In a chalk pit at Upware.	
Rev. Mr	. Holme.
pauciflorus. Hinton, & Foulmire Moors	
Rev. Rd.	Relhan.
PHALARIS phleoides. Barrington Hill, near Hil	u.
dersham, in great abundance. Relban.	
On Chippenham Park Wall. Rev. J. 1	Hemsted.
PANICUM wiride. In the gravel pits near Chip-	
penham Park; and in the cornfields ad-	
joining.	Relhan.
* PHLEUM paniculatum. On Newmarket Heath.	
Hudson. Near Bourn Bridge. Mr.	Crowe.
ACROSTIS Spica-venti. Sandy cornfields.	Relhan.
CYNOSURUS cristatus. y. Fl. Cant. Fields and	
meadows near Cambridge, plentifully.	Id.
FESTUCA loliacea. Grauchester Meadow: Hin-	
ton, &c. not uncommon.	Id.
BROMUS multiflorus. Paradise, near Cambridge:	
Audrey Caufeway.	Id.
arvensis. By the side of the road to Im-	
pington: in a meadow by the side of the	
road to Barton: Long Stanton: not un-	
common near Cambridge.	Id.
pinnatus. Hedges and dry fields.	Id.

* Mr. Echan has often diligently sought for this plant in vain

Avena fatua. Cornfields at Barnwell, Hinton,
&c. Relhan.
pubescens. Dry pastures, &c. Id.
ARUNDO epigejos. Madingley, Kingston, and
Eversden woods: Comberton: Wood
Ditton : Isle of Ely. I.I.
Calamagrostis. In the bushy Closes at
the North side of Hardwick Wood: Wood
Ditton Park Wood, in the corner next
to the Common. Relban. Prickwillow,
Isle of Ely. Rev. Dr. Goodenough.
colorata. B. picta. Fulbourn, on the
left hand side of the footway leading to
the Mill. Relban.
LOLIUM arvense. Corn fields near the Hill of
Health, &c. Id.
HORDEUM maritimum. By the river side, below
Wisbech. Id.
TRITICUM loliaceum. Salt marshes near Wisbech. Id.
DIPSACUS pilosus. In Love Lane, Chesterton:
East side of Trumpington : Linton, by
the first bridge in the footway leading to
Fuldersham. Id.
GALIUM tricorne. Cornfields at Gogmagog
Hills: Hinton: Fulbourn: Linton. Id.
anglicum. Chippenham, on the Park
Wall: Wisbech. Id.
RUBIA perigrina. Frequent, plentiful, and lux-

uriant in the quick hedges at Crabmarsh, near Wisbech, in which neighbourhood it has not been cultivated above 50 years. Mr. W. Skrimshire, jun. CENTUNCULUS minimus. Gamlingay Bogs. Relhan. SANGUISORBA officinalis. King's Hedges: Whitwell: Ditton: Shelford: Long Stanton: Cottenham: &c. Id. POTAMOGETON beterophyllum. Burwell Fen. Id. ---- compressum. B. Fl. Cant. Queen's Green, &c. Relhan. (The leaves of this are fo much longer and broader than those of P. compressum, and the habit so different, that I cannot but suspect it is a distinct species. D. T.) RUPPIA maritima. Ditches below Wisbech. Mr. Skrimsbire. ANCHUSA semper-wirens. Emanuel College, in the Master's Close, under the Walnut Trees : Back yard of Christ's College. Relban. SYMPHYTUM officinale. B. Side of Dead-man's Pond, Wisbech. Mr. Skrimshire. ---- tuberosum. On the Fen Banks, mixed with S. officinale. MIr. Woodward. Asperugo procumbens. It used to grow near Newmarket, but has not been found there for many years. Rev. J. Hemsted. (Dr. Withering has quoted Mr. Woodward, as his authority for its growing in

the Church Yard, Newmarket, which
that Gentleman has desired us to say is a
mistake, as he never found it there.)
MENYANTHES nymphaoides. Rivers about Strea-
tham Ferry. Merret. In the Cam, be-
low Cambridge : Fens, near Ely : Little-
port : Old Bedford river. Relhan.
ANAGALLIS arvensis. y. In the Corn on the left
hand side of the road to Histon, a little
beyond the first closes: Linton, between
the Town and the Wood : near Kingston
Wood. Id.
CAMPANULA latifolia. Comberton, in a coppice
called Gaynes, and the field adjoining.
Rev. Mr. Newton. About Cheveley.
R.v. G. Cralbe.
VIOLA birta. Moor Barns: Hinton. Relban.
VERBASCUM Lychnitis. Waste places and rub-
bish. Mr. Lyons.
DATURA Stramonium. Wisbech. Mr. Skrimshire.
CHIRONIA pulchella. Hinton Moor: Tever-
sham: Hauxton. Relhan
LONICERA Caprifolium. Hinton, Chalk-pit Close;
and in another Coppice. Id.
RHAMNUS catharticus: Woods and hedges. Id.
THESIUM linophyllum. Gogmagog Hills: Trip-
low Heath: between Hinton and Ful-
bourn : Teversham : Bottisham : between

Linton and Hildersham: Newmarket Heath : Chippenham Moor. Relban. Abundant near Snailwell, 3 miles East of Newmarket. Rev. Fames Dalton. VINCA major. Girton: Madingley: Coton: Whittlesford : Rampton : Bottisham. Relban. HERNIARIA glabra. Near Newmarket. Rev. J. Hemsted. CHENOPODIUM hybridum. Banks of some watery pits between Ely and Stuntney : not uncommon in gardens. Relhan. ----- olidum. Under Jesus College Wall: under the wall of the Botanic Garden : Pound Hills, &c. Relhan. Cuscura europea. Madingley on Nettles: Barton on Caucalis anthriscus : Oakington on Humulus lupulus: Swavesey on Beans. Relban. Near Newmarket. Rev. J. Hemsted. * GENTIANA Amarella. var. flor. prolif. Coldhams: Gogmagog Hills: Borders of Hinton Moor: Chalk-pit close; &c. Relban.

* Mr. Relhan justly observes that the number of the segments of the Calyx and Corolla in this plant is not to be depended on. as he has seen quadrifid, and quinquefid flowers on the same individual. I have noticed the same thing, but Profeffor Martyn's remark that *G. campestris* and *Amarella* are not distinct is by no means well founded, for the habit of the two plants is very different, and in the latter the segments of the Calyx are equal, whereas in the former they are always alternately large and small. *D. T.*

BUPLEURUM rotundifolium. Gogmagog Hills,
by the lower road : Stapleford : Tever-
sham : between Quey Church and Bot-
tisham. Relhan.
tenuissimum. Salt water ditches
at Wisbech. Id:
CAUCALIS daucoides. Among the wheat, on the
left hand side of the road leading from
Cambridge to Gogmagog Hills, between
the 2d. and 3d. Mile-stone: Kingston:
Devil's Ditch: Anglesey; &c. Id.
latifolia. With the former King's
Hedges: Trumpington : Histon : Tever-
sham: Comberton: Kingston, &c. Id.
SELINUM palustre. On the sides of the Drain
running Eastward from Prickwillow.
Rev. Dr. Goodenough.
ATHAMANTA Libanotis. Chalk-pit Close, Hin-
ton: in some old chalkpits on the right
and left hand side of the road leading
from Hinton towards the road to Gog-
magog Hills, and on the Balks. (Not
now found on Gogmagog Hills.) Relkan.
HERACLEUM Sphondylium. B. Hinton, in a Close
on the right hand side of the Lane leading
from the Church towards Teversham. Id.
SIUM latifolium. Rivers and ditches. Id.
repens, Coldham Common, Id.

SISON S	egetum. Madingley Road, near Cam-	,
	bridge : Haddenham.	Relbac.
OENANT	HE pimpinelloides. Marshes and Moors,	
	common near Cambridge.	Id.
	- peucedanifolia. Granchester Meadows	s. Id.
CORIAND	DRUM sativum. In the cornfield, by the	:
	road to Hinton, between Cambridge and	
	the New Mill.	Id.
CICUTA	virosa. Fens between Ely and Prick-	-
	willow, in a creek over against the tiled	ł
	house on Rimney Bank. Relban. In	1
	the river about a mile from Prickwillow	F
	Bridge. Sir T. G.	Cullum.
CARUM	Carui. Christ College Pieces : Stourbridge	2
	Fair Green: Meadows near the Wind	
	mills, Wisbech.	Relhan.
PIMPIN	ELLA magna. Gamlingay wood : Hatley	y
	St. George: Linton.	Id.
SAMBUC	vs Ebulus. Madingley, near the Well	•
	near the road to Hinton: Oakington	:
	Barrington : Eversden ; &c.	Id.
	migra. β. fruciu albo. Orwell: Lin	-
		Id.
	y. laciniata. Near Wisbech.	. 1d.
	E reticulata. Tydd Marsh. Mr. Ski	
	usitatissimum. Cornfields, Wisbech: Out	
	well.	Relhan.

LINUM perenne. Gogmagog Hills: Chalk-pit
Close: between Hinton and Fulbourn. Relban.
DROSERA longifelia. Hinton, Teversham, and
Sawston Moors. Id.
anglica. Hinton and Sawston Moors. Id.
MYOSURUS minimus. Stourbridge Fair Green:
Oakington, in a lane leading from the
village to the Huntingdon road: Hazen-
field, in the road leading to Cambridge,
near the village, plentifully : Eversden. Id.
ALLIUM oleraccum. Hinton. Rev. Rd. Relhan
FRITILLARIA Meleagris. In some Closes at West
Hoe, near Linton. Relhan
ORNITHOGALUM pyrenaicum. Little Eversden, in
a bushy Close, near the Church. Ray.
umbellatum. Fulbourn. Relhan
Acorus Calamus. In the ditch opposite to the
great Gates of Trinity College walks: in
a ditch by Great Founders Closes near
the house in the fields.
FRANKENIA lævis. Tydd Gote, near Wisbech. Id.
RUMEX sanguineus. Eversden, Kingston, and
Hall Woods.
* maritimus. Hinton Moor : behind

* Mr. Relhan observes that it appears to him *Rumex palastris* would more properly be made a variety of this species 'The same' idea, I must confess, has repeatedly occurred to me, and as often as I have endeavoured to fix limits between the two plants, both which

Granchester Mill: Isle of Ely: Cam-	
bridge Common: Ditch near Christ	
College : Wisbech, &c.	Relhan.
COLCHICUM autumnale. Wood Ditton, in a Close	
on the South side of Mr. Eaton's House.	1d.
ALISMA ranunculoides. Hinton, and Shelford	
Moors: Isle of Ely.	Id.
CHLORA perfoliata. Hinton, and Teversham Moors:	
pastures near Madingley Gap, &c.	Id.
VACCINIUM Oxycoccos. Gamlingay Bogs.	Id.
POLYGONUM minus. Waterbeach.	Id.
Bistorta. Closes near How's House:	
Whitwell: Hinton: Shelford.	Id.
PARIS quadrifolia. Woods at Kingston; Evers-	
den; and Wood Ditton.	Id.
MONOTROPA Hypopithys. Madingley Plantations.	I.l.
SAPONARIA officinalis. Barnwell: Paper-Mills:	
Granchester: Madingley: Shelford:	
Hinton: Comberton, near the Church:	
Whittlesford.	Id.
flore pleno. Impington, near Cam-	
bridge. Rev. Rd.	Relhan.
DIANTHUS deltoides. Hildersham, on the furze	
hill next to Linton : near Whitewood,	
Gamlingay.	Relhan,

grow near Varmouth, I have always found the marks of distinction hitherto given uncertain, and not to be relied upon. D. T.

DIANTHUS caryophyllus. On the walls of Chippen-	
ham Park.	Relhan.
SILENE anglica. Chippenham Gravel-pit : New-	
market, in the ploughed land, on the	
right hand side, near the Turnpike.	Id.
Otites. Gravel-pits on the North side of	
Newmarket : Chippenham, on the Balks,	
plentifully.	Id.
noctiflora. Newmarket, in the ploughed	
fields on the right hand fide, near the	
Turnpike : near Catlidge Hall : between	
Snailwell and Chippenham: and between	
Chippenham Park wall and the Gravel-pit.	Id.
STELLARIA glauca. Between Ely and Stuntney :	
Isle of Ely: Bottisham Load: Bardolph	
Fen, near Wisbech.	Id.
ARENARIA tenuifolia. Near Gogmagog Hills:	
borders of Triplow Heath, &c.	Id.
SEDUM dasyphyllum. Fulbourn, on the north side	
of the Church, and on the Buttresses:	
on the Garden Wall of R. G. Townly,	
Esq. and the opposite wall.	Id,
album. Chatteress. (Not found there for	
many years.)	IJ.
LYTHRUM byssopifolium. Near the Brick Kilns	
in the road to Chesterton: In the ditches	
on the right hand side of the road to	
Histon, near the first pond : Hinton and	

Teversham Moors: Oakington, &c.
Relban. Near Newmarket. Rev. J. Hemsted.
EUPHORBIA platyphylla. On the north side of
Eversden Wood, in a field called the
Three-cornered ground. Relhan.
β . With α , and near
Gransdon Lodge. Id.
PRUNUS Padus. Hedges, near Wimpole. Id.
Pyrus torminalis. Gransden. Ray.
Rosa villosa. By the side of the swampy ground,
near the water running from the Trum-
pington road, by the first mile stone.
Rev. Mr. Holmes. White Wood, near
Gamlingay. Relban.
tomentosa. In the lane leading from the
Barton road towards Granchester: not
uncommon near Cambridge. Id.
POTENTILLA verna. Gogmagog Hills, near the
two small hills, as soon as the τ oad is
left. Id.
Pastures near Wood-
Ditton: Hall, Cat-
GEUM rivale. lidge, and Wood-
B. fl. plenis. Ditton Park Woods:
Shudy Camps.
GEUM rivale. ————————————————————————————————————
GLAUCIUM violaceum. Cornfields between Swaff-
ham and Burwell: Newmarket, in a

	field behind Mr. Vernon's house, probably	•
	not wild.	Relhan.
PAPAVE	R bybridum. Cornfields at Paradise, near	
	Cambridge: Hilton: Fulbourn: An-	
	glesey.	Id.
	- somniferum. Waterheath Fen: on the	
	banks of the Closes, which separate	
	Denny Farm from the Ely road : Ramp-	
	ton.	Id.
TILIA	microphylla. * Ventenat. White Wood,	
	near Gamlingay.	Id.
CISTUS	Helianthemum. B. fl. albis. Wood-	2100
	Ditton.	Id.
DELPHI	NIUM Consolida. By the lower road to	100
	Gogmagog Hills: Hinton: Trumping-	
	ton: Teversham: Swaffham: Bottisham,	
	&c.	Id.
STRATI	OTES aloides. Near the Bridge at Ely:	1.00
	Stretham Ferry: Rampton: Audrey	
	Causeway: Mepole: beyond Littleport:	
	March, &c.	Id.
ANEMOR	E Pulsatilla. Gogmagog Hills: Hilder-	10.
	sham: Bartlow: Barrington Hill, near	
	Linton : Newmarket Heath.	Id.
THALIC	TRUM minus. Saxon Lane, Newmarket;	Ille.
	and in a field behind Mr. Vernon's	
	house : Linton : Bartlow.	ŢĴ
		Id.

* See Annals of Botany, I: p. 209.

RANUNCULUS Lingua. Teversham Moor:	
Anglesey Abbey : Isle of Ely : Stretham	
Ferry: Audrey Causeway: Gamlingay. R	Relhan
parviflorus. Madingley : Trump-	
ington: Shelford: Toft: Gamlingay, &c.	Id.
HELLEBORUS viridis. Dry Pastures at Coton:	
Ditton: Bigwin Closes, near Ditton:	
Whitwell.	Id,
fatidus. Pastures and hedges at	
Hinton: Fulbourn: Triplow.	Id.
CALTHA palustris. flore pleno. Granchester	
Meadow.	Id.
Ajuga Chamapitys. On the Lays about the	
borders of Triplow Heath.	Id.
TEUCRIUM Scordium. In a ditch on the left	
hand side of the road leading from Cam-	
bridge to Histon, near the first pond:	
Waterbeach: Cottenham: Isle of Ely,	
&c.	Id.
MENTHA sylvestris. a. Behind the Mills at	
Granchester, Harston, and Barrington:	
Linton, near the first bridge in the road	
to Hildersham.	Id.
β. Behind Granchester Mill:	
Bottisham Load, a quarter of a mile	
below the Mill.	Id.
rotundifolia. Cottenham, in the outer-	

¥.

most Town-Close, next to Westwick:	
Exning.	Relhan
MENTHA piperita. Hauxton, in the running water,	
in the Village.	Id.
odorata. On the Milton side of the River,	
a little below Ditton.	Id.
birsuta. 8. Behind Granchester Mill:	
Pampisford.	Id.
near Stretham.	Id.
side of the road to Impington, a little	
way beyond the Pond : Fordham, on the	
South side of the River below the Town,	
in the way to Soham.	Iđ.
gracilis. Hauxton, in the Village, and	
on the Common : Shelford : by the river	
side, nearly opposite to Horningsea.	Id.
Pulegium. Near the Castle Hill, Cam-	
bridge : Denny Abbey : Harlston, on the	
Common, by the road side : Wisbech.	Id.
GALEOPSIS versicolor. Audrey Causeway: Wis-	
bech.	Id.
BALLOTA nigra. B. Near the Paper Mills: Barn-	
well: Ditton.	Id.
LEONURUS Cardiaca. On the back of the New-	
market road, beyond the Paper Mills: At	

the back part of Trumpington, towards Shelford. Relhan. THYMUS Serpyllum. E. Near the Nine Wells, by the footway to Shelford. Id. ----- Nepeta. Granchester : Hildersham : Linton, &c. Id. MELAMPYRUM cristatum. Madingley, Kingston, Eversden, Linton, Hall, and White Woods Id. ANTIRCHINUM Cymbalaria. Walls about Cambridge. Smith .- On the walls of the Great House at Drayton: Bartlow Church. Id. ------ spurium. Cornfields at Barnwell, Teversham, Histon, Madingley, &c. common near Cambridge. Id. ----- minus. Gogmagog Hills: Babraham : Hinton : Newmarket Heath. Id. LIMOSELLA aquatica. Watery places at Milton. Id. OROBANCHE elatior. Hauxton: Comberton Hill: Hill between Shelford and Stapleford. Id. ramosa. Hemp fields at Wisbech and Upwell. Mr. Skrimshire. ALYSSUM sativum. Crab Marsh, Wisbech: Chippenham Gravel Pit. Relban. LEPIDIUM latifolium. Leverington, near Wisbech. Mr. Skrimshire. LEPIDIUM ruderale. Wisbech, on the bank of the

river above and below the Town: near
the Turnpike at Tydd Gote. Relban.
COCHLEARIA anglica. Tydd Marsh, below Wis-
bech. Mr. Skrimshire.
IBERIS nudicaulis. Gamlingay, near the Wind-
mills, and by White Wood. Relhan.
ISATIS tinctoria. New Barns near Ely: Isle of
Ely. Relban. Near Wisbech. Mr. Woodward.
SISYMBRIUM sylvestre. Cambridge Common:
on the bank of the river Cam, between
Chesterton Sluice, and Barnwell Pool:
Ifle of Ely. Relhan.
Irio. On walls at Wisbech. Mr. Skrimshire.
Eaysimum cheiranthoides. In the Ofier Holts,
near Ely: and on the Bank of the river
between the bridge and the Town:
Prickwillow, near the bridge: Chatte-
refs. Relhan.
ARABIS Turrita. On Trinity and St. John's
College walls. Id.
TURRITIS birsuta. Near Bournbridge Turnpike :
Chippenham: Newmarket, on the walls
of the Red Lion Inn: Deadman's Lane,
Wisbech. Id.
GERANIUM phaum. Teversham, in the first Close
on the East side of the Church, near the
Fence of the Churchyard. Rev. Robert Forby.
pratense. Chesterton : Ditton : Hin-
ton: Histon: Girton: Madingley. Relhan.

x

GERANIUM sanguineum. By the Devil's Ditch:
Wood Ditton : in a wood between Stock-
worth and Chidley. Relhan.
LATHYRUS Aphaca. Meadows and Cornfields.
Relhan. Near Newmarket. Rev. J. Hemsted.
Nissolia. Near the House in the
fields, a little beyond the Turnpike, on
the left hand side of the road : Hadden-
ham. Relhan.
sylvestris. Borley Wood, near Lin-
ton: Kingston and Eversden woods:
Hatley St. George: near Castle-Camps,
in the way to Bartlow. Id.
latifolius. Borley wood, on the South-
west side: in the woody part of the
Devil's Ditch, near Canvafs Hall. Id.
palustris. Little Eversden: Swaff-
ham Fen; by the side of Reche Load,
about 100 yards before the Cut divides. Id.
VICIA sylvatica. Hall Wood, Wood Ditton.
Rev. J. Hemsted.
ASTRAGALUS glycyphyllos. Pastures and hedges
at Madingley; Trumpington; Shelford;
Hinton. Relban.
hypoglottis. Gogmagog Hills:
Hildersham: Barrington Hill, near
Linton: Newmarket Heath: Linton:
Teversham. Id.

TRIFOLIUM ochroleucum. Gravel pits near Gran-
chester Meadow: Comberton: Hin-
ton : Chalk-pit Close, &c. Relhan.
scabrum. Near Trinity Conduit
Head: Hill of Health: Newmarket:
Chippenham : Gamlingay. Id.
MEDICAGO falca:a. Quey: Chippenham Gravel-
pit: Boarn-Bridge: Bartlow: Wilbra-
ham: Linton. Id.
polymorpha. β. Chippenham Gravel-pit.
Relhan. Near Newmarket. Rev. J. Hemsted.
Sonchus palustris. Near Stretham Ferry, (Not
found there for many years). Relban.
LACTUCA scariola. Hazenfield, in the lane lead-
ing into the Village from Cambridge:
between Histon and Bampton: In a lane
leading from Long Stanton towards
Swavescy: Burwell Pit: Isle of Ely.
Relban. Near Denny Abbey, between
Cambridge and Ely. Rev. J. Hemsted.
Town in the road to Cottenham: on the
bank a little way below Clayhithe. Relhan.
LEONTODON palustre. On Hinton, and Tever-
fham Moors. Id.
CREPIS sætida. Between Little Shelford, and
Whittlesford, ' Id.

CREPIS liennis. In the Horse Road from Bottis-	
ham to Newmarket : Linton.	Relhan.
HYOSERIS minima. Gamlingay, near the Wind-	
mills, and the bogs: in a field adjoining	
to White Wood.	Id.
HYPOCHÆRIS maculata. Gogmagog Hills: be-	
tween Triplow Heath and Foulmire:	
Hildersham : Newmarket Heath : Relban.	
In the Devil's Ditch, near the Turnpike	
Gate. Rev. J. H	cmsted.
glabra. Gamlingay, under the	
pales of the Park, between the Town and	
Sir J. Downing's Gateway.	Relhan.
CARDUUS eriophorus. Madingley: Hinton: Chalk-	
Pit Close: Kingston Wood, &c.	Id.
pratensis. Hinton Moor: Fulbourn:	
Anglesey Abbey: Littleport: Bardwell	
· Fen.	Id.
GNAPHALIUM luteo-album. Between Hauxton	
and Little Shelford, by the road side;	
and in a gravel-pit, on the right hand	
side of the road, leading towards Little	
Shelford.	Id.
dioicum. Gogmagog Hills : New-	
market Heath.	Id.
rectum. Gamlingay Park.	Id.
SENECIO viscosus. Sandy Grounds at Gamlingay;	
Mepole : Chatteress.	Id.
*	

SENECIO paludosus. Stretham Ferry; and in the	
Isle of Ely. Ray. (Not found there for	
many years. Relhan.)	
INULA Helenium. Madingley, in a Close near the	
road leading to Drayton: Eversden	
wood : Lolworth, in a Close N. W. of	
	Rethan
CINERARIA palustris. March : Chatteress : Chip-	
penham, in a Ditch, on the edge of the	
Moor, next to the Park.	Id.
integrifolia. Gogmagog Hills : New-	
market Heath.	Id.
ORCHIS ustulata. Gogmagog Hills: Devil's	
Ditch : Chippenham.	Id.
SATYRIUM wiride. King's Hedges: Hinton, in	
a Pit near Chalk-Pit Close: Wood-	
Ditton: Catlidge.	Id.
OPHRYS Nidus-avis. Madingley Wood: Hall	
Wood.	Id.
Loeselii. Hinton, Teversham, Ful-	
bourn, and Sawston Moors.	Id.
monorchis. Chalk-Pit Close: pits be-	
tween Hinton and the road to Gogma-	
gog Hills: Westhoe, in an old gravel-	
pit near Mr. Keene's House.	Id.
anthropophora. In a Close near Linton.	Id.
muscifera. Hinton : Teversham : Ful-	
bourn : Devil's Ditch : Chippenham :	
Linton: Langley Wood,	Id.

OPHRYS apifera. Madingley Wood: Hinton:
Teversham: Fulbourn: Burrough-Green:
Chippenham: Linton: near Hall Wood:
Catlidge. Relban.
aranifera. Chalky Pastures at Shelford ;
Abingdon; Hildersham; Bartlow. Id.
MALAXIS paludosa. Gamlingay Bogs. Relhan.
On Hinton Moor. Dr. Manningham.
SERAPIAS longifelia. Hinton, Teversham, and
Sawston Moors: Triplow Heath, &c. Id.
grandiflora. Quey: Isle of Ely. Ray.
(Not found for many years. Relhan.)
ARISTOLOCHIA Clematitis. Woods and Hedges
at Milton, and Whittlesford. Id.
Турна angustifolia. Brick-kilns on the Chester-
ton road: sides of the Cam, below
Ditton: Swan Pond, &c. Id.
SPARGANIUM natans. Stream by the edge of
Sawston Moor: Wilbraham Moor:
Burwell Fen. Id.
CAREX dioica. Hinton Moor: Gamlingay Bogs:
Moor between Snailwell and Exning. Id.
B. Fl. Cant. (C. capitata. Huds.)
Hinton Moor. Relhan.
curta. Gamlingay Bogs. Id.
axillaris. Hall Wood, Wood-Ditton. Id.
- teretiuscula. Among some low bushes

in Granchester Meadow: Fulbourn,
near the footway to the Mill. Relhan,
CAREX pendula. Wood-Ditton: Catlidge. Id.
* Oederi. In a ditch which crosses the
foot-path to Stapleford near the Nine
Wells: Bottisham Load. Id.
- distans. Watery Places and Meadows. Id.
Pseudo-cyperus. Fulbourn, by the foot-
path to the Mill: Pampisford: Angle-
sey Abbey: Burwell: Wood-Ditton:
Catlidge. Id.
vesicaria. Paradise, near Granchester
Meadow. Rev. Mr. Newton. Wis-
bech, in a field near the New Common
Bridge. Mr. Skrimshire.
ampullacea. In the ditch by the road to
Hinton, on the edge of the Fen : Foul-
mire Common: Chippenham Moor.
Ely: Isle of Ely: Gamlingay Bogs. Relhan.
biner vis. Hinton Moor: Chesterton near
Cambridge. Real D. D. H
LITTORELLA lacustris Comit D
* This is Carex extensa Fl. Gant. as I know by specimens which

Mr. Relhan has been so obliging as to shew and give mc. D. T. VOL. 1.

AMARANTHUS Blitum. Near Parker's Piece. Relhan.
MYRIOPHYLLUM verticillatum. In a Ditch near
Coldham Common, by the footway to
Teversham: Quey. Id.
SALIX Russelliana. By the first mile stone from
Cambridge, on the London Road. Id.
vitellina. Queen's College Grove, Cam-
bridge. Id.
rubra. Near Ely. Rev. Dr. Goodenough.
Prickwillow, in an Ofier Holt on the
right hand side of the river, opposite to a
public house, two miles from Ely.
Rev. J. Hemsted.
Ofier Holts by the river side, a
little above Cambridge, plentifully. Rev. R. Relhan.
amygdalina. Banks of rivers and ditches. Relhan.
fissa. Prickwillow, near Ely. Rev. J. Hemsted.
ATRIPLEX pedunculata. Wisbech, by the river
side, a little way below the town; and
in the salt marshes. Mr. Skrimsbire.
EQUISETUM sylvaticum. Madingley Wood : Ches-
terton, &c. Relhan.
fluviatile. Banks of rivers and sides
of ditches, &c. Id.
byemale. Stretham Ferry: Gamlin-
gay Bogs. Id.
OSMUNDA Lunaria. 1n Little Linton Warren:
Chippenham, gravel-pits on the outside

of the Park, near the Gate: near Balsham. Relban. OSMUNDA regalis. Gamlingay Park, on both sides of the pales in the corner next to Sandy. Ray. (Not found since his time. Relban.) LYCOPODIUM inundatum. Gamlingay. Id. ASPIDIUM Lonchitis. Gamlingay. Ray. (Not found since his time. Relban.) ------ Thelypteris. Fulbourn, by the footpath to the Mill, on both sides: Fulbourn, and Teversham Moors : Gamlingay. Id. PHASCUM serratum. Newmarket. Rev. J. Hemsted. _____ curvicollum. In the small plantation on Gogmagog Hills, by the road to Babraham. Id. rectum. Graffy places at Newmarket. Rev. J. Hemsted. GRIMMIA calcarea. In a chalk-pit near Gogmagog Hills, and near Newmarket. Mr. Dickson. ENCALYFTA lanceolata. Mole Hills between Hinton and Teversham. Relhan. TORTULA rigida. Near Newmarket, in the road to Wood Ditton. Rev. J. Hemsted. Wisbech. Mr. Skrimsbire. ---- tortuosa. Gravel Hill: Gogmagog Hills: Newmarket Heath. Relban, F 2

(It never bears fruit in this county. Mr. Hemsted.)	
,	Relhan.
BRYUM aureum. Hot-house in the Botanic Gar-	20010 0000-
den.	Id.
	7
JUNGERMANNIA sphærocephala. Wisbech, on the brinks of ponds. Mr. Skri	mehire
brinks of policies	11131351 6 0
pusilla. Gamlingay Heath, near	Relhan.
SPHÆROCARPUS terrestris. Barnwell Gravel-pit.	100
MARCHANTIA hemispharica. Hinton, Fulbourn,	73
and Sawston Moors.	Id.
cruciata, Sawston Moor.	Id.
conica. Hinton Moor.	Id.
RICCIA natans. Sawston Moor: in a pond in a	
grove of oaks near Hall-wood, Wood-	
Ditton.	Id.
LICHEN roseus. Ach. On the inside of decayed	
willows behind Granchester Mill : Haux-	
ton.	Id.
Hoffmanni. Stones on Gogmagog Hills.	Id.
coardatus. On the Garden Wall between	
the Pound Hills and the Huntingdon	
turnpike.	Id,
conspurcatus. On the Churches at Hinton;	
Teversham; Hauxton; Swavesey: and	
on old chalk walls, not uncommon.	Id.

LICHEN Turneri. Madingley Wood.	Relbano
Ebrartianus. Old pales at Madingley,	
&c.	Id.
calvus. Stones on Gogmagog Hills:	
Gamlingay, in a gravel-pit near White	
Wood.	Id.
cretaceus. Pieces of Chalk on Gogmagog	
Hills.	Id.
cinereo-fuscus. Trunks of trees and old	
pales.	Id.
cæsio-rufus. Walls in Deadman's-lane,	
Wisbech. Mr. Skr	imshire.
crenulatus. On the walls of St. Peter's	
	Relhan.
pezizoides. On the wall of Gamlingay	
Park, a little beyond the Great Gate-way.	Id.
impolitus. On an Elm Tree in Hinton.	
corticola. Madingley Wood.	Id.
epipolius. On the walls of Christ, and	
St. Peter's Colleges: Hinton Church,	
&c.	Id.
sanguinarius. On the Pales of Madingley	
Park, not far from the Churchyard.	Id.
inquinans. Madingley Park: West	
Wratting.	Id.
bypnorum. Gogmagog Hills, on the	
newly ploughed up land.	Id.

LICHEN caruleo-nigricans. Gogmagog Hills : near	•
Hare Park : Newmarket Heath.	Relhan.
scalaris. Madingley Park Pales, near the	
entrance from the Caxton road.	Id.
decipiens. Gogmagog Hills : Newmarket	:
Heath.	Id.
subimbricatus. On Clare Hall Bridge:	
on the walls of Granchester, and Imping-	
ton church-yards, &c.	Id.
lentigerus. Gogmagog Hills : Newmarket	
Heath.	Id.
granulatus. Moist sandy places.	Id.
fascicularis. Hinton, on old Elms near	
the spring: Gogmagog Hills: New-	
niarket Heath.	Id.
trapeziformis. Gravel-pit near Hare Park	. Id.
pulmonarius. White Wood, near Gamlin-	
gay.	Id.
polydaclylos. Dry sandy heaths.	Id.
jubatus. On Gamlingay Park pales, near	
the Public House, by White Wood.	Id.
Fucus amphibius. Tydd Marsh. Mr. Skr.	imshire.
ULVA pruniformis. Madingley Bath : Hinton	
Moor, in the springs and the ditches.	Relhan.
CONFERVA confragosa. River Cam: ditches in	
Garrét Hostel Lane.	Id.
fucicola. Wisbech, on Fucus vesicu-	
losus. Mr. Skr.	imshire.

CARDIGANSHIRE.

AVENA fatua. At Hafod. Dr. Smith. ELYMUS arenarius. In a flat salt Marsh about five miles from Aberystwith towards Dolgelle. Mr. A. Aikin. RUBIA peregrina. Near the Devil's Bridge. Mr. 7. Woods, jun. PULMONARIA maritima. About Aberystwith. Evans. SCANDIX odorata. Behind a farm house, about a mile West of Hafod. Dr. Smith. SCILLA verna. About Aberystwith. Evans. Juncus Forsteri. In a wood opposite the South front of Hafod House. Dr. Smith. SILENE Otites. Marshy banks of the Rhyddol near Aberystwith. Mr. A. Aikin. (Surely from the situation this must have been some other species.)

EUPHORBIA Peplis. Neighbourhood of Aberyst-
with. Evans. I have received a speci-
men from the shore at Aberystwith.
Mr. T. F. Forster, jun.
Spir ÆA salicifolia. In a wood at Hafod, near a Gate,
in the Eastern approach to the House. Dr. Smith.
OROBUS sylvaticus. At Hafod. Mr. Todd. (Dr.
Smith mentions in Flor. Brit. III. p.
1403, that what he takes for a singular
variety of this plant, with large simple .
leaves, but never bearing flowers, has also
been found near Hafod by Mr. Todd.)
HYPERICUM dubium. At Hafod, plentifully. Dr. Smith.
GNAPHALIUM dioicum. On Plinlimmon. Merrett.
SATYRIUM albidum. On the expofed grassy hills,
which furround Hafod. Miss Johnes.
β. At Hafod. Mr. Todd.
Aspidium Oreopteris. 7
DICRANUM polyphyllum. } At Hafod. Dr. Smith.
BUXBAUMIA foliosa. On moist rocks near the
River, below the House at Hafod. Id.
LICHEN plumbeus. Trunks of trees in this County. Id.
Burgessii. Trunks of trees about the
Devil's Bridge, and at Hafod. Id.
perlatus. Found in fruit in this County by Id.
glomuliferus. Very abundant in Col.
Johnes' woods at Hafod. Id.

LICHEN aquaticus.	Bed of an alpine river at Ha-	
fod.	Dr.	Smith.
fuliginosus.	2-	
limbatus.	Rocks and trees at Hafod.	Id.
inclusus. P	lentiful at Hafod, on trees	
upon the	Hill immediately before the	
	nt of the House.	Id.

CARMARTHENSHIRE.

ERIOPHORUM vaginatum. Frequent in moun-	
tainous Bogs. L. W.	D.
CAMPANULA hederacea. Shady places near Car-	
marthen, and other parts of the County,	
not unfrequent. Ia	<i>!</i> .
VIOLA lutea. On the Black Mountains. Mer	rett.
RIBES alpinum. It propagates itself in the Woods	
at Golden Grove, but I doubt its being	
indigenous. Mr. M	ilne.
SIUM verticillatum. Moist Meadows, plentifully.	
Sir J. Ba	
JUNCUS acutus. Sandy Meadows about Kidwelly. R	lay.
VACCINIUM Oxycoccos. On the Mountains, fre-	
quent. L. W.	D.
LYTHRUM hyssopifolium. On the road side near	
Gelli towards Carmarthen. Eve	7115.
ANTIRRHINUM repens. About Llandovery and	
Llangaddock, plentifully. L. W.	D.

.

 HYPERICUM Androsamum. Several places near

 Carmarthen.
 L. W. D.

 SALIX argentea. In the Sand Burrows at Laugharn.

 harn.
 Mr. Hurlock.

 EMPETRUM nigrum. Mountainous parts of the

 County, frequent.
 L. W. D.

 SCOLOPENDRIUM Ceterach. Frequent in this

 County.
 Id.

 DICRANUM aciculare.
 Rocks near Pontardy

 polyphyllum.
 Iais.
 Id.

CARNARVONSHIRE, in a botanical point of view, may certainly be considered the best cultivated, as well as the most interesting of the Welsh Counties. Its Mountains surpassing in height every other in the Principality, afford inexhaustible fields for discovery; and from the days of Dillenius, when they were investigated by that able Botanist and his Friends, to the present, have never ceased to attract the attention of travellers of taste and science. For the greater part, however, of what we know respecting their Botany, especially in the Class Cryptogamia, we are indebted to two Naturalists of our own time, Rev. Hugh Davies and John Wynne Griffith, Esq.the latter of whom in particular, joining to a most acute mind an ardor that cannot be surpassed, continues to explore the Botany of this and the neighbouring Counties; and it is to be hoped, will shortly favor the public with observations upon the British Lichens, which his situation and knowledge so well enable him to illustrate.

VERONICA spicata. Woods near Gloddaeth. Rev. H. Davies. On Penmaen Mawr. Bingley. Side of Llandidno Rocks. Mr. Griffith. ----- hybrida. Hyssva Bengam, and Trygyvylchi Rocks near the Glyder. Evans. SCHOENUS albus. Bog West of Dolbadarn Castle, near Llanberris. Mr. Griffith. Peat Bogs among the hills, not rare. Rev. H. Davies. ERIOPHORUM vaginatum. Mountainous Peat Bogs throughout the County. Rev. H. Davies. Turbaries near Llyn Idwel. Bingley. ----- polystachion. With the former, but not so common. Id. POA alpina. Rocks of Snowdon; where it varies with * viviparous flowers. Rev. H. Davies. --- glauca. High Mountains about Llyn Idwel, and Llanberris; and on Clogwyn y Garnedd, one side of Snowdon. Bingley. FESTUCA vivipara. On Snowdon. Rev. H. Davies. About Beddgelart; Nant Francon; and the mountainous parts of the County, abundantly. D. T. About Capel Cerrig, plentiful. Mr. Grifhth.

* Mr. Griffith observes that the viviparous plant is certainly a distinct species, as he has cultivated it more than 18 years, during which it has been constantly viviparous. I have seen it in the Cambridge Botanic Garden, still more viviparous than on the Welch Mountains. D. T.

FESSUCA rubra. B. (F. Cambrica, Huds.) On
the ledges of Clogwyn y Garnedd. Herb.
Banks. Crib y Ddescil. Mr. Griffith.
bromoides. Common in waste places
round Bangor. D. T.
AVENA fatua. Very Common in this County. Mr. Griffith.
ARUNDO colorata. B. Near Llanberris. Ray.
y. Bank of the Rivulet, that
runs by Dolbadarn Castle into the Lake
in the vale of Llanberris. Bingley.
ELYMUS arenarius. Near the Orme's Head. Mr. A. Aikin.
GALIUM boreale. Trigiwulchau Rocks: also
about half a mile from Llanberris, in the
gravel by the side of the second Rivulet
in the way to Llanrwst; and on Clog-
wyn y Garnedd. Mr. Griffub. Rocks
near Twll Dû above Llyn Idwel. Bing-
ley. Sides of Snowdon. Rev. H. Davies.
RUBIA peregrina. Gloddaeth. Pennant. Llan-
didno Rocks; and in the fissures of the
Rocks on the North side of Penmaen
Mawr. Mr. Griffith.
RUPPIA maritima. Ditches between Traeth Mawr
and Pont Aberglaslyn. Bingley.
PULMONARIA maritima. Shore by the Orme's
Head. Mr. A. Aikin. By the river
Llyfni, that runs from Llyniau Nantle
into the sea, about half way between
Llandwrog and Clynog. Bingley.

ASPERUGO procumbens. North side of Llandidno Rocks, as you descend down to the Llêch; in a most perilous situation, and certainly wild. Mr. Griffith.

CAMPANULA rotundifolia. var. flor. unico. Common on the higher parts of all the Mountains about Llanberris and Snowdon. Bingley.

hederacea. In the wood, called Pancretch between Llanrwst & Capel Cerrig. *Ray.* Swampy places in the meadows of the Vale of Llanberris, about a quarter of a mile beyond the Village : by the road side near Llanrwst; and near the Cataract Rhaiadr y Wenol, 5 miles from Llanrwst. *Mr. Griffith.* Meadows below Carnedd Llewelyn. *Rev. H. Davies.*

LOBELIA dortmanna. Llyn y Cwn, one of the Lakes of Snowdon. Pennant. Plentifully in the river Seiont, between the lower Pool of Llanberris and Cwm y Clo. Bingley. Upper Lake at Llanberris: Llyn Ogwen: Ffynnon frêch, and

* This is a distinct species from C. rotundifolia, as I have raised it from seed. Mr. Griffith.

most of the mountainous Lakes in Wales. Mr Griffith. VIOLA lutea. By the road side from Llanrwst to Festiniog, near the bridge about a mile from Penmachno. Bingley. RIBES alpinum. Hedges between Bangor Ferry and Bangor. Rev. H. Davies. GENTIANA campestris. About Llanberris. Hudson. Very common in most mountainous pastures in Wales. Mr. Griffith. SIUM verticillatum. In a meadow immediately on the entrance of Nant Hwynant Vale from Beddgelart. Rev. H. Davies. PIMPINELLA dioica. Llandidno Rocks. Mr. Griffith. OENANTHE pimpinelloides. Salt marsh near Aber. Rev. H. Davies. DROSERA longifolia. Between Beddgelart and the Salmon Leap. Id. SCILLA verna. Gloddaeth. Pennant. Llandidno Rocks; and on a Promontory between Llandûlas and Groes yn Eirios. Mr. Griffith. autumnalis. Llandwyn. Ray. ANTHERICUM serotinum. On Snowdon; and on the Mountains in the neighbourhood of Llanberris : on the west side of Trigvilchau. Ray. Abundant on rocks above Twll dû: I have alfo found it in one place only on the rocks of Crib y Ddescil, VOL. I.

but could never find it on Clogwyn ddu'r Arddû. Mr. Griffith. On Clogwyn ddû'r Rev. H. Davies. Arddû. Id. JUNCUS acutus. Morfa-menog, Llanengan. ----- triglumis. Near Llyn y Cwn on Glyder Vawr. Pennant. In the afcent from Llanberris to Glyder, but only in one small spot, that lies between a small eminence called Bryn Brâs, and a rivulet cal-Mr. Griffith. ed Avon Lâs. _____ spicatus ? Cwm Idwel; and Iscolion duon. Id. RUMEX digynus. Cwm Idwel: Clogwyn y Garnedd: and Crib y Ddescil. Mr. Griffith. Moist rocks near Llyn Fynnon Las. D. T. ALISMA natans. In the great Lake below the old Castle at Llanberris. Brewer. In the Rivulet that runs past the Castle into the Mr. Griffith. Lake. EPILOBIUM angustifolium. Among the Bushes in several of the meadows in Cwm y Clo, at the bottom of the Lower Llanberris Bingley. Lake. * angustissimum? Rocks near Twll Mr. Griffith. Dû in Cwm Idwel.

• " I have fubjoined a mark of doubt to this plant, never having found it in flower, but, as I have been informed by my Friend, Dr. Smith, that *E. angustissimum* grows in similar situations abroad, this Note may be productive of further investigation upon the sub-

83

CHLORA perfoliata. Near the Orme's Head. Ewans. VACCINIUM uliginosum. Moist high woods about Gwydir, near Llanrwst. Bingley. - Vitis idea. Snowdon and other Mountains. Rev. H. Davies. Top of the Glyder; also Rocks about Llyn y Cwn. Mr. Griffith. .---- Oxycoccos. Peat Bogs at Llyn. Rev. H. Davies. Bogs, very common. Mr. Griffith. DAPHNE Cneorum. Said to have been found by Mr. Meyrick, a Surgeon near Birmingham, about two miles from Beddgelart, by the road side leading to Carnarvon, just at the place where the ascent to Snowdon begins. It has since been sought for in vain by many Botanists. Bingley. POLYGONUM viviparum. Rocks above Ffynnon frêch. Mr. Griffith. _____β. Mountainous pastures on the rock Crib y Coch, above Ffynnon

ject. Until I faw Dr. Smith, I always considered it as a starved wariety of *E. angustifolium*, but now I think it very likely to prove *E.* angustissimum. This is the *E. angustifolium* of *Bingley's Tour*, which he mentions to grow on the rocks of Hysfa Bengam; but he is wrong in placing those Rocks between Llanberris and Cwm Idwel, for they are in fact the upper part of Cwm Idwel, close to Twll Dû." Mr. Griffub.

frêch, by Llanberris. Lhwyd. SAXIFRAGA stellaris. On Snowdon. Ray. In the neighbourhood, very common. Mr. Griffith. Near the foot of the waterfall at Aber. Bingley. Nant Francon, abundantly. D. T.

SAXIFRAGA nivalis. On the rocks of Clogwyn y
Garnedd: Crib y Ddescil: Clogwyn dû
Umhen y Glyder, &c. near Llanberris.
Lbwyd. On Clogwyn dû'r Arddû.
Bingley. Near the summit of Clogwyn
y Garnedd, which is by the top of
Snowdon, but very scarce: in Cwm Idwel, and Crib y Ddescil, but more rare.

Mr. Griffith.

oppositifolia. With the former, in abundance. Mr. Griffith. Among the rocks about Twll dû; and Glyder: on Glogwyn y Garnedd; and Clogwyn dû'r Arddû. Bingley.

- caspitosa. Twll dû, Cwm Idwel.

Mr. Griffith.

palmata. Rocks of Cwm Idwel. *Id. hypnoides.* Frequent on the mountains. *Ray.* (The shape of the Petals by no means forms a sufficient character for the distinguishing of this plant, as in some

specimens which I found on Snowdon,
near Llyn Ffynnon Lâs, instead of being
elliptico-oblonga, they were late ovata
ct ferè subrotunda. D. T.)
SAXIFRAGA ajugifolia? Cwm Idwel; and Cwm
y ffynnon Lâs. Mr. Griffith.
SILENE anglica. In a cornfield on the Coast, not
far from Llanfaglan Church, 3 miles from
Carnarvon. Bingley.
nutans. Near Gloddaeth. Rev. H. Davies.
acaulis. On the steep and higher rocks of
Snowdon, every where. Lbwyd. Cwm
Idwel. Mr. Griffith.
* ARENARIA verna. On Snowdon. Id.
β. (A. juniperina. With.)
Near Llanberris. Mr. Griffith. Rocks
near Twll dû, and on Clogwyn y Garnedd.
Bingley.
Near Twill da with the former Die I
Near Twll dû, with the former. Bingley. On Snowdon. Mr. Griffith
SEDUM dasyphyllum. On the wall as you enter Con-
way Church Yard, by the Eastern Gate. Id.
different fait, by the Eastern Gate. Id.

* "I am of opinion that the Snowdon Arenaria is perfectly distinct from the Holywell plant, but which of the two is the true A. *:rna I leave others to determine." Mr. Griffith.

SEDUM rupestre. North side of Penmaen Mawr,
among the loose stones above the Turn-
pike Road, very near the inaccessible
Rocks : among the Rocks near Twll Dû,
called Creigiau Hyssva Bengam : and on
a wall south of Gwydir Chapel, near
Llanrwst. Mr. Griffith.
CERASTIUM alpinum. Sides of Snowdon, plenti-
fully. Ray. Clogwyn dû 'r Arddû.
Rev. H. Davies. Highest Rocks of Snow-
<i>Latifolium.</i> With the former. Mr.
Griffith. On Clogwyn y Garnedd.
Dr. Richardson.
Spergula subulata. Not uncommon on the
rocks or this County. Rev. H. Davies.
EUPHOREIA Portlandica. On the Coast near Car.
narvon. Hudson.
Pyrus Aria. The North End of the Great Orme's
Head: on the Rocks of Penmaen Mawr
Head: on the Rocks of Penmaen Mawr above the Turnpike Road. <i>Bingley</i> . Near Gloddaeth, and Penrhyn. <i>Rev. H. Davies</i> .
Head: on the Rocks of Penmaen Mawr above the Turnpike Road. <i>Bingley</i> . Near Gloddaeth, and Penrhyn. <i>Rev. H. Davies</i> .
Head: on the Rocks of Penmaen Mawr above the Turnpike Road. Bingley. Near
Head: on the Rocks of Penmaen Mawr above the Turnpike Road. Bingley. Near Gloddaeth, and Penrhyn. Rev. H. Davies. Rosa villosa. Very common all over Wales.
Head: on the Rocks of Penmaen Mawr above the Turnpike Road. Bingley. Near Gloddaeth, and Penrhyn. Rev. H. Davies. Nery common all over Wales. Mr. Griffith.

loose stones about the Great Orme's Head. Bingley. RUBUS chamemorus. In Peat Bogs on the sides of the mountains. Hudson. POTENTILLA verna. Llandidno rocks. Mr. Griffith. At Gloddaeth. Pennant. TORMENTILLA reptans. Sandy places about Carnarvon. Bingley. GEUM rivale. Moist spots among the Hills. Rev. H. Davies. On Snowdon. Ray. PAPAVER Cambricum. Near the Bridge at Aber, in the Bed of the River. Lbwyd. On the Back of Snowdon, going from Carnarvon to Llanderris, not far from the Castle. - Ray. CISTUS marifolius. Gloddaeth. Pennant. South side of Llandidno Rocks, abundantly. Mr. Griffith. THALICTRUM alpinum. Very frequent on the mountains. Lhwyd. Rocks about Llyn y Cwn: Twll dû: and on Clogwyn y Garnedd. Bingley. Abundant near the summit of Snowdon. D. T. ____ minus. Frequent in the County. Hudson. About Cwm Idwel; and Snowdon. Mr. Griffith. RANUNCULUS L'ngua. Common in wet Pastures about Carnarvon. Bingley .

RANUNCULUS * acris? On the higher rocks of Snowdon, particularly by the waterfall near Llyn Fynnon Lâs, abundantly. D. F. TROLLIUS curopæus. Meadow below Penrhyn. Rev. H. Davis. In the hollow, immediately below the Cataract in Caunant Mawr: near Dolbadarn Castle: in the Vale of Llanberris; and in the Meadows near Llyn Cowlid, a lake in the mountains above, and nearly North of Capel Cerig. Bingley. Rocks in Cwm Idwel; Clogwyn dû'r Arddû; and Crib y Ddescil. Mr. Griffith. AJUGA alpina. On the flat near the fummit of Carnedd Llewelin. Id. MENTHA odorata. In the upper pool, and ditches about Capel Cerig. Id. GALEOPSIS villosa. About Bangor. Hudson. Cornfields between Bangor and Carnar-

* This Ranunculus seems generally to have been taken by Botanists for an alpine variety of *R. acris*, but appears to me sufficiently distinct, and nearly allied to *R. Gouani*, of Willdenow: the whole plant is not 3 inches high: the radical leaves on long petioli, shaped like those of *R. Acris*, but with wider segments; the cauline ones narrow, palmate, sessile; the stem undivided, bearing only a single flower.' A root, which I brought into my garden is in the first year quite altered, and has produced branching stems 2 or 3 feet high; but is still less like *R. acris*: in so difficult a genus, however, I do not venture to make a new species. *D. T.*

von: and on the beach between Aber and Ogwen rivers, plentifully. Rev. H. Davies. SCUTELLARIA minor. Swampy Meadows and Pastures about Carnarvon and Llanberris. Bingley. SCROPHULARIA vernalis. Gloddaeth. Sir 7. Cullum. OROBANCHE minor. Among the ruins of Diganwy Castle; near Conway; and in two or three places among those of Conway Castle. Bingley. SUBULARIA aquatica. Llyn Ogwen; and all the Lakes about Snowdon. Mr. Griffith. Llyn y Cwn. Pennant. DRABA incana. Sides of Snowdon. Rev. H. Davies. Crib y Ddescil and Cwm Idwel. Mr. Griffith. On the high rocks East of Twll Dû, above Llyn Idwel, called Creigiau Hyssva Bengam. Bingley. Frequent in the County. Hudson. LEPIDIUM petraum. Walls at Gloddaeth. Rev. H. Davies. COCHLEARIA officinalis. On the Sea Coast, near the Great Orme's Head : and on the rocks of Clogwyn y Garnedd. Bingley. - y. (C. groenlandica. With.) Plentifully in the mountains about Llanberris. Hudson. ---- danica. On a wall by the side of the hill between Carnarvon and Pont Sciont. Bingley. Stone walls near the old Mill

at Llandrillo; and Llandidno Bay. Mr. Griffith. Abundant on the rocks by the Menai near Bangor Ferry. D. T. CRAMBE maritima. On the Coast in various parts of the Promontory of Llyn. Bingley: Beach near Crickaeth, plentifully. Rev. H. Davies. In the most inaccessible Rocks of the lesser Orme's Head, near Conway, facing North. Mr. Griffith. SISYMBRIUM menense. In the Sand of the Sea Coast near the Great Orme's Head. Bingley. CHEIRANTHUS sinuatus. Morfa Bychan, near Penmorfa. Rev. H. Davies. At Aberdaren. Ray. ARABIS hispida. On Clogwyn dù 'r Arddû; and Clogwyn y Garnedd. Lbwyd. Ascent of Snowdon from Cwm Brynog. Rev. H. Davies. I never found this in the neighbourhood of Snowdon, except on the Rocks of Clogwyn dû 'r Arddû, which are synonymous to those above Lake Layn-du, meaning Llyn dû. Mr. Griffith. BRASSICA oleracea. Aberdaron, in the Promontory of Llyn. Bingley. On the perpendicular Rocks at the extremity of the Great Orme's Head. Rev. H. Davies. ERODIUM maritimum. Very abundant about Carnarvon. Ray. Every where, on stone fences about Bangor. D.T.

GERANIUM sanguineum. At Gloddaeth. Pennant.	
Llandidno Rocks. Mr. Griffith. Near	
the Orme's Head. Mr. A.	Aikin.
OROBUS sylvaticus. Side of the River near the	
	Griffith.
LATHYRUS sylvestris. Hedge by the Beach just	
below Aber. Rev. H. Davies. About	
0	Audson.
latifolius. Near Gyffin Mill, about	
half il c a	Singley.
palustris. Moist pastures near Beddge-	ungiey.
lart.	Id.
HYPERICUM androsæmum. In a hedge near the	2000
Menai, between Carnarvon and Llanfair	
Iscaer; and in the woods at Glangconna,	
Door Con	Id.
the TT I I TT I	Id.
LEONTODON palustre. Boggy Ground about	A
LEONTODON palustre. Boggy Ground about HEDYPNOIS Taraxaci. Llyn y Cwn. Mr. Gr	i Æ. L
HIERACIUM alpinum. Rocks West of Llyn y Cwn.	Jup;
Mr. Griffith. Rocks about Cwm Idwel.	
Rev. H. Davies. On the Glyder, and	
Tricfulate TI	udson.
sylvaticum. Sides of Llyn y Cwn.	643 678a
Dr. Richardson. (As this plant is quite	
excluded from the British List in the	
3d Vol. of Flora Britannica, it would be	

curious to ascertain what species was here designed by Dr. Richardson.) HIERACIUM * paludosum. Cwm Idwel; and rocks of Iscolion duon. Mr. Griffith. Meadows in Nant Hwynant Vale near D.T. Beddgelart. SERRATULA alpina. On the highest rocks of Snowdonia, as Clogwyn y Garnedd; Iscolion dûon. Ray. Rocks of Crib y Ddescil and Cwm Idwel. Mr. Griffith. $----\beta$. Near a rivulet in the rock Rhiw 'r Glyder, above Llyn y Cwn. Ray. CARDUUS tenuiflorus. Dry bank sides near Carnarvon Castle. Bingley. ---- heterophyllus. In the ascent from Llanberris to Glyder. Bingley. As you ascend from Llyn y lâs to Crib y Ddescil. Mr. Griffith. GNAPHALIUM dioicum. Rocks in the upper parts of the clefts of Twll dû above Llyn Idwel. Bingley. Llandidno rocks; and rocks above Llyn y Cwn. Mr. Griffith. Heathy Commons. Rev. H. Davies. SOLIDAGO virga aurea. y. (S. cambrica. Huds.) About Llanberris, plentifully. Hudson. Pennant. Near Llyn y Cwn.

* Mr. Griffith observes, there are also two or three species of *Hieracium* about Snowdon which he has not yet settled.

INULA Helenium. Very plentiful in the fields and lanes about Aber. Waring.

SATYRIUM albidum. Roadside from Bangor to Llyn Ogwen. Rev. H. Davies. In moist meadows near Lord Penrhyn's flate quarries in Nant Francon: plentifully in Glyn Meadow between Cwm y Clo and Dolbadarn Castle; and near Llyn Cowlid in the meadows N. W. of Capel Cerig. Bingley. Rocky pastures near Bluen y Nant near Llyn Ogwen. Mr. Griffith. On the back of Snowdon on the way from Llanberris to Carnarvon. Ray. MALAXIS paludosa. In a field where a wood has formerly stood, called Coed y Tû Dû, near Llanberris. Mr. Griffith. SERAPIAS longifolia. Marshy Meadows in the County. Bingley.

SPARGANIUM natans. Lakes about Llanberris, plentifully. Hudson. Pools on the side of Snowdon. Sir 7. Banks.

CAREX dioica. In a bog at the upper end of Llyn Idwel. Mr. Griffith.

stellulata. I found in the meadows opposite the Inn at Beddgelart a curious variety of this plant, with a small male spike at the top of the androgynous ones.

My friend, Professor Mertens, had previoufly sent me a similar specimen from Germany. D.T. CAREX curta. At the foot of Snowdon, between Capel Cerig and Llanberris. Rev. W. Wood. pendula. Near the Beach at Bangor Rev. H. Davies. Ferry. atrata. Upper rocks of Crib y Ddescil. Mr. Griffith. Plentifully on the Mountains about Llanberris. Hudson. fulva. South end of Llyn Idwel. Mr. Griffith. binervis. On all the moors of Wales. Mr. Griffith. Very abundant in meadows opposite the Inn at Beddgelart. D. T. rigida. Rocks near the summit of Carnedd Llewelin. Mr. Griffith. On the top of Hudson. Snowdon. ampullacea. Bogs near Llyn Idwel. Bingley. LITTORELLA lacustris. Shores of all the alpine Mr. Griffith. Lakes in Wales. QUERCUS sessiliflora. Very common in Wales. Id. SALIX herbacea. Near the summit of Carnedd Llewelin; and about Bwlch Glas, as vou ascend Snowdon. Mr. Griffith. Sherard. Plentiful on Snowdon. reticulata. In the crevices of the rocks on the North side of Penmaen Mawr : also

:04

North side of Llandidno. Mr. Griffub. On the highest Mountains. Ray. EMPETRUM nigrum. On all the Moors in Wales. Mr. Griffith. Mountains above Aber. Rev. H. Davies. RHODIOLA rosea. On Snowdon. Ray. Rocks of Cwm Idwel, and Iscolion duon. Mr. Griffith. Abundant on the ridge of rocks immediately above Llyn Fynnon Lâs. D.T. JUNIPERUS communis. B. On the Rocks near Llyn y Cwn, above Llanberiis. Bingley. OSMUNDA lunaria. On Penmaen Mawr. Ray. regalis. Watery places about Pont Aberglasllyn. Bingley. Near Ystymllyn, abundantly. Rev. H. Davies. Lycopodium selaginoides. Rocks in Cwm Idwel; and Crib y Ddescil. Mr. Griffith. Not uncommon about the middle regions of Snowdon. D.T.annotinum. Plentiful on the Mountain Rhiw'r Glyder, above the Lake, Llyn y cwn; and not uncommon in the lower parts of the same mountain, opposite to the Rock, y Trivan. Ray. Found in great abundance about ten years ago near Llyn y Cwn, but never met with there since, though repeatedly sought for.

Mr. Griffith.

Lycopodium alpinum. Mcors, very common.
Mr. Griffith. On Snowdon. Ray.
POLYPODIUM vulgare. S. (P. cambricum. L.)
Gloddaeth. Bingley. Walls of Conway.
Rev. H. Davies.
arvonicum. On a moist black rock,
almost at the top of Clogwyn y Garnedd,
facing North West, directly above the
lower Lake. Dr. Richardson.
Phegopteris. In a moist dell at the
foot of Snowdon, near Llauberris.
Aikin. Clefts of Rocks in the higher
parts of Clogwyn y Garnedd. Bingley.
Cwm Idwel. Mr. Griffith.
Dryopteris. Cwm Idwel. Rev. H. Davies.
ASPIDIUM Lonchitis. Cwm Idwel. Mr. Griffith.
Plentiful on the Mountains above Llan-
berris. Hudson. On Snowdon, near
the foot of Crib y Ddescil; and on the
Glyder. Eolton.
Thelypteris. In a moist dell near the
village of Llanberris. Bingley.
Oreopteris. One of the most common
species in all mountainous pastures.
Rev. H. Davies.
Asplenium viride. On the moist rocks in the
clefts of Twll Dû: on Clogwyn du 'r y
Arddû; and on Clogwyn y Garnedd.

1 .

Bingley. Crib y Ddescil; and Cwm Idwel. Mr. Griffith. ASPLENIUM viride. B. Rocks in this County. Robson. marinum. By the well on the Llêch at Llandidno. Mr. Griffith. septentrionale. Carnedd Llewelyn. Lhouyd. SCOLOPENDRIUM Ceterach. Moist Rocks on the N.W. Coast. Bingley. Walls at Vainol. Rev. H. Davies. PTERIS crispa. Moist Rocks on Snowdon. Ray. Rocks about Llyn du yr Arddû. Mr. Griffith. At Lord Penryhn's Slate Quarries near Nant Francon. D.T. CYATHEA regia. Cwm Idwel. Mr. Griffith. On Snowdon, but very rare. Rev. H. Davies, HYMENOPHYLLUM tunbrigense. Moist Rocks on Snowdon. Rev. H. Davies. Cwm Idwel; and about Capel Cerrig. Mr. Griffith. Rocks under Dolbadarn Castle. Bolton, ISOETES lacustris. In Ffynnon frêch; and most of the other Lakes. Mr. Griffith. Llyn Ogwan; and Llyn y Cwn. Pennant. -- -- B. In Llyn Ogwan. Mr. Griffith, BUXBAUMIA foliosa. Near Llanberris. Dillenius. Cwm Cennog Rocks : also about Craig Dû, by the road side near Llanberris. Mr. Griffith. Abundant at Beddgelart, on the right hand side of the way to Pont Aberglâslyn. D, TYQL. I. H

GYMNOSTOMUM luteolum. Rocks of Clogwyn du	
yr Arddû: Crib y Ddescil; and Cwm	
Idwel. Mr. G	riffith.
fasciculare. Cwm Idwel:	
Crib y Ddescil; and near the summit	
of Snowdon.	Id.
Griffithianum. On the East side	
of Snowdon, about a hundred and fifty	
yards from the summit.	Id.
ANICTANGIUM lapponicum. Cwm Idwel.	Id.
ciliatum. Stone Fences and Rocks	
in Nant Francon, and every where in the	
County.	D.T.
ANDRÆA rupestris. On Snowdon; and the Gly-	
	lenius.
alpina. With the former, and near	
Llanberris. Dillenius. Very common	
on Rocks at Beddgelart.	$D. \mathcal{T}.$
SPLACHNUM minioides. Near Llyn Idwel, fre-	
quently on the dung of Foxes and	
	Griffith.
Turnerianum. About Llyn Idwel :	
also on the Bog by the Old Castle near	
Llanberris.	Id.
tenue. On Cow-dung about Cwm	
Idwel.	Id.
* sphæricum. On Cow-dung near	
the Old Castle at Llanberris.	Id.

* I very much doubt how far the true S. spharicum is really a native of our Islands: for all the specimens I have seen under that

ENCALYPTA ciliata.		
GRIMMIA recurvata.	Cwm Idwel.	Mr. Griffith.
striata. Crib y	Ddefcil; and Cwi	m Idwel. Id.
beteromalla. H	About Llyn Ogw	on 7.7
, ,)	At Beddgelart, by	the side
bomomalla.	of the Road fi	rom the
bomomalla. Donniana.	Inn to Pont Abo	erglâslyn. D. T.
acuta. Crib y	Ddescil. Mr.	Griffith.
On Snowdon.		Dillenius.
β. Mu.	sc. Hib. Near the	summit _
of Snowdon.		Mr. Griffith.
Para verticillata. Mari	ine Rocks near Ba	ingor. Brewer.
PTEROGONIUM gracile. A	bundant through	out the
County; but I f	ound it in fruit o	nly on
the moist rocks b	by the side of the	e river
at Pont Aberglâs	yn. (It was here,	, not in
that I gathered t	eously stated in	E. B.
that work.)	ine ipecimen ingu	
TRICHOSTOMUM capillace	um Cribu Ddaa	D. T.
colion dûon: an	d Cwm Idwel	11:1s-
Griffith. Llanbe		. H. Davies.
	- var. 2. With.]	
above Llyn Fynn	on Velan, abund	antly.
		Mr. Griffith.
ame have been S. sphæricum B.		

name have been S. sphæricum B. Musc. Hib. (Mr. Dickson's S. gracile) and the figure in English Botany undoubtedly represents that plant. Sec. t. 785. D. T.

TRICHOSTOMUM polyphyllum. Rocks about Gwydir Chapel. Mr. Griffith. Aber. Rev. H. Davies. Nant Francon; and every D. T.where about the Mountains. ---- var. with black capsules. On Snowdon. Dillenius. ----- fasciculare. microcarpon. > On Snowdon. Id. ____ obtusum. ericoides. Rocks of Clogwyn y Garnedd. Mr. Griffith. On Snowdon, D.T. not rare. lineare. Llanberris. Rev. H. Davies. ------- * fontinalioides. River at Beddge-D. T. lart. TORTULA tortuosa. Moist rocks at Nant Francon. Id. DICRANUM osmundaceum. Nant Francon, on moist rocks. Id. majus. Rocks by the Menai near Bangor Ferry, with G. verticillata. Id. _____ interruptum. On Snowdon. Mr. Griffith.

* I must beg leave to observe that in my opinion Dr. Smith has done wrong in making Fontinalis minor, Huds. and T. fontinalioides, Hedw. the same plant. 'The former is certainly Dillenius', T. 33. f. 2. the latter F. alpina Dicks. If therefore the name T. fontinalioides is to be preserved, Mr. Dickson's plant must be quoted as a synonym, and Dillenius', if not distinct, referred to var. β as in Musc-Hib. D. T.

DICRANUM patens. Rock above Gwydir Cha-
pel. Mr. Griffith. Rocks on Snowdon
washed by Llyn Fynnon Lâs. D. T.
ellipticum. In Cwm Idwel. Mr.
Griffith. Abundant on Snowdon, on
bare rocks at the Base of Moel y Wyddfa. D. 7.
aciculare. About Llanberris. Dil-
lenius. Abundant on Rocks near the
river at Beddgelart. D.T.
. B. Plentiful about Llan-
berris. Dillenius.
squarrosum. On the borders of Lakes
on Glyder and Snowdon. Id.
flexuosum. y. Abundant on Snowdon.
Dillenius. Llanberris. Rev. H. Da-
vies. At Nant Francon and Beddge-
lart, often covering very large pieces of
ground. D. T.
POLYTRICHUM strictum. Llanberris. Rev. H. Davies.
alpinum. On Snowdon. Dillenius.
urnigerum. Plentiful on Snow-
don, in higher regions than P. alpinum. D. T.
hercynicum. About Llanberris.
Rev. H. Davies. Not uncommon in
the higher parts of Snowdon. D. T.
NECKERA curtipendula. About Snowdon. Dil-
lenius. Plentiful every where in the

neighbourhood, but I never could find it D. T.in fruit. NECKERA crispa. Between Beddgelart, and Pont Aberglâslyn. Id. ----- heteromalla. Wood near Bangor. Id. BARTRAMIA Halleriana. Boggy ground in the afcent to Crib y Ddefcil from Llanberris; and about Llyn Idwel in the ascent to Twll dû, in great abundance. 6 Mr. Griffith. arcuata. On the Glyder. Dillenius. By the side of Avon Lâs, near Pystill, in Llanberris Parish. Mr. Griffith. Very abundant about Beddgelart. D. T. BRYUM nutans. About Llanberris, Rev. H. Davies. Cwm Idwel; and Crib y Ddescil. Mr. Griffith. elongatum. On Clogwyn y Garnedd. Herb. Dill. Abundant on Snowdon, where a small variety is also found. D. T. ____ aureum. On Snowdon. Mr. Wood. _____ var. 2. With. About Llanberris. Mr. Griffith. ----- crudum. Crib y Ddescil. Mr. Griffith. Rev. H. Davies. Llanberris. Zierii. Cwm Idwel : Crib y Ddefcil; and common on all the wet rocks about Snow-Mr. Griffith. don.

BRYUM alpinum. Rocks covered with a little earth, near the ruined Castle at Llanberris. Dillenius. About Llyn Ogwen; and Capel Cerig: also on rocks on Cwm Idwel; and Crib y Ddescil. Mr. Griffith. ----- * cubitale. Near Snowdon. Id. HYPNUM subtile. Under the Walls of the ruins of Segontium, near Carnarvon, not in Anglesea, as stated in Fl. Brit. Rev. H. Davies. ----- myosuroides. Very frequent about Snowdon. D.T. filamentosum. Crib y Ddescil; and Cwm Idwel. Mr. Griffith. ----- undulatum. On the Summit of Snowdon. Dillenins. ----- rufescens. Crib y Ddescil, wet rocks, but very rarely with capsules. Mr. Grifhth. ----- piliferum. Llanberris. Rev. H. Davies. ----- loreum. Very common about Beddgelart, and on heaths in the County. D. T._____ alpinum. Rocks on Snowdon, washed by Llyn Fynnon Lâs. Id.

* "That B. alpinum appears at first sight specifically distinct from B. cubitale I readily allow, but that it is so still I doubt. What two plants appear more distinct, or vary more in size than B. fragile, Dicks. and Sphagnum alpinum, Huds. yet they were not made a variety of Dic. flexuosum, nor do I know that they were suspected to be so, until the publication of my Note in Withering. 3rd. Ed. III. 834?" Mr. Griffith.

HYPNUM commutatum. Rocks by the Menai, near Bangor Ferry. D. T. FONTINALIS squamosa. Brooks near Llanberris. Dillenius. By the old Castle. Mr. Griffith. JUNGERMANNIA trilobata. On Crib y Ddescil. Hudson. - julacea. Moilt Rocks of Snowdon, and Glyder. Dillenius. guinquedentata. In the ascent to Ffynnon frêch, near Llanberris. Mr. Griffith. minuta. Summit of Carnedd Llewelin. Id. _____ spinulosa. Crib y Ddescil. Mr. Griffith. Snowdon. Dillenius. - purpurea. On Snowdon and Glyder. Id. ----- concinnata. On Snowdon. Mr. Griffith. curvifolia. Many places in the ascent to Crib y Ddescil from Llanberris : on the summit of Carnedd Llewelin. Id. pumila. Cwm Idwel. Id. LICHEN Oederi. Common about Beddgelart ; and on the very highest rocks of Moel y Wyddfa. D. T.------ silaceus. With the former, but most plentifully in Nant Hwynant Vale. Id. geographicus. About Bangor and Pentir. Mr. S. Brequer. flavo-virescens. var. 2. With. On the top

of Carnedd Llewelin. Mr.	Griffith.
LICHEN luteus. Wood near Bangor.	D. T.
Acharii. Rocks by the side of the River	r
at Beddgelart.	Id.
	t
Port Penrhyn.	Id.
escharoides. SummitofCarnedd Llewelyn	•
Mr. Griffith. Rocks by the Menai, near	r
Bangor Ferry.	D. T.
pezizoides. Rocks by the Menai near Bangor Ferry.	r
luridus. 5 Bangor Ferry.	Id.
crassus. On the Glyder.	Dillenius.
gelidus. Between Llanberris and Pen y	7
Gorphwysfa: also on stones in Cwm Id-	-
wel, particularly near Twll Dû. Mr.	
Griffith. Stone fence at Beddgelart, be-	-
tween the Inn and Pont Aberglâslyn.	D. T.
polyphyllus. Rocks near Perfeddgoed	l
House by Bangor.	Dillenius.
proboscideus. Rocks and Stones near Llan-	
berris. Brewer. Summit of Carnedd	
Llewelyn. Mr. Griffith. Snowdon, on	
rocks by Llyn Llydaw; and near Llyn	i -
Cwm y Fynnon Lâs: also on the tops of	
Mountains from Cwm Brwynog towards	
Ardhu near Llanberris.	Dillenius.
torrefactus. With the former.	Id.
pustulatus. On 2 large Mass of Rock on	

the right of the road from Penmorvay to Dolbelmen. Dillenius. On an old wall about half way between Carnarvon and Beddgelart. Rev. H. Davies. Rocks in Nant Hwynant Vale at the foot of Snowdon. D. Ta

LICHEN deustus. About Llanberris. Rev. H. Davies. - pellitus. On Snowdon, by Llyn Llydaw; and about Llyn Cwm y Fynnon Lâs: also on the tops of Mountains, from Cwm Brwnog towards Ardhu, by Llanberris. Dillenius. On the top of a high hill called Moel Shiabod, near Capel Cerig. Mr. Grifhth. amphibius. With. Stones near Llyn Idwel; and on many of the Stones under the water in the Lake. Id. --- aquaticus. With the former. Mr. Griffith. Rocks in a rivulet near Perfeddgoed House by Bangor. Dillenius. - tiliaceus. Rocks in this County. Rev. H. Davies. _____ diatrypus. Stones immediately over the Bridge at Beddgelart, at the entrance of the Vale of Nant Hwynant. D. T. physodes. In fruit, but rarely, on the _____ perlatus. S stone fences near Beddgelart. D. T. aquilus. Dry rocks about Bangor. Dillenius.

conspersus. Ach. Abundant on rocks and

roofs of cottages about Beddgelart, and
Nant Francon. D. T.
LICHEN fablunensis. Stones near the summit of
Carnedd Llewelin. Mr. Griffith.
islandicus. Between Bwlch Lâs, and
the summit of Snowdon, but not in fruit.
Mr. Griffith. On Clogwyn y Garnedd.
Dillenius.
plumbeus. Trees near Pentir and Bangor. Id.
* rupestris? On the bridge at Beddgelart
and rocks in the neighbourhood. D. T.
tenuissimus. Crib y Ddescil, and Clog-
wyn y Garnedd. Mr. Griffith.
fluviatilis. In the Alpine Torrents of
Snowdon. Dillenius.
on Rocks, by the fide of the
River adjoining Pont Aber-
glaslyn: the two first in
fruit · (not upon Snowdor
glish Botany.) D. T.
saccatus. On Snowdon, on the rocks of
Llyn Cwn y Ffynnon Velan; and about
Clogwyn y Garnedd. Dillenius. Crib y
Ddescil. Mr. Griffith.
aphthosus. Near the summit of Snowdon,

* I have subjoined a mark of doubt to this plant, not having found it in fruit. D. T.

where Mr. Griffith has found it in fruit,	
though I did not.	D. T.
LICHEN lanatus. Rocks about Llyn y Cwn, but	
never in fruit. Mr. Griffith. Rocks	
about Borth, a mile from Bangor. Dil-	
lenius. About Llanberris. H	ludson.
pubescens. On Snowdon. Dillenius. Car-	
nedd Llewelin, and Carnedd Dafydd.	
Mr. Griffith. Glyder Vawr. P	ennant.
exilis. Frequent on Rocks about Beddgelart.	D. T.
tristis. On Snowdon, from Cwm Brwy-	
nog towards Ardhu. Di	llenius.
fragilis. On the Summit of Carnedd	
Llewelin.	Id.
globiferus. On Snowdon.	Id.
CONFERVA confragosa. Near Llanberris.	Id.
myochrous. Dillav. In Alpine torrents at	
Beddgelart, and the lower regions of	
	D. T.
corallina. On an Island, Ynys y Moch	
C C	llenius.
gelatinosa. S. Huds. In Llyn Fynnon Ve-	
	D. T.
fluviatilis. Near Bangor, in a stream of	
fresh water, called Afon Cegin, between	
the Mill called Pander Maes y Geirchan,	
and Melun Escob, growing thickly on	
the stones. Dil.	lenius.

FOR almost the whole of what we are here enabled to offer respecting the Botany of Cheshire, we have to express our obligations to Edward Hawkins, junr. Esq. of Court Herbert, by Neath, and Luke Howard, Esq. of London; the former of whom procured us the habitats followed by the name of Mr. Bradbury, the latter those of Mr. G. Holme, through the medium of Mr. Ollive Sims, of Stockport. We are persuaded, however, that very much is yet to be done in this interesting county, which is in many respects peculiarly favourable to the cultivation of Natural History, and we trust a more perfect catalogue of its vegetable productions will not be suffered long to remain a desideratum to the Flora Britannica.

CHARA hispida. In Ponds, frequent.	Mr. Bradbury.
CIRCÆA alpina. On the Borders of Yo	orkshire in
Greenfield, but whether within	the bounds
of Cheshire, I cannot say.	Id.
UTRICULARIA minor. In old Turf-pit	ts, Dean's
Moss. Mr. Bradbury. Whit	tley Reed.
	Mr. A. Aikin.
CROCUS vernus. Meadows near Gorton,	four miles
from Stockport.	Mr. G. Holme.
SCHOENUS albus. On almost all the Bog	s. Mr. Bradbury.
ERIOPHORUM vaginatum. On the Me	oors above
Stayley Wood, &c,	Id.
MELICA nutans. In Woods frequent	: as Early
Banks-Wood, &c.	Id.
BRIZA minor. In high Pastures, Stay	ley. Id.
(surely this cannot be correct.)	
AVENA fatua. Often found with the	cultivated
Oat.	Id.
DIPSACUS Fullonum. Sometimes culti-	vated near
Stayley.	Id.
pilosus. Near Broken Brow	by Stock-
port.	Mr. G. Holme.
GALIUM anglicum. On Wild Bank, ne	ar Stayley
Bridge.	
	Mr. Bradbury.
SANGUISORBA officinalis. Dry Meadow	
SANGUISORBA officinalis. Dry Meadow New Bridge at Stockport.	vs near the
	vs near the Mr. G. Holme.

LYSIMACHIA thyrsiflora. Woods near Stockport.
Mr. G. Holme.
CAMPANULA latifolia. Near the rivers Mersey
and Tame at Stockport. Id.
IMPATIENS noli angere. Near Lyme Hall. Id.
VERBASCUM Lychnitis. Between Chester and
Wrexham. Rev. W. Wood.
Blattaria. Between Stockport and
Reddish. Mr. G. Holme.
DATURA Stramonium. Cross Road between
Chorley and Chelford. Id.
RIBES alpinum. Stayley Wood, in a place called
Stocks. Mr. Bradbury.
CUSCUTA europea. Has been found in Cheshire,
but is rare. Mr. G. Holme.
GENTIANA pneumonanthe. In a Marsh on a Heath
near Holmes Chapel. Mr. Hunter. Not
uncommon about Park-Gate. Mr. Griffith.
campestris. In a high Pasture above
Stayley Hall, called Flax-field. Mr.
Bradbury. Tatton Park, near Knuts-
ford. Mr. A. Aikin. Common on the
Sea Coast and other parts of the County. Ray.
HERACLEUM sphondylium. B. Hedges round Mr.
Ward's Meadow at Macclesfield. Mr. Bradbury.
CICUTA virosa. Pits near the Little Moor, a
mile from Stockport. Mr. G. Holme.
Pond two miles from Northwich, on

the road to Chester. Rev. W. Word. Brereton Mere. Ray. SCANDIX odorata. Banks of rivulets and near Farm Houses. Mr. Bradbury. Banks of the rivers near Stockport. Mr. G. Holme, ---- cerefolium. Lanes near Bageley by Stockport. Id. IMPERATORIA Ostruthium. Near J. Oldham's Garden at Hyde, probably an outcast. Mr. Bradbury. PIMPINELLA magna. Woods and Fields belonging to Stayley Hall. Id. SAMBUCUS Ebulus. In Rainow, road side Bridge Id. near Goodwin's Mill. DROSERA longifolia. Moors above Stayley Wood. Mr. Bradbury. Knutsford, and other Mr. L. Howard. Moors. anglica. In a Marsh on a Heath near Mr. Hunter. Holme's Chapel. NARCISSUS biflorus. In some Closes at Bellow Hill, near Whitechurch. Blackstone. ORNITHOGALUM luteum. In a Close near Willow Moor at Bellow Hill. Id. umbellatum. Meadows near Cheadle Bridge, three miles from Stock-Mr. G. Holme. port. Acorus calamus. In the River Dee, but rare. Mr. G. Holme. At Holford, near Northwich, plentifully. Blackstone. Abundant Raye in the county.

Juncus filiformis. In marshy places, or where water has stood. Mr. Bradbury. COLCHICUM autumnale. Meadow betwixt Castle Hall, and River, Stayley. Id. ALISMA ranunculoides. Pond in Tatton Park. Id. EPILOBIUM angustifolium. In Longdendale, above Tintwistle, in a wild state. Id. CHLORA perfoliata. Fields from New Ferry to Park-Gate. Id. VACCINIUM Vitis idea. Abundantly in old Slate Quarries on Stayley Moors, where it bears fruit in Summer and at Christmas. Mr. Bradbury. Common on the Moors. Blackstone. ---- Oxycoccos. On Gate Moor, and several other below Bellow Hill, near Whitechurch. Blackstone. Moors near Stockport. Mr. L. Howard. POLYGONUM Bistorta. Frequent in meadows. Mr. Bradbury. Near Stockport. Mr. G. Holme. PARIS quadrifolia. In a wood near the Bolling; and not far from Seale Moor. Mr. Bradbury. Wood in Brinnington, near Stockport. Mr. G. Holme. ANDROMEDA polifolia. Moors and Bogs, frequent. Mr. Bradbury. Moors near Stockport. Mr. I. Howard. ARBUTUS Uva Ursi. On the hills betwixt VOL. I. I

Cheshire and Yorkshire, near Greenfield. Mr. Bradbury. CHRYSOSPLENIUM alternifolium. Very common in the County. Mr. Bradbury. Banks of the river, one mile above Stockport. Mr. G. Holme. SAXIFRAGA Hirculus. Found by Dr. Kingston on Knutsford Moor, and there shown by him to Dr. Richardson, now growing Ray. in plenty. ---- aizoides. On Beeston Castle, and on a high Hill in Wyrswall Town, near Blackstone. Malpas. SAPONARIA officinalis. On the Banks of the Mersey, near Stockport, but rare. Mr. G. Holme. SILENE Armeria. On the Banks of the river, half a mile below Chester. Dr. Richardson. STELLARIA Nemorum. Shady Woods near Stock-Mr. G. Holme. port. ARENARIA verna. y. High Pastures at Stayley. Mr. Bradbury. PRUNUS Padus. Common in woods. Mr. Brad-Mr. G. Holme. bury. Near Dishley. MESPILUS Germanica. In all the hedges about Du Bois. Muishull. RUBUS Chamamorus. Mountains above Stayley Wood, where I have often gathered the Mr. Bradbury. fruit.

TORMENTILLA reptans. Moors and waste ground
on all sides of Stockport. Mr. G. Holme.
GEUM rivale. Frequent in the Wems, a marshy
place near Stayley Hall. Mr. Bradbury.
STRATIOTES aloides. In several ponds near Gor-
ton, four miles from Stockport. Mr,
G. Holme. Pools about four miles from
Holme's Chapel. Withering.
RANUNCULUS Lingua. Bridgewater Canal, from
Manchester to Warrington. Mr. Bradbury.
TROLLIUS europæus. In great plenty in a wood
betwixt Stayley Hall and Scout Mill, at
a place called Wems, from which I sent
two or three hundred plants to Lee of
Hammersmith. Id.
MENTHA piperita. In rivulets. Id.
MENTHA piperita. In rivulets. Id.
MENTHA piperita. In rivulets. Id.
MENTHA piperita. In rivulets.
MENTHA piperita. In rivulets. Id.
MENTHA piperita. In rivulets. Id.
 MENTHA piperita. In rivulets. Id. Odorata. Very common by the sides of rivers and brooks in Cheshire, especially about Aston House. Mrs. Walmesly. Pulegium. Godly Hill, between Mottram and Gee Cross. Mr. Bradbury. GALEOPSIS versicolor. Northwich. Rev. H. Davies.
 MENTHA piperita. In rivulets. Id. Odorata. Very common by the sides of rivers and brooks in Cheshire, especially about Aston House. Mrs. Walmesly. Pulegium. Godly Hill, between Mottram and Gee Cross. Mr. Bradbury. GALEOPSIS versicolor. Northwich. Rev. H. Davies. SCUTELLARIA minor. Boggy places on Stayley
 MENTHA piperita. In rivulets. Id. odorata. Very common by the sides of rivers and brooks in Cheshire, especially about Aston House. Mrs. Walmesly. Pulegium. Godly Hill, between Mottram and Gee Cross. Mr. Bradbury. GALEOPSIS versicolor. Northwich. Rev. H. Davies. SCUTELLARIA minor. Boggy places on Stayley Moor. Mr. Bradbury.
 MENTHA piperita. In rivulets. Id. odorata. Very common by the sides of rivers and brooks in Cheshire, especially about Aston House. Mrs. Walmesly. Pulegium. Godly Hill, between Mottram and Gee Cross. Mr. Bradbury. GALEOPSIS versicolor. Northwich. Rev. H. Davies. Scutellaria minor. Boggy places on Stayley Moor. Mr. Bradbury. BARTSIA viscosa. Near Chester. Rev. H. Davies.
 MENTHA piperita. In rivulets. MENTHA piperita. In rivulets. odorata. Very common by the sides of rivers and brooks in Cheshire, especially about Aston House. Mrs. Walmesly. Pulegium. Godly Hill, between Mottram and Gee Cross. GALEOPSIS versicolor. Northwich. Rev. H. Davies. SCUTELLARIA minor. Boggy places on Stayley Moor. Mr. Bradbury. BARTSIA viscosa. Near Chester. Rev. H. Davies. North West of Park-Gate. Mr. Pennant.
 MENTHA piperita. In rivulets. Id. odorata. Very common by the sides of rivers and brooks in Cheshire, especially about Aston House. Mrs. Walmesly. Pulegium. Godly Hill, between Mottram and Gee Cross. Mr. Bradbury. GALEOPSIS versicolor. Northwich. Rev. H. Davies. Scutellaria minor. Boggy places on Stayley Moor. Mr. Bradbury. BARTSIA viscosa. Near Chester. Rev. H. Davies.

I 2

MELAMPYRUM sylvaticum. Woods near Disley.
Mr. G. Holme. In many parts of the
county. Mr. Bradbury. (Is it not more
probable that M. pratense is intended?)
LATHRÆA squamaria. In Harnicroft Wood, near
Chadkirk. Mr. Bradbury.
ANTIRRHINUM Cymbalaria. On the road side at
Bank-Top, near Stockport. Mr. G. Holme.
Garden Walls at Addington. Mr. Bradbury.
LIMOSELLA aquatica. Brought me from near
Frodsham. Id.
THLASPI arvense. Corn fields, frequent. Id.
COCHLEARIA officinalis. Sea-coast below Park-
gate. Id.
ISATIS tincloria. Near Broken Brow, Stockport.
Mr. G. Holme.
CARDAMINE impatiens. A common weed in
Gardens. Mr. Bradbury.
amara. Brooks and shady places in
woods. Id.
SISYMERIUM sylvestre. Brook sides-about Maccles-
field. Id.
ERYSIMUM cheiranthoides. Near Stockport. Mr. G. Holme.
ERODIUM moschatum. In the Lane between Stock-
port and the New Bridge. Id.
maritimum. Coast near Park-gate. Mr. Bradbury.
GERANIUM pratense. Common in the County.
Mr. Bradbury. Meadows near Stock- Mr. G. Holme.

* FUMARIA lutea. On old walls near Marple, four miles from Stockport. Mr. G. Holme. VICIA lathyroides. Macclesfield Common; and road to Broken Cross. Mr. Bradbury. HYPERICUM androsemum. In a wood near Broken Brow. Mr. G. Holme. Woods near Scout Mill. Mr. Bradbury. CARDUUS eriophorus. A little short of Gatehouse Green, in the road from Middlewich. plentifully. Waring. ------- heterophyllus. Near Stayley Bridge, in a place called Wharf. Mr. Bradbury. BIDENS cernua. B. About Tapperley, and elsewhere in the County. Hudson. GNAPHALIUM dioicum. Mountains above Stayley, &c. Mr. Bradbury. Stockport. Mr. G. Holme. TUSSILAGO hybrida. Near Stockport. Mr. G. Holme. Cheshire side of the Tame below Stayley Bridge. Mr. Bradbury. SENECIO viscosus. On dry Banks; Walls; and

* " This is a Habitat of the Plant, which Withering calls F. capnoides, and inserts as first found by me in Derbyshire. The latter fact is somewhat doubtful, as I rather think the specimen came from Marple, and, if so, the situation being on walls in a Village, it might originally have been cultivated; being found in a similar state near Gardens hereabout." Mr. L. Howard.

Rocks at Ince. Waring. Road side from Northwich to Delamere Forest. Mr. Bradbury. SENECIO paludosus. By the Eastern side of the Dee, just above Eaton Boat, sparingly. Waring. INULA Helenium. About old Farm Houses. Mr. Bradbury. SATYRIUM viride. 7 Both found in Cocker's albidum. SFields, Stayley Wood. Id. OPHRYS cordata. Stayley Moors. Mr. Bradbury. On Longside Hill, near Disley. Mr. G. Holme. SERAPIAS long folia. Woods near Stayley Hall, Mr Bradbury. Knutsford Moor. Mr. A. Aikin. TYPHA angustifolia. In a pond by the Foot-path from New Ferry to Park-gate, about half way. Mr. Bradbury. CAREX curta. Bogs, not uncommon. Id. pendula. River side above Stayley Bridge. Id. ------ atrata. On Higher Edge; and Wild Bank. Id. ------- distans. (binervis?) Common. Id. MYRIOPHYLLUM verticillatum. Stockport little Moor, in ponds. Mr. G. Holme. SALIX pentandra. Woods about Stayley Hall, and Common. Mr. Bradbury. EMPETRUM nigrum. Stayley and other Moors, Id. common. EQUISETUM sylvaticum. Woods near Stayley Hall; and in many other places. Id.

EQUISETUM hyemale. Near Arden Hall, in a	
Wood close to the River. Mr. Br	adbury.
OSMUNDA Lunaria. Fields about Hooly's Folly,	
near Macclesfield.	Id.
regalis. Bog near Gawsworth.	Id.
LYCOPODIUM alpinum. Moors above Mickle-	
hurst.	Id.
POLYBODIUM Phanetteria Zarly Banks Wood,	
POLYPODIUM Phegopteris. Dryopteris. Early Banks Wood, near Stayley Bridge; and elsewhere.	
Dryopteris.) and elsewhere.	Id.
ASPIDIUM Thelypteris. Harnicroft Wood, near	
Wernith; and other woods.	Id.
ASPLENIUM viride. Among stones and rubbish	
thrown out of the Quarries on Carr Edge.	Id.
SCOLOPENDRIUM Ceterach. Old Walls at Carr	
Edge, going to the Stone Quarries.	Id.
PTERIS crispa. On the top of Tag's Ness, a Hill	
near Macclesfield.	Id.
HYMENOPHYLLUM Tunbrigense. On mossy	
rocks near Croyden Brook, the Eastern	
extremity of Cheshire; and on the hills	
from Macclesfield to Buxton, in similar	
situations.	Id.
ANDRÆA alpina. ————————————————————————————————————	
Higher Edge, a high hill	
Jabove Stayley Bridge.	Id.
SPLACHNUM vasculosum. Moors above Stayley	
Wood, on Sheep's dung.	Id.

SPLACHNUM ovatum. Deans Moss. Mr. Br	adbury
Brewerianum. Delamere Forest.	·
Mr. J. De	roburst
GRIMMIA pusilla. On a Stone on the Hill above	
Macclesfield. Mr. Br	adbury
TORTULA tortuosa. Common on the top of old	
Walls in Brushes Woods.	Id.
DICRANUM pellucidum. Near Johnson's Pond,	
Macclesfield.	Id.
POLYTRICHUM alpinum. Moors and Heaths,	
Alderly Edge.	Id.
NECKERA pumila. Trunks of trees in Brookbank	
Wood.	Id.
curtipendula. Stones in Brushes Woods.	Id.
BARTRAMIA arcuata. First found in fruit in a	
Valley, called the Brushes, in Stayley	
Wood.	Id.
BRYUM nutans. Deans Moss: Stayley Moors,	
&c.	Id.
HYPNUM fluitons. In fruit in old Turf-pits on	
Deans Moss: the pedicles vary in length	
in proportion to the depth at which the	
plant swims : I have seen them six inches	
long.	Id.
rugosum. On Alderly Edge.	Id.
JUNGERMANNIA quinquedenteta. In the Woods	
under Buckton Castle, a Hill near	
Mofley.	Id.

MARCHANTIA cruciata. Shady Garden Walks,	
frequent: moist rocks near Stockport.	
Mr. Br.	adlury.
conica. Common on all moist	~
rocks covered with earth.	Id.
BLASIA pusilla. On breaches made by the rivulet,	
about half a mile above Stayley Mill.	Id.
LICHEN corallinus. Frequent on Stone Walls,	
Stayley.	Id.
geographicus. On Pudding Stone on all	
the Cheshire Hills.	Id.
sanguinarius. Armfield Moor near Tint-	
wistle.	Id.
ventosus. Only seen on Higher Edge on	
Pudding Stone.	Id.
conspersus. Stone Walls near Stayley	
Moor: Barn in Bollington, &c.	Id.
omphalodes. Frequent on Stones on the	
Mountains.	Id.
plumbeus. Bank Wood near Mottram in Longdendale, frequently on the same tree.	
in Longdendale, frequently	
late-virens.) on the same tree.	Id.
furfuraceus. Common on Stone Walls:	
in fruit near Hooly's Folly by Maccles-	
field.	Id.
glaucus. Frequent and fine in Brushes	
TTT J. TTT 11	Id.

often found in fruit. Mr. Bradbury. serobiculatus. Stunted Oaks in Brushes Woods. Id. borizontalis. Bank Wood near Mottram in Longdendale. Id. resupinatus. Near Broad Bottom Bridge, scarce. Id. deustus. On the Borders of Yorkshire, on Biddistone. Id. polyphyllus. On large Stones near pellitus. Greenfield. Id. pustulatus. Near Biddistone Light House, Id. in plenty. Id. Id.	LICHEN pulmonarius. Great Wood, Stayley	:
scrobiculatus. Stunted Oaks in Brushes Woods. Id. borizontalis. Bank Wood near Mottram in Longdendale. Id. resupinatus. Near Broad Bottom Bridge, scarce. Id. sepincola. On the Borders of Yorkshire, on some large Stones above Micklehurst, near Greenfield. Greenfield. Mr. Dickson informed me these are the only specimens found in England. England. Id. <i>deustus.</i> On Stones near the Light House, Biddistone. Id. <i>polyphyllus.</i> On large Stones near <i>pellitus.</i> Greenfield. Id. <i>pustulatus.</i> Near Biddistone Light House, in plenty. in plenty. Id. Id. <i>fragilis.</i> Large Stones on the Moors globi/erus. globi/erus. above Micklehurst. Id. <i>pascbalis.</i> Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id. Id. Id.		
 borizontalis. Bank Wood near Mottram in Longdendale. resupinatus. Near Broad Bottom Bridge, scarce. Id. resupinatus. Near Broad Bottom Bridge, scarce. Id. sepincola. On the Borders of Yorkshire, on some large Stones above Micklehurst, near Greenfield. Mr. Dickson informed me these are the only specimens found in England. deustus. On Stones near the Light House, Biddistone. polyphyllus. polyphyllus. polyphyllus. forcenfield. for large Stones near pellitus. pustulatus. Near Biddistone Light House, in plenty. fragilis. Large Stones on the Moors globiferus. above Micklehurst. pascbalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. tristis. Windberry Stones, West-side of 		
in Longdendale. Id. resupinatus. Near Broad Bottom Bridge, scarce. Id. sepincola. On the Borders of Yorkshire, on some large Stones above Micklehurst, near Greenfield. Mr. Dickson informed me these are the only specimens found in England. Id. deustus. On Stones near the Light House, Biddistone. Id. polyphyllus. On large Stones near pellitus. Id. polyphyllus. On large Stones near pellitus. Id. pustulatus. Near Biddistone Light House, in plenty. Id. fragilis. Large Stones on the Moors globiferus. Id. pasebalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id.	Woods.	Id.
 resupinatus. Near Broad Bottom Bridge, scarce. sepincola. On the Borders of Yorkshire, on some large Stones above Micklehurst, near Greenfield. Mr. Dickson informed me these are the only specimens found in England. deustus. On Stones near the Light House, Biddistone. deustus. On Stones near the Light House, Biddistone. polyphyllus. On large Stones near pellitus. polyphyllus. Greenfield. torrefatus. pustulatus. Near Biddistone Light House, in plenty. fragilis. Large Stones on the Moors globi/erus. above Micklehurst. pascbalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. tristis. Windberry Stones, West-side of 	horizontalis. Bank Wood near Mottram	
scarce. Id. sepincola. On the Borders of Yorkshire, on some large Stones above Micklehurst, near Greenfield. Mr. Dickson informed me these are the only specimens found in England. Id. deustus. On Stones near the Light House, Biddistone. Id. polyphyllus. On large Stones near pellitus. On large Stones near pellitus. Con large Stones near pellitus. Id. pustulatus. Near Biddistone Light House, in plenty. Id. fragilis. Large Stones on the Moors globi/erus. above Micklehurst. Id. paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id.	in Longdendale.	Id.
scarce. Id. sepincola. On the Borders of Yorkshire, on some large Stones above Micklehurst, near Greenfield. Mr. Dickson informed me these are the only specimens found in England. Id. deustus. On Stones near the Light House, Biddistone. Id. polyphyllus. On large Stones near pellitus. On large Stones near pellitus. Con large Stones near pellitus. Id. pustulatus. Near Biddistone Light House, in plenty. Id. fragilis. Large Stones on the Moors globi/erus. above Micklehurst. Id. paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id.	resupinatus. Near Broad Bottom Bridge,	
some large Stones above Micklehurst, near Greenfield. Mr. Dickson informed me these are the only specimens found in England. Id. deustus. On Stones near the Light House, Biddistone. Id. polyphyllus. On large Stones near pellitus. On large Stones near pellitus. Id. pustulatus. Near Biddistone Light House, in plenty. Id. fragilis. Large Stones on the Moors globiferus. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id.	_	
Greenfield. Mr. Dickson informed me these are the only specimens found in England. Id. deustus. On Stones near the Light House, Biddistone. Id. polyphyllus. On large Stones near pellitus. On large Stones near pellitus. Greenfield. Id. torrefactus. Vear Biddistone Light House, in plenty. Id. pustulatus. Near Biddistone Light House, in plenty. Id. fragilis. Large Stones on the Moors globiferus. above Micklehurst. Id. paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id.	sepincola. On the Borders of Yorkshire, on	
these are the only specimens found in England. Id. Id. Id. Id. Id. Id. Id. Id.	some large Stones above Micklehurst, near	
England.Id.deustus.On Stones near the Light House, Biddistone.Id.polyphyllus. pellitus.On large Stones near Greenfield.Id.pellitus. pustulatus.On large Stones near Greenfield.Id.pustulatus. pustulatus.Id.Id.fragilis. globiferus.Large Stones on the Moors above Micklehurst.Id.pascbalis. five miles from Macclesfield, on the Buxton road.Id.	Greenfield. Mr. Dickson informed me	
deustus. On Stones near the Light House, Biddistone. Id. polyphyllus. polyphyllus. On large Stones near foreenfield. Id. pellitus. Greenfield. Id. pustulatus. Near Biddistone Light House, in plenty. Id. fragilis. Large Stones on the Moors globiferus. Id. paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id.	these are the only specimens found in	
Biddistone. Id. polyphyllus. On large Stones near pellitus. Greenfield. Id. torrefactus. Id. pustulatus. Near Biddistone Light House, Id. in plenty. Id. fragilis. Large Stones on the Moors Id. globiferus. above Micklehurst. Id. pascbalis. Stone Walls at Brushes; and Id. five miles from Macclesfield, on the Buxton road. Id. tristis. Windberry Stones, West-side of Id.	England.	Id.
polyphyllus. pellitus. pellitus. forcenfield. Id. pustulatus. Near Biddistone Light House, in plenty. fragilis. globiferus. babove Micklehurst. Id. paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id.	deustus. On Stones near the Light House,	
— pellitus. Greenfield. Id.	Biddistone.	Id.
 pellitus. forrefa&us. pustulatus. Near Biddistone Light House, in plenty. fragilis. Large Stones on the Moors globiferus. above Micklehurst. Id. paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id. 	polyphyllus.] On large Stones near	
 torrefactus. pustulatus. Near Biddistone Light House, in plenty. fragilis. Large Stones on the Moors globiferus. above Micklehurst. Id. paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id. 	hellitus	F.J
in plenty. Id. ————————————————————————————————————	torrefactus.	a w e
 fragilis. Large Stones on the Moors globiferus. above Micklehurst. Id. paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id. tristis. Windberry Stones, West-side of 	pustulatus. Near Biddistone Light House,	
— globiferus. S above Micklehurst. Id. — paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id. — tristis. Windberry Stones, West-side of	in plenty.	Ida
— globiferus. S above Micklehurst. Id. — paschalis. Stone Walls at Brushes; and five miles from Macclesfield, on the Buxton road. Id. — tristis. Windberry Stones, West-side of	fragilis. 7 Large Stones on the Moors	
five miles from Macclesfield, on the Buxton road. Id. tristis. Windberry Stones, West-side of	globiferus. 5 above Micklehurst.	Id.
Buxton road. Id. ————————————————————————————————————	paschalis. Stone Walls at Brushes; and	
tristis. Windberry Stones, West-side of	five miles from Macclesfield, on the	
	Buxton road.	Id.
Greenfield. Id.	tristis. Windberry Stones, West-side of	
	Greenfield.	Id.

CHESHIRE.

LICHEN	articulatus.	Old	Oaks, .	Lime	Park, in	
	plenty.				Mr. Br.	adbury.
(lanatus.	Large	Stone	s on	Stayley	~
	Moors.					Id.

CHARA flexilis. Between Bodmin and Wadebridge. L. W. D. VERONICA spicata. About Pensance. Withering. montana. In Brown's hill wood, Bo-Mr. E. Forster, jun. connoc. PINGUICULA lusitanica. Plentiful in this County. Hudson. In the marshes at Kilkhampton. Heath. VALERIANA dentata. Cornfield at Boconnoc. Mr. E. Forfter, jun. SCHOENUS mariscus. By the sea side between Pensance and Marazion. Ray. ---- albus. Many boggy places in this

County. Ray. In Torr Valley on Bra-Mr. E. Forfler, jun. doc Downs. SCIRPUS multicaulis. With the former. Mr. E. Forster, jun. About the Wherry Mine Mr. W. Allen. near Pensance. ERIOPHORUM vaginatum. On wet Moors, not uncommon. Withering. Near Pensance, by the Cromlech. D.T.PANICUM Dactylon. On the sandy shore between Pensance and Marazion. Ray. AGROSTIS setacea. Scarcely a heath in this County, on which it does not abound. Curtie. BRIZA minor. Cornfields between Pensance and Marazion. Lightfoot. At Boconnoc. Mr. E. Forster, jun. AVENA strigosa. On St. Winnoe Downs, by the roadside from the Tollgate, Boconnoc, to Lostwithiel. IJ. TRITICUM loliaceum. On the Coast throughout the County. Id. EXACUM filiforme. Near the Land's End; and between Pensance and St. Ives. Martyn. Sandbanks between St. Blazey Bridge and the Par Sand. Mir. Stackhouse SANGUISORBA officinalis. Goonhilly Downs. Mr. W. Allen. RUPPIA maritima. Salt ditches in this County. Mr. Stackhouse,

.

ANCHUSA semper-virens. About Fawy. Mr. E.
Forster. jun. Near Liskeard. D. T
CAMPANULA latifolia. Moist shady places, very
frequent. L. W. D
hederacea. Moist shady places near
Boconnoc, not uncommon. Mr. E.
Forster, jun. Near Falmouth; Pen-
sance; Camelford, &c. D. T.
VIOLA lactea. Bradoc Downs. Mr. E. Forster, jun.
Pendarvis. Mr. Stackhouse. Heaths be-
tween Liskeard and Lostwithiel. D. T.
VERBASCUM Blattaria. By the roadside near
Morval. Mr. E. Forster, jun.
CHIRONIA pulchella. On the Downs, near the Sea,
at Port Owen, on the North Coast. Mr. Watt,
ILLECEBRUM verticillatum. In watery places be-
tween St. Columb and St. Michael : also
about Pensance; and towards the Land's
End. Heath. On Talloe Water, Bradoc.
Mr. E. Forster, jun. Among the rocks
at Castle Treryn. Borlase.
HERNIARIA glabra. At the Lizard Point. Hud-
son. By the soap rocks. Mr. E. Forster, jun.
birsuta. Sent from this County to Dr. Smith.
GENTIANA campestris. About Perian Round. Hudson.
DAUCUS Carrota, var. 5. With. (D. Polygamus.
Gouan.) Sent from this County by Mr.
Giddy. Withering.

DAUCUS * maritimus. With. Rocks about Pensance. Rev. J. T. Thomson. At Tintagel Castle: the Lizard, and elsewhere.

Mr. E. Forster, jun. Castle Treryn. D. T. LIGUSTICUM Cornubiense. In a wheat field, and in an adjoining coppice called Marget, or Margaret, Wood, about three furlongs from the Bodmin Turnpike, that leads to Launceston. Sir. T. Cullum. Shown to Mr. Sowerby and me in fields about half a mile north of Bodmin, by Dr. Hall, plentifully. D. T. In a wood, and cornfield at Cardinham Parsonage; and in Draw-wood, Bradoc. † Mr. E. Forster. jun.

* Dr. Smith in the Flora Britannica has reduced this plant to a var. of D. Carrota, and observed that the leaves " in maritimis carnosa, lucida, subinde hirsutissima. vel lanata." I must, however, differ from him in opinion. I never saw on any part of the Coast, except in Cornwall, a plant approaching to D. maritimus, of which the whole habit is distinct from the common species; and the leaves very unlike in form, as well as appearance and hairiness, besides other differences pointed out by Dr. Withering. Unless it suffers very material alteration from culture, which I understand it does not, it appears to me to have as clear a right to be considered a species as any other in our Flora. D. T.

† "These are the only places in which I have seen it; but I have heard of it elsewhere, though always within a few miles of Bodmin. It has not yet been found in other parts of the County, so that "in Cornubia" generally should not stand. I suppose Dille-

TAMARIX Gallica. Plentiful on St. Michael's mount, and every where about the Lizard Point, but chiefly on the Banks of Earth called Hedges. Mr. Giddy. CORRIGIOLA littoralis. On the Beach near the Tin Mine at Helston. Mr. F. Borone. On the Shore of Loe Pool, near Helston, opposite Penrose on each side of a low Mr. E. Forster, jun. wall. LINUM usitatissimum. Falmouth Cliffs. D. T.----- angustifolium. Pastures by the sea side about St. Ives and Truro. Ray. DROSERA longifolia. In Torr Valley on Bradoc Mr. E. Forster, jun. Downs. SCILLA verna. About St. Ives. Hudson. Near Redruth. Mr. Watt. Near Pensance, Withering. Abundant on the heathy ground by the soap rocks, the Lizard, and the Land's End. D. T.- autumnalis. Common near Chace Water Mine. Mr. Watt. Lizard Point. Heath. ASPARAGUS officinalis. Cliffs at the Lizard Point. Borlase.

nius did not recollect the exact place, and Dr. Smith imagined it to be a common Cornish plant, which it is not, but a local one, though, I dare say, thousands of Acres might be produced." Mr. E. Forster, jun.

129

JUNCUS acutus. Sandy hills near Gwythien, and Philac. Martyn. ALISMA ranunculoides. Bog by the road side, between Pensance and Marazion. Mr. E. Forster, jun. Marsh at Marazion. Withering. * ERICA vagans. Goonhilly Downs, going from Helston to the Lizard Point. Ray. Near Pendarvis, on the road from Camborn to Hayle, in the low ground between Tregothan and Connor Downs. Mr. Stackhouse. I have seen it near the Land's End. Borlase. - var. flore albo. Near Mullein. Maton. SAXIFRAGA stellaris. Fissures of rocks near Castle Treryn. Id. STELLARIA glauca. About Falmouth. Withering. ARENARIA tenuifolia. Side of a watery bank about three miles and a half from the Land's End. Ray. EUPHORBIA Peplis. On the sandy shore between Pensance and Marazion, plentifully. Id.

* "Ray's habitat " in cricetis in Cornubia" is a great mistake, as it is quite a local plant, and none grows in the Eastern part of the County. Withering has mentioned it at the Land's End: this I cannot deny, though I did not see it there, and the people put one of the common species into my hands as something curious, and therefore, I suppose, deceive those who know no better. I judge it cannot abound in that part, if it grow there at all, and there are extensive heaths in the East on which not a morsel is to be seen." Mr. E. Forster, jun.

VOL. I.

EUPHORBIA Portlandica. Between Pensance and
Marazion. Mr. W. Allen.
Pyrus domestica. In the mountainous parts of this
County. Ray.
HELLEBORUS viridis. Hedge in Bradoc Church
Yard. Mr. E. Forster, jun.
MENTHA rotundifolia. By the road side from
Trengoff to Panter's Bridge, Warleggon:
near Looe; and between Bossiney and
Tintagel Castle. Id.
gentilis. St. Berian. Martyn.
MELITTIS melissophyllum. Road side between Lis-
keard and Callington. Withering. (As
Dr. Withering does not distinguish be-
tween this species and the following, and
the habitat here given seems to be the
same in which Mr. Sowerby and I found
M. grandiflora, I cannot doubt but that
plant is intended. D. T.)
grandiflora. Near Liskeard on hedges
in the road to Lostwithiel. D. T.
BARTSIA viscosa. Boggy and watery places towards
the farther end of the County. Ray.
Cornfields near Plengwarry: In a field
close by Perran, near Penrhyn, plentifully.
Mr. W. Allen Moist places at Bocon-

noc. Mr. E. Forster, jun. Marshes by

Pensance.

D. T.

ANTIRRHINUM repens. Hedges near Penrhyn.
Ray. Stone wall between Penrhyn and
Falmouth. Mr. Wutt.
C. I
SCROPHULARIA scorodonia. Near the sea shore
of St. Trees 1 . C. H
Hudson.
SIETHORPIA curopza. Plentiful in the County.
Dr. Sibthorpe. Near St. Austle. Dr.
Maton. Dark hedges and boggy ground
about Lostwithiel; Falmouth; Pensance;
Camelford, &c. D. T.
LEPIDIUM ruderale. About Truro. Ray:
COCHLEARIA officinalis. Rocks at Polperro, and
elsewhere. Mr. E. Forster, jun.
danica. Rocks near Tintagel Castle. Id.
Coronopus didyma. About Truro and Penrhyn.
Hudson Rubbish in the start of D
Hudson. Rubbish in the streets of Pen- rhyn.
CRAMBE maritima. Near Mevagissey. Mr. Watt.
CHEIRANTHUS sinuatus. On the Coast of Corn-
walt. Ray.
riesperis inodora. About Falmouth. Withering
DRASSICA oleracea. About Pensance, and various
other places in Cornwall. Hudson. Be-
tween Fawy and St. Catharine's Tower.
Mr. E. Forster, jun. Rocks at King's
Corre many D.C. I
ERODIUM moschatum. Near Pensance. Rev.
a cubance. Key,

J. T. Thomson. At Portluney and Carhayes. Mr. E. Forster, jun. ERODIUM maritimum. About Pensance. Ray. St. Michael's Mount. Sir T. G. Cullum. Tintagel and Polruan. Mr. E. Forster, jun. Lizard Point. D. T. GERANIUM sanguineum. Kynance Cove. Dr. Maton. About the Soap rocks and the Lizard Point. D. T. LAVATERA arborea. Frequent in this County. Hudson. Godrevy Island, near Portreath ; Mullion-gull rock in St. Ives' Bay, &c. Borlase. GENISTA pilosa. On the Soap Rocks by the Lizard Point. Sir T. G. Cullum. PISUM maritimum. Beach near Pensance. Heath. ANTHYLLIS vulneraria. B. Near Padstow. * D. T. TRIFOLIUM ornithopodioides. About Marazion and Pensance. Hudson. ---- maritimum. Crevices of rocks at Tintagel. Maton. ----- scabrum. On the shore between Pensance and Marazion. Mr. E. Forster, jun. Lorus diffusus. Sands near Pensance. Mr. W. Allen. HYPERICUM androsæmum. Near Pengwarry. Mr. Watt, Very common on moist hedges.

* The plant is in my herbarium with this habitat affixed; but I have forgotten from whom I received it. Certainly neither Mr. Sowerby nor I gathered it. D. T.

Mr. E. Forster, jun. Lanes near Saltash. D. T. TRAGOPOGON porrifolius. In many parts of the County, not uncommon. Dr. Gunthorpe. CARDUUS eriophorus. Near Truro. Withering. SANTOLINA maritima. On the gravelly shore between Pensance, and St. Michael's Mount. Ray. GNAPHALIUM dioicum. Frequent in the County. Hudson. INULA Helenium. Near St. Ives. Id. PYRETHRUM maritimum. On the Coast of this County. Mr. E. Forster, jun. ANTHEMIS nobilis. Common on almost every hill, as well as the banks of rivers. Borlass. CAREX lavigata. In a wood between Boconnoc Parsonage and Crickapit Mill. Mr. E. Forster, jun. LITTORELLA lacustris. Watery lane near Pensance. Mr. Stackbouse. URTICA pilulifera. In a shady ditch at Velinvran. Borlase. OSMUNDA regalis. Near Launceston. Ray. Boggy Vallies at Boconnoc, and elsewhere, not uncommon. Mr. E. Forster, jun. Roadside between Marazion and Pensance. Mr. W. Allen. LYCOPODIUM inundatum. Tor Valley on Bradoc Downs. Mr. E. Ferster, jun,

ASPIDIUM Oreopteris. N	loist places about Bo-
	here. Mr. E. Forster, jun.
spinulosum.	Moist places in the
county.	I Id.
Asplenium marinum. R	ocks at Tintagel. Mr.
	. In a lane near Tre-
vorgy, towards C	arn Brè; and on the
	. Mr. W. Allen. Near
	E. Forster, jun. Li-
	e Treryn, &c. D. T.
TRICHOSTOMUM polypl.ylla	
	where. Mr. E. Forster, jun.
	Temple Moor. Id.
FONTINALIS squamosa, F	
	Mr. Stackhouse.
JUNGERMANNIA Tomentella	
MARCHANTIA conica.	· Iteat Doconnoc.
	Mar E Franken
	Mr. E. Forster, jun.
LICHEN geographicus. R	ocks near the Wring
LICHEN geographicus. R Cheese, on Temple	ocks near the Wring Moor. Mr.E.Forster,
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel.	ocks near the Wring Moor. Mr.E.Forster, D.T.
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel. atro-albus. Stones	ocks near the Wring Moor. Mr.E.Forster, D.T. of a Druid circle at
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel. ————————————————————————————————————	ocks near the Wring Moor. Mr.E.Forster, D.T. of a Druid circle at Mr. E. Forster, jun.
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel. ————————————————————————————————————	ocks near the Wring Moor. Mr.E.Forster, D.T. of a Druid circle at Mr. E. Forster, jun.
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel. ————————————————————————————————————	ocks near the Wring Moor. Mr.E.Forster, D.T. of a Druid circle at Mr. E. Forster, jun.
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel. ————————————————————————————————————	ocks near the Wring Moor. Mr.E.Forster, D.T. of a Druid circle at Mr. E. Forster, jun.
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel. ————————————————————————————————————	ocks near the Wring Moor. Mr.E.Forster, D.T. of a Druid circle at
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel. — atro-albus. Stones Duloe. — varians. Davies. — ferrugineus. — escharoides. Moist	ocks near the Wring Moor. Mr.E.Forster, D.T. of a Druid circle at Mr. E. Forster, jun.
LICHEN geographicus. R Cheese, on Temple jun. At Tintagel. — atro-albus. Stones Duloe. — varians. Davies. — ferrugineus. — escharoides. Moist	ocks near the Wring Moor. Mr.E.Forster, D.T. of a Druid circle at Mr. E. Forster, jun. Plentiful on Stones about the SoapRocks, and Lizard Point. D.T. rocks covered with a

LICHEN niger. Plentifully on rocks and grea	it
stones about St. Ives.	Hudson.
fulvus. Rocks in this County.	Dickson.
aquilus. On Roche Rock; and on Templ	e
Moor. Mr. E. Forster, jun. Rock	5
about the Lizard, Pensance, St. Mi	-
chael's Mount: on St. Michael's Chair	;
and other places in the neighbourhood	,
frequent.	D. T.
conspersus. Rocks on Temple Moor	•
Mr. E. Forster, jun. About Castle	
Carn Brè.	D. T.
J Trees near Boconnoc; and	l
affinis.] Trees near Boconnoc; and Respring, near Lanhydroc.	*
plumbeus. Mr. E. Forster, jun. Near	
Bodmin ; and Liskeard.	
omphalodes. On 'Temple Moor. Mr.	
E. Forster, jun. Common on rocks on	
the mountainous Heaths.	D. T.
scopulorum. Granite rocks of Lemorna	
Cove, near the Land's End. Mr. Knapp.	
Rocks near Fawy: Tintagel; and else-	
where near the Sea. Mr. E. Forster, jun.	
King's Cove; and every where about	
Pensance.	D.T.
pulmonarius. Plentiful about Camelford;	
Launceston, &c. but Mr. Sowerby and	
I found no fruit.	Id.

LICHEN pulmonarius. var. with black shields.
Trees at Boconnoc. Mr. E. Forster. jun.
glomuliferus. Trees in the avenue at Bo-
connoc. Mr. E. Forster, jun. Near
Launceston. D.T.
lælè virens. Trees at Boconnoc.
Mr. E. Forster, jun.
scrobiculatus. Trees at Boconnoc; and
Respring, near Lanhydroc. Mr. E.
Forster, jun. Near Bodmin, and Lis-
keard. D. T.
caperatus. Abundantly in fruit on rocks
near Pensance; and in a Wood between
Camelford and Bodmin. Id.
perlatus. Mr. Sowerby and I found this
Lichen in fruit only in a wood between
Camelford and Bodmin. Id.
miniatus. Rocks near Tintagel Castle.
Mr. E. Forster, jun.
sylvaticus. Rookwood, Boconnoc. Id.
polydactylos. Near Boconnoc. Id.
resupinatus. Rocks at Boconnoc, and
clsewhere. Mr. E. Forster, jun. Trunks
of trees near Liskeard. D. T.
fuliginosus. Rookwood, Boconnoc. Mr.
E. Forster, jun. Plentiful on trees and
moist rocks about Launceston; Bodmin,
&c. D. 7.

LICHEN limbatus. On a tree near the river Fawy,
above Respring. Mr. E. Forster. jun.
polyphyllus. Rocks near the Wring Cheese,
on Temple Moor. Id.
globiferus. Rocks and stones on Temple
Moor, near Arthur's Bed, and the Wring
Cheese. Mr. E. Forster, jun. Stones
near Castle Carn Brè. D. T.
fragilis. Stones at St. Cleere, near Lis-
keard. Id.
paschalis. Stones on Temple Moor be-
tween Arthur's Bed and the Wring
Cheese. Mr. E. Forster, jun. Rocks
by the roadside between Lostwithiel and
St. Austle. D.T.
exilis. Rocks at St. Cleere, near Lis-
keard. Id.
* roccella. Rocks near Tintagel Castle.
Mr. E. Forster. jun.
fuciformis. Rocks near Tintagel Castle. D. T.
articulatus. Trees about Liskeard; and
between Camelford and Bodmin. Id.
* It is not a little singular that in the very spot, where Mr.

Sowerby and I found only *L. fuciformis*, E. B. Mr. Forster should meet with nothing but *L. roccella*, though at the same time both of us gathered specimens in some degree intermediate. This circumstance very much strengthens the opinion, I have long entertained, that these two plants are merely varieties of each other, but the Indian *L. fuciformis*, which is the true plant figured in Dillenius, seems from its texture and habit to be distinct. *D. T.*

LICHEN plicatus. With the former. D.	T.
wulpinus. Apple Trees, frequent. Mr.	
Stackhouse. Near Boconnoc. Mr. E.	
Forster, jun. Rocks at Castle Treryn;	
and Trees about Liskeard and Laun-	
ceston. D.	Τ.
Fucus H. poglossum. At Polruan. Mr. E. Forster, ju	
β . All along the Coast.	
Mr. Stackhou.	se-
Palmetta. On the stems of F. digitatur. Id.	
ovalis. Acton Castle near Pensance, &c.	
Stackhouse. Lizard Point. Mr. E.	
Forster, jun. St. Michael's Mount, and	
elsewhere, not uncommon. D. 2	Γ.
dasyphyllus. Coast of this County. Mr. Stackhous	с.
natans? Among the rejectamenta of the	
bacciferus? Sea at Falmouth. Hudson	12.
ligulatus. Common in deep waters. Mr.	
Stackhouse. Lizard Point; and Ky-	
nance Cove. Mr. E. Forster, jui	2.
siliquosus. B. Near Fawy. Stackhous.	е.
mucronatus. Rocks in this County. Hudson.	
Looe. Mr. E. Forster, jun. Falmouth. D. 7	ی ۵
tamariscifolius. Plentifully near Marazion;	
and elsewhere in the County. Hudson.	
Portluney; Carhayes. Mr. E. Forster, jun.	
Falmouth; and St. Michael's Mount. D. T.	•
$ fibrosus.$ $\beta. Falmouth. Id.$	
β . β . β . β . β . f falmouth. Id.	

.

-

Fucus esculentus. At the Lizard Point : Kynance Cove; and Tintagel Castle. Mr. E. Forster, jun. King's Cove; and St. Michael's Mount. D.T. serratus. y. Coast of the County. Ś wesiculosus. 2, n, 9, e. Mr. Stackhouse. ----- ceranoides. In the Fawy River near St. Winnoe. Mr. E. Forster, jun. ---- laciniatus. Coast of the County. Mr. Stackhouse. Portluney, Carhayes: Pensance: Lizard Point; and elsewhere. Mr. E. Forster, jun. ciliatus. S. Mount's Bay: Pensance; &c. Mr. Stackhouse. King's Cove. D. T. edulis. Menabilly : Fawy : Acton Castle : Pensance, &c. Mr. Stackhouse. Portluney, Carhayes. Mr. E. Forster. jun. ---- bulbosus. Here and there in the County. Hudson. Looe: Fawy; Carhayes. Mr. Stackhouse. St. Michael's Mount. D.T. ---- bifidus. Common on the Coast. Stackhouse. ____ crispus. ζ. Near Marazion. Goodenough. ------ n, ---- mammillosus. y, z. Falmouth. D. T. pygmæus. On the Coast, frequent. Mr. Stackhouse. Rocks at Falmouth. D. T. ----- pinnatifidus. β. & δ. Pensance. Mr. Stackhouse. corneus. Here and there in the County.

Hudson. Acton Castle : Marazion, &c.
Mr. Stackhouse.
Fucus corneus. E, King's Cove. D. T.
* gigartinus. First found in this County
by Hon. Dr. Wenman. On a bank of
Sabella in Mounts Bay, near Newlyn.
Dr. Maccullock.
coronopifolius. All along the Coast. Stack-
house. Looe: Pensance; and elsewhere.
Mr. E. Forster, jun.
plumosus. Common in this County. Stackhouse.
tomentosus. On the long rock between Ma-
razion and Pensance, plentiful. Hon. Dr.
Wenman. St. Ives; and Menabilly.
Mr. Stackhouse. Lizard Point. Mr.
E. Forster, jun. King's Cove. D. T.
tuberculatus. Near St. Ives. Hudson. By
Acton Castle, in pools left by the Tide,
plentiful. Mr. Stackhouse. Lizard
Point. Mr. E. Forster, jun. St. Mi-
chael's Mount. D. T.
rotundus. Pensance. Mr. E. Forster, jun.
y. Falmouth. D. T.

* It has already been pointed out by Dr. Smith in English Botany, that the habitat, given to this plant in the Linnæan Transactions, "copiosissime ad scalas ascensus in ponte marino St. Ives," is a mistake originating in Leofling's imperfect handwriting; and that the place intended is, not St. Ives in Cornwall, but St. Ubes in Portugal.

Fucus conferwoides. G. S. e. Coast of the County.
Mr. Stackhouse.
subfuscus. Pensance: Tintagel; and else-
where. Mr. E. Forster, jun.
pinastroides. Thrown on the Coast in thick
lumps after a storm. Mr. Stackhouse.
capillaris. Submersed rocks, here and
there. Hudson.
kaliformis. Coast of the County. Mr.
Stackhouse. Falmouth: and St. Mi-
chael's Mount. D. T.
asparagoides. Among the rejectamenta of
the sea, very rare. Mr. Stackhouse.
Opuntia. On the naked Rocks at St. Mi-
chael's Mount Pier. Id.
β. St Michael's Mount; and the
adjacent coast. Id.
fruticulosus. Near Acton Castle, Mount's
Bay. Id.
ULVA coccinea. Rocks near Falmouth. Hudson.
fistulosa. In the sea at Looe. Mr. E. Forster, jun.
rubens.)
filiformis. Coast of the County. Mr. Stackhouse.
elminthoides.
dichotoma. Sea near Pensance and Car-
hayes. Mr. E. Forster. jun. A very
narrow variety grows at King's Cove. D. T.
plantaginea. Roth. Rocks at Falmouth.
Mr. G. Sowerby.

CONFERVA villosa. Shores of the County. 2 byssoides. 5 Mr. Stackhouse. fluviatilis. Rivulets near Pensance; where it reaches the length of two or three feet, and is much larger in every other respect than I have seen it in other places. D. T. - parasitica. Here and there on the shore.

Hudson,

THE County of Cumberland, which may justly be called the most alpine district of England, abounds so much in wild rocky situations, contains so great a variety of surface, and has on account of its beautiful scenery so continually been visited by travellers, that it might be expected we should be able to enumerate a catalogue of its rare plants at least equal to that furnished by any other County. This, however, is by no means the case; and our list would have been very deficient had it not been for Mr. Hutchinson's History of Cumberland, from the Flora of which, prepared by the Rev. William Richardson, we have made very copious extracts; and had not our friend Mr. Harriman kindly procured us from the Rev. John Dodd, of Wigton, the habitats followed by his name, and that of Mr. Wood of Whitehaven. A few only of Mr. Hutton's habitats were communicated by himself: the rest were extracted from Mr. Hutchinson's History, were they stand avowedly upon his

authority. Some of the plants are certainly not what we should have looked for in similar situations, so that it is impossible not to suspect that he has fallen into errors; but as this indefatigable Naturalist is a resident at Keswick, and acts as Guide to the Lakes, it will be easy for any Botanist to avail himself of his assistance, and visit the stations of his most rare plants; his readiness to shew which, and indeed to render himself useful by all means in his power, we ourselves experienced on a Tour to the Lakes nine years since.

CHARA flexilis. Ditch near the foot of Keswick Lake. Rev. J. Harriman.

and the second

CIRCAEA alpina. Lorton, on the road between Keswick and Cockermouth. Rev. J. Harriman. Banks of Ullswater. Mr. J. Woods, jun. Bank wood by Chapel Well. Hutchinson. By the borders of Derwent Water, and elsewhere about Keswick. D. T.

VERONICA spicata. · Mr. Salkeld's Cow Pasture, Kirkland. Ilutchinson.

UTRICULARIA minor. Turfy bogs in this County,

Hudson. Near Keswick. Mr. Hutton. In Cooper, Parish of Bromfield. Rev. J. Dodd. SCHOENUS mariscus. Gelt-bridge Farm. Hutchinson. _____ compressus. Hell-beck and Tindale Fell, Brampton. Id. ---- albus. Side of river Gelt. Hutchinson. By Cromack Water. Rev. W. Wood. Common in this County. Mr. Crozue. SCIRPUS acicularis. Tindale Fell. Hutchinson ----- Holoschænus. Maryport. Id. _____ sylvaticus. Bank wood. Id. ERIOPHORUM vaginatum. On all the hills and peat-mosses in low grounds. Id. AGROSTIS Spica venti. Blencarn; and Skirwith: Kirkland; and Langwathby Banks. Id. SESLERIA carulea. Bank-rigg, Stoop-band, on Cross-Fell; Mr. Salkeld's fell pasture, Kirkland; and Tarn House, Brampton. Id. FESTUCA vivipara. Summit of Cross-Fell; Helvellyn; Saddleback; and Skiddaw; and many lower parts of the Fells; islands in Ullswater; and mountains by the sides; and Lattrigg, Keswick. Hutchinson. Scale Force. Rev. W. Wood. BROMUS arvensis. Road from Brampton to Naworth-Castle: Walton House, by the garden. Hutchinson. VOL. I. L

BROMUS pinnatus. Woods at Kirk-house: Bank-	
wood by Chapel Well. Hutch	insona
STIPA pennata. Said to grow on hills between	
Ullswater and Hawswater, but we have in	
vain sought for it.	Id.
ARUNDO epigejos. Dunmallet, by front avenue,	
rare.	Id.
calamagrostis. Dalemain, and Kirkland	
woods.	Id.
ELYMUS arenarius. Abbey Holm; and Allonby.	Id.
TRITICUM loliaceum. In Barley crops, Abbey	
Holm. Rev. J.	
GALIUM tricorne. Edenhall. Hutc	hinson.
boreale. Ullswater; and Keswick.	
Hutchinson. Near Wastdale-head. Mr.	
Wood. Near Pooley Bridge, by Ulls-	
water. Mr. Woodward. Hedges near	
Penrith, common. Mr. Har	riman.
pusillum. Common in this County.	
· Mr. Wood	
I S S S S S S S S S S S S S S S S S S S	Hutton.
* EPIMEDIUM alpinum. On Saddleback, near	
Threlkeld.	Id.
ALCHEMILLA alpina. Mountains at the head of	

* We had never heard of any other Botanist who had been fortunate enough to meet with this plant, till we were lately assured by Mr. Rudge, that he had received a specimen from a Lady, who gathered it herself in the above habitat.

Ullswater. Mr. Brunton. Rocks in Borrowdale. Mr. Woodward. Mountain at the head of Ennerdale. Mr. Fardon. Swarth-beck Gill, Ullswater; and Rosthwaite, by the road to Buttermere Lake. Hutchinson. In St. John's, in the neighbourhood of Keswick, on the sides of the Fells above Hill-top; also on the Fells near Buttermere.

Rev. J. Harriman. POTAMOGETON gramineum. Keswick. Mr. Hutton. _____ setaceum. Peat moss by Talkin tarn. Hutchinson. LITHOSPERMUM purpuro-caruleum. , Wood at Castle-carrock. Id. PULMONARIA officinalis. Keswick. Mr. Hutton. maritima. Near Whitehaven. Ray. Here and there between Maryport and Worthington. Rev. J. Harriman. Abbey Holm. Hutchinson. Near Workington. Rev. J. Dodd. Coast between Ravenglass and Bootle. Mr. Wood. PRIMULA farinosa. Kirkland, wettish ground, frequent. Hutchinson. Near Aspatria, where a white variety is also found. Rev. J. Dodd. Common in mountainous Pastures. Rev. J. Harriman. L 2

MENYANTHES nymphæoides. Keswick. Mr. Hutton. Lysimachia thrysiflora. CAMPANULA patula. ---- latifolia. Woods at Newbiggin: by Hutton John; and Highgate, Graystock. Hutchinson. Isell. Rev. 7. Dodd. Common in this County. Rev. 7. Harriman. LOBELIA dortmanna. Ullswater: Keswick: and Talkin-tarn. Hutchinson. Tarn in Wast-Mr. Wood. dale. IMPATIENS noli-tangere. Keswick. Mr. Hutton. Mr. 7. Woods, jun. At Scale Hill. VERBASCUM Blattaria. Aspatria Church Yard. Rev. 7. Dodd. DATURA Stramonium. Wallow Crag, Keswick. Mr. Hutton. GENTIANA pneumonanthe. Field between Maryport and Flimby, 200 or 300 yards from the latter. Rev. J. Harriman. Howgill Hutchinson. castle woods. ---- campestris. Mr. Salkeld's Grounds; and side of the river Gelt. Hu chinson. Fields adjoining Black Cock, near Cockermouth. Rev. J. Dodd. Near the rope-walk, Maryport. Rev. J. Harriman. Mr. Hutton. ATHAMANTA Libanotis. Keswick. MEUM athamanticum. Near the Vicarage at Kes-Rev. W. Wood. wick.

SIUM latifolium. Keswick. Mr. Hutton. ----- verticillatum. Near Ramsey, Isle of Man. Rev. H. Davies. (Who here first added it to the British Flora.) CICUTA virosa. Keswick : banks of the Irthing at Walton, and Irthington. Hutchinson SCANDIX odorata. Common in Orchards. Hutchinson. About Keswick. D. T. PIMPINELLA dioica. Keswick. Mr. Hutton. SAMBUCUS Ebulus. Alston Moor. Rev. J. Harriman. Very near Aspatria, and only in one field on the East side of the Town. Rev. 7. Dodd. STATICE reticulata. Whitehaven. Id. DROSERA longifolia. Keswick; and Mr. Carlyle's, Culgaith Moor, rare. Hutchinson. anglica. Keswick, with the preceding, rare: near Talkin Tarn, and Unity Farm, Brampton. Hutchinson. Three miles from Carlisle towards Scotland. Ray. NARCISSUS poeticus. Keswick. Mr. Hutton. ALLIUM ampeloprasum. * House-holm Island, Ullswater; and Keswick. Hutchinson. arenarium. Lowdore Water Fall. Sir

* Do not these two habitats belong to the following species, which, however, Hutchinson also enumerates, and assigns as a station "Buckam wood, Lowther, Westmoreland ?"

T. Frankland. Borders of Ullswater.

Mr. Jos. Woods, jun. ALLIUM oleraceum. Borders of Derwent water. D. T. FRITILLARIA Meleagris. ORNITHOGALUM luteum. Keswick'. Mr. Hutton. umbellatum. CONVALLARIA Polygonatum. JUNCUS filiformis. Margin of Derwent water. Mr. Dickson. _____ triglumis. Rocks above Borrowdale. Mr. Hutton. By Lowes water. Mr. Wood. RUMEX digynus. By black-lead mine, Borrowdale. Mr. Hutton. In a ravine of the Screes Mr. Wood. near Wastwater. ALISMA natans. In Keswick Lake. Rt. Hon. C. Grewille. ranunculoides. Patterdale; and Waverton. Hutchinson. Aspatria Moss. Rev. J. Dodd. EPILOBIUM angustifolium. Banks of Eden, Hutchinson. Corby. _____ alpinum. Keswick; and Gowbarrow Park by Ullswater. Id. CHLORA perfoliata. Spade Adam, near the house. Id. VACCINIUM uliginosum. Cross Fell; Hartside Fell, near Kirkhouse; Talkin Tarn. Hutchinson. At Gamblesby, six miles from Penrith. Ray. Alston Moor. Rev. 7. Harriman. ---- Vitis idea. Keswick ; and Waverton. Hutchinson. On Skiddaw. Rev. J.

Dodd. Upon Great Gable, in the West side of the County. Mr. Word. VACCINIUM Oxycoccos. Bromfield. Rev. J. Dodd. Colgaith Moor : Tarn Waddalyne, Hesket. (A bog here has produced £ 20 worth of these berries in one year.) Hutchinson. Common in the peat Bogs. Rev. J. Harriman. PARIS quadrifolia. Bank Wood: Naworth Woods: and roadside between Hutton Moor and Penrith. Hutchinson. Isell Wood, near the river. Rev. 7. Dodd. ANDROMEDA polifolia. Kirkland: Cross Fell: Keswick: and Brampton. Hutchinson. Moss near Bromfield. Rev. 7. Dodd. ARBUTUS Uva Ursi. Martindale Dale-Head, Ullswater. Hutchinson. PYROLA minor. Dunmallet, foot of the Avenue fronting Ullswater. Id. secunda. } Near Keswick. Mr. Hutton. rotundifolia. * SAXIFRAGA stellaris. Cross Fell: Patterdale:

* The difference occasioned in this plant by its place of growth is very striking. Specimens I gathered in the valley near Keswick Lake were nearly a foot high, with leaves above an inch long, and panicled stems of many flowers, while those on the very summit of Snowdon did not rise to an inch in height, and were in all parts proportionably small, except the flower, which was single, and considerably larger than those of the vailey. D. T.

Swarth Fell: Place Fell: Kirkstone; and Keswick. Hutchinson. On Skiddaw, plentiful. D.T. SAXIFRAGA nivalis. Gowbarrow Park: and Keswick. Hatchinson. ---- aizoides. Patterdale : Gilsland. near Spawell; and Keswick. Hutchinson. On the moist parts of Skiddaw, and the adjacent Mountains, frequent. D. T. On the Fell, between Borrowdale and Watendlath. Rev. 7. Harriman. oppositifolia. In a ravine of the Screes, near Wastwater, about 600 feet in perpendicular height; with S. stellaris, aizoides, and hypnoides. Mr. Wood. --- caspitosa. Kirkstone; and Keswick. Hutchinson. meschata. Moist rocks, near Keswick. Mr. Hutton. ---- hypnoides. Swarth-Beck, and Gowbarrow, Ullswater : Keswick. Hutchinson. SAPONARIA officinalis. Usemire, and Howtown, Ullswater; and Ake-beck-bridge, by Pooley. Hutchinson. On a hedge in Aspatria Village, no garden near. Rev. 7. Dodd. Mr Hutton. DIANTHUS deltoides. B. Keswick. STELLARIA nemorum. Cooms Wood; and Dunmallet. Hutchinson. By Aspatria Mill. Rev. 7. Dodd,

ARENARIA verna. y. Hallen-hag: Swarth-Fell; and Place Fell, Ullswater. Hutchinson. SEDUM dasyphyllum. Keswick. Mr. Hutton. ----- rupestre. ----- villesum. By Mill Stream, Melmerby Moor; and Hartside Fell. Hutchinson. OXALIS acetosella. B. Near Keswick. Mr. Hutton. ASARUM eurspæum. Ramskin, Martindale; and Keswick. Hutchinson. PRUNUS Padus. Frequent in this County, Hudson Aspatria. Rev. 7. Dodd. ROSA villosa. Plentiful in this County. Ray. Near Ullswater. Mr. Woodward. Pooley. Hutchinson Rubus chamamorus. On Cross Fell, abundant: from Kirkland to Tindale Fell; and How town, Ullswater. Id. POTENTILLA verna. Bank Wood. Id. GEUM rivale. Kirkland; and Brampton. Hutchinson. Roadside between Penrith and Keswick. Mr. Fardon. ACTÆA spicata.. Sandwicke, Ullswater. Hutchinson. PAPAVER cambricum. Keswick. Mr. Hutton. CISTUS marifolius. Stybrow; and Blowicke, Ullswater. Hutchinson AquileGIA vulgaris. B. Near Kes-ANEMONE nemorosa. var. flore purpureo. wick. THALICTRUM alpinum. Mr. Hutton.

THALICTRUM minus. Keswick. Hutchinson. Wastdale Head. Mr. Wood. Margin of Ullswater, near Gowbarrow Park. D. T. majus. Near Isell Church, 2 single root. Rev. J. Dodd. On the margin of Ullswater. Mr. Robson. In a ravine of the Screes near Wastwater. Mr. Wood. RANUNCULUS Lingua. Emont, by Carleton Hall, Hutchinson. Abbey Holm. Rev. 7. Dodd. Ditches in Low Holm Mire. Rev. J. Harriman. TROLLIUS europæus. Kirkland, abundant in moist woods, meadows, &c. Hutchinson. As-Rev. 7. Dodd. patria Mill. HELLEBORUS viridis. Threepland Gill. Id. fætidus. Keswick. Mr. Hutton. LEONURUS Cardiaca. Langrigg, Broomfield. Hutchinson. SCUTELLARIA minor. Culgaith Moor; and Kes-Id. wick. PRUNELLA vulgaris. var. fl. albo. Aspatria Moss. Rev. 7. Dodd. MENTHA Pulegium. In Low Holm Mire. Rev. J. Harriman.

* MELAMPYRUM sylvaticum. Near Scale Hill.
Mr. J. Woods, jut
LATHRÆA squamaria. In a field of Mr. Rede's,
Hole-house, Baggray. Hutchinson. Near
Akebank, in the Parish of Wigton.
Rev. J. Doda
SIBTHORPIA europæa. Glencoin, and Gowbarrow
Park, by Airey Force; and Keswick.
Hutchinson
THLASPI arvense. Isle of Man, but I do not re-
collect the spot. Rev. H. Davies
perfoliatum. On most limestone pastures
in this County.
alpestre. Mr. Salkeld's fell pasture;
and Bank-rigg, Kirkland. Hutchinson.
COCHLEARIA officinalis. Among rocks, Cross
Fell. Id.
danica. Coast between Ravenglass
and Bootle. Mr. Wcod.
IBERIS nudicaulis. Between Corby Castle and
Carlisle. Dr. Stokes.
CRAMBE maritima. Coast between Ravenglass

* When I was in this County, in 1796, I remarked in the woods two very remarkable, and apparently diffind varieties of M. pratense; one with deep yellow, the other very pale flowers, but I unfortunately omitted to note down the differences, and I have since lost my specimens, so I can only request the attention of other Botanists to the subject. D.7.

and Bootle. M . Wood. Between Maryport and Flimby. Rev. 7. Harriman. CARDAMIME impatiens. Not unfrequent in this County. Mr. Woodward. Keswick. Mr. Hutton. amara. Gelt river, Irthing; and Maryport. Hatchinson. Fitz, near Aspatria. Rev. 7. Dodd. SISYMBRIUM sylvestre. Keswick. Mr. Hutton. monense. Found by Mr. Lawson in Sella fields, Seabank. Ray. On the Bent Hills, Maryport. Rev J. Harriman.' Allonby. Rov. 7. Dodd. Near Peel, and Point of Eyre, Isle of Man. Rev. H. Davies. HESPERIS inodora. Banks of rivulets about Dale Head. Ray. ARABIS stricta. Kirkland Fell pasture. Hutchinson. ----- bispida. Ravine of the Screes, near Wastwater, about 600 feet in perpendicular height. Mr. Wood. ERODIUM moschatum. Hedge bank at Mr. Yates's, Skirwith, Kirkland. Hutchinson. ---- maritimum Sea shore in the Isle of Robson. Man. GERANIUM phæum. Kirkland and Keswick. Hutchinson.

GERANIUM * nodosum. Mountainous situations in	
Cumberland.	Ray.
Hudson. About Corby Castle. Dr.	Stokes.
pyrenaicum. Skirwith; and Keswick.	
Hutc	hinson,
pratense. Common in this County.	
Rev. J. Har	riman.
sanguineum. Near the road leading	
from Allonby to Maryport. Rev. J.	
Dodd. Common on the shore near Mary-	
port. Rev. J. Harr	iman.
FUMARIA solida. At John Johnson's Esq. Wal-	
ton House, by farm yard. Hutchinson.	
Catsteads, near Brampton. Rev. J. 1	Dodda
OROBUS sylvaticus. Hedges and pastures, plenti-	
fully, at Gamblesby, six miles from Pen-	
rith, in the way to Newcastle. Ray.	
Under Cross Fell: from Kirkland to Gam-	
blesby. Hutch	insono
+ LATHYRUS sylvestris. On the rocks by Red	
Neese, near Whitehaven,	Raÿ.

• Dr. Withering's observation, that Mr. Woodward has since found it, is erroneous, as that Gentleman never met with it, and merely pointed out Ray's habitat to his notice.

† The plant here found was by Ray, and has continued to be by all subsequent writers, called *L. latifolius*, but Rev. J. Harriman detected the error and has been so good as to favor us with specimens.

VICIA sylvatica. Culgaith woods, by Eden; and Keswick. Hutchinson. At Gamblesby, Mr. P. Harrison. Isell woods, where it covers the rocks in most beautiful festoons. Rev. J. Dodd. ----- Intea. ASTRAGALUS bypoglottis. Keswick. Mr. Hutton. glycyphyllos. Culgaith-pie: Keswick. Hutchinson, TRAGOPOGON porrifolius. Fields about Carlisle; and Rose Castle. Robson. Said to grow ncar Rose Castle, where we have sought for it in vain. Hutchinson HIERACIUM alpinum. Keswick. Mr. Hutton. dubium. Keswick; and Patter-Auricula. I dale. Hutchinson paludosum. Kirkland : Ullswater : Keswick. Hutchinson. Common about the Falls in this County. Mr. J. Woods, jun. HYPOCHÆRIS maculata. Keswick. Mr. Hutton. SERRATULA alpina. Castle, and near Church, Bewcastle. Hutchinson CARDUUS eriophorus. Bewcastle. Id. ----- heterophyllus. Kirkland. Id. GNAPHALIUM dioicum. Kirkland : Brampton; and Penrith Fell. Hutchinson. Ravine of the Screes near Wastwater. Mr. Wood. sylvaticum. Brampton. Hutchinson.

SENECIO viscosus. Nunnery walks: Brampton:
Keswick. Hutchinson,
- saracenicus. Very plentifully at Salkeld.
Ray. In the hedges at Long Town,
and in the side of the river below Carlisle,
Mr. Jackson. Newby Bridge. Mr.
J. Woods, jun. Near Ullswater. Mr. Brunton.
Solidago Virgaurea. y. Mr. Salkeld's fell pas-
ture. Hutchinson.
ORCHIS ustulata. Blencarn; and Keswick. Hut-
chinson. Blinderake, parish of Isell.
SATYRIUM bircinum.
repens. J Loo when it is
albidum. Keswick. Mr. Hutton.
Mountainous pastures aboye Borrowdale. D. T. OPHRYS Nidus avis. Keswick. Mr. Hutton.
OPHRYS Nidus avis. Keswick. Mr. Hutton.
cordata. Kirkland. Hutchinson. On Snafield Isla of Mar
Snafield, Isle of Man. Rev. H. Davies. CYPRIPEDIUM Calceolus. Warm shade near Kes-
wick
SERAPIAS longifolia. Banks of Waver. Hutchin-
CAREX dioica. } Brampton. Hutchinson.
Id.
- ampullacea. Tindale Tarn: Eel stank,
hy Pooler

SPARGANIUM natans. Old Eden. Hutchinson LITTORELLA lacustris. Ullswater : Cullgaith Moor, School, and Mr. Carlyle's Land. Hutchinson. Low Holm Mire. Rev. 7 Harriman. At Tyne Head; and in many places at Alston; Gray-SALIX pentandra. ---- reticulata. stock; Abbey Holm, &c. Hutchinson. ---- herbacea. Summit of Skiddaw, and Saddleback. Id. - cinerea. There are specimens of this plant in Lightfoot's Herbarium from Cumberland. Smith. EMPETRUM nigrum. Cross Fell: Kirkhouse; and Brampton. Hutchinson. On Skiddaw. D. T. RHODIOLA rosea. Keswick. Mr. Hutton. Ravine of the Screes, near Wastwater. Mr. Wood. JUNIPERUS communis. B. Near Stye-head. Id. TAXUS baccata. Ullswater : undoubtedly indigenous, as the Yew has been found in peat mosses in Matterdale and Patterdale : we have seen large pieces of a beautiful dark Hutchinson. red colour. ATRIPLEX laciniata. Allonby. Rev. 7. Dodd. Shore near Maryport. Rev. J. Harriman. EQUISETUM fluviatile. Old Eden; Banks of Emont; and Keswick. Hutchinson.

151

EQUISETUM hyemale. Low-gelt-bridge. Hutchinson. OSMUNDA lunaria. Keswick, in a field on the right hand of the road to Penrith : How Hill: Castle Sowerby; and Naworth Park, rare. Hutchinson. In a field of Mr. Allison's near Aspatria, Rev. J. Dodd. Sea dike between Workington and the Coin House. Rev. J. Harriman. ---- regalis. Low-gelt-bridge; and Keswick. Hutchinson. Allonby Moss. Rev. J. Dodd. LYCOPODIUM selaginoides. Frequent on Hills. Hutchinson. alpinum. Place Fell; and Swarth Fell. Id. POLYPODIUM fontanum. Keswick; and Saddleback. Id. ----- Phegopteris. Airey-beck, Glencoin : Lodore, Keswick. Hutchinson. Plentiful in Borrowdale, and under Bowder-Stone. D. T. ---- Dryopteris. Cross Fell, Ullswater; and Keswick. Hutchinson. Not uncommon in Borrowdale. D. T. Rocks at Lowdore. Mr. Woodward. Aspidium Lenchitis. Keswick. Mr. Hutton. ----- Thelypteris. Glencoin, and Blowike, Ullswater; and Keswick. Hutchinson, ------ Oreopteris. Ullswater; Cross Fell: VOL. I. M

Keswick. Hutchinson. Near Aspatria. Rev. J. Dodd. Frequent among the waterfalls, particularly at Cromack and Buttermere. D.T.ASPLENIUM viride. Place Fell, and Swarth Fell, Ullswater : Cross Fell Edge; and Keswick. Hutchinson. --- marinum. Whitehaven; and Maryport. Id. ---- septentrionale. Patterdale ; and Keswick. Hutchinson. Ravine of the Screes near Wastwater, about 600 feet in perpendicular height. Mr. Wood. SCOLOPENDRIUM Ceterach. Dry rocks near Keswick. Mr. Hutton. PTERIS crispa. Frequent in Martindale, &c. Hutchinson. Very common about Borrowdale. Milin. D. T.HYMENOPHYLLUM tunbrigense: Gowbarrow, and Keswick. Hutchinson. Ravine of the Screes near Wastwater. Mr. Wood. Scale Force. Rev. W. Wood. ISORTES lacustris. In Derwent Water. Mr. Woodward. Ullswater, lower end; and Gowbarrow-wike. Hutchinson. GYMNOSTOMUM curvirostrum, Rocks in this Dickson. County.

*POLYTRICHUM urnigerum.	
NECKERA crispa. Near Keswick. Mr.	Hulton.
pumila. Barton Park, Ullswater. Hute	binson.
BARTRAMIA arcuata. Near Keswick. Mr.	Hutton.
FONTINALIS squamosa. Swarth-beck. Hute	binson.
JUNGERMANNIA ciliaris. Woods and moist	
Heaths; and near rivulets.	Judeon
LICHEN geographicus. Common in the mountain-	<i>L </i>
ous parts of this County. Rev. J. Har-	
Timan. Rocks about V and 1	
riman. Rocks about Keswick.	D. T.
atro-rufus. On a Hill near Penrith,	
growing on the sandy ground among	
mosses. Mr. Dickson. Near St. John's	
Chapel in the neighbourhood of Keswick.	
Rev. J. Harr	iman,
ventosus. Common in mountainous si-	
tuations	Id.
coccineus. I have found this in Cumber-	
and, but forget when	Id.
Oederi. In the neighbourhood of Keswick.	ι <i>ι</i> , τ ,
tartareus. Common in the mountainous	ld _a
Conspersus in and	
parts. 1	'd.
* Our knowledge of the cryptogamic productions of the m	

bourhood of Keswick, which stands on the authority of Mr. Hutton, is derived almost wholly from a box of Lichens and mosses, which at the request of a friend he sent to me, and in which the species here enumerated were included, though chiefly with very wrong names. D.T.

LICHEN	* quadricolor. In the neighbourhood of	f
	Keswick. Rev. J. Ha	arriman.
	gelidus. In Borrowdale.	Id.
-	† cæsio-rufus. On a sand stone wall be	-
	tween Bankend and the salt pans in th	e
	neighbourhood of Maryport.	Id.
	pruinatus. Limestone walls between	n
	Penrith and Penruddock.	Id.
	Westringii. Rocks in Thornthwaite	г,
	near Keswick.	Id.
	squamatus. Rocks in this County. Mr.	Dickson.
	‡ carnosus. Near St. John's Chapel i	n
	the neighbourhood of Keswick.	
	Rev. J. H	arriman.

bourhood of Keswick. (I have seen spe-

* " I sent specimens of this to Acharius, which he suspects to be a variety of his *Lecidea granulosa*, and yet it is not of the same genus." *Mr. Harriman*.

† "I do not think this distinct from *L. cinereo-fuscus*. Ach. Prod." Mr. Harriman.—I must dissent from this opinion, for *L. cinereo-fus*cus, according to authentic specimens, is only *L. ferrugineus*. D. T

‡ " I sent Acharius a specimen of this, and he thinks it a variety of his Lecidea microphylla: he certainly, however, is wrong." Mr. Harriman.—Unquestionably so: for his L. microphylla, on the authority of specimens from himself and Dr. Swartz, is Lichen escharoides. E. B. t. 1247; the error lies in his having quoted Schrader's L. microphyllus, which really is our L. carnosus, as a synonym to his plant of the same name. D. T.

cimens of this that approached very near,
at least, to L. encaustus.) Rev. J. Harriman.
LICHEN pulmonarius. glomuliferus. latè-virens. scrobiculatus. Common near Keswick; and glomuliferusisgene- rally in fruit : I never found lætè-virens when it was not so. Id.
trees in mountainous situations, but I never found it in fruit in this County. Id.
<pre>plumbeus plumbeus scopulorum proboscideus Near Keswick. Mr. Hutton.</pre>
- pellitus. Dry rocks near Keswick. Id.
islandicus. Upon Great Gable on the
west side of the County. Rev. W. Wood.
vermicularis. Near the summits of moun-
tains. Mr. Hutton.
sterilis. Ach. } Near Keswick. Id.
Fucus esculentus. Coast of this County. Hudson.
Isle of Man. Rev. H. Davies.
CONFERVA fluviatilis. Rivers in this County. Hudson.

DENBIGHSHIRE.

IT may be observed that we have frequently in our list of this County, as well as occasionally in Carnarvonfhire and Flintshire, copied articles verbatim from Mr. Bingley's Tour, and instead of his name subjoined that of Mr. Griffith. In all these cases the habitats were originally communicated by this gentleman, though inserted without an acknowledgment; and we have considered that we should best discharge our duty to the public by referring them to the original source, as it is impossible for any information to be so authentic, as that derived from those Naturalists who have long been resident in a county, and bestowed a continued application to the subject. Upon the Botany of Denbighshire we are not aware that any thing has yet been published, nor much observed, unless by Mr. Griffith : its features however bear so strong a resemblance to those of North Wales in general, that, excepting the few species which are peculiar to the highest Mountains, it may fairly be expected that its vegetable productions should not materially differ from those of the adjoining counties.

DENBIGHSHIRE.

CHARA flexilis. In a pool, called Llyn Aled, in
the Parish of Llanfannon, where I have
seen it many yards in length. Mr. Griffith
VERONICA montana. Near the rivulet in Garn
Dingle, three miles Westward of Den-
bigh: as common in woods about Garn
as V. Chamædrys. Id.
IRIS fatidissima. Woods and hedges about the
Farm House of Pen ucha'r Green, in the
parish of Henllan. Id.
SCHOENUS albus. By the side of the old road from
Capel Cerig to Llanrwst in an inclosure
above Pencraig House. Id.
ERIOPHORUM vaginatum. Peat bogs at Llyn
Aled, and on all the mountains. Id.
polystachion. With the former, but
less frequent. Bingley.
MILIUM lendigerum. Pastures East of Merllin
Farm House, in the Parish of Llanyfydd.
Mr. Griffith.
MELICA nutans. Garn Dingle. Id.
FESTUCA vivipara. Mountainous pastures. Id.
bromoides. About Denbigh castle. Bingley.
BROMUS arvensis. Not uncommon in Wales.
Mr. Griffith.
AVENA fatua. Neighbourhood of Llanrwst. Id.
pubescens. About Garn, as common as
A. flavescens. Id.

ELYMUS europaus. Garn Dingle, under Garreg wen rocks. Mr. Griffith. DIPSACUS pilosus. In the hedge of the meadow on the South side of the River Elwy, belonging to Mr. Lloyd of Wygfair above Pont y rallt goch, S. E. of his house adjoining the Hanging Wood. Id. CENIUNCULUS minimus. In a piece of moist ground about a mile from Llanwrst, and within three or four yards of the Turnpike road leading from thence to Conway. Id. LITHOSPERMUM purpuro-caruleum. On the top of a bushy hill North of Denbigh. Ray. It certainly does not grow there. Mr. Griffith. CAMPANULA latifolia. About Eglwyseg, and Llangollen. Waring. ----- bederacea. By the roadsides near Llanrwst; and near the Cataract Rhaiadr v Wenol, five miles from Llanrwst. Mr. Griffith. VIOLA hirta. Very common about Garn. Id. ------ lutea. Between Llanrwst and Denbigh; and about Cerig y Druidion. Mr. Griffith. Pretty Plentifully on the moun. tains about Llanarmon Dyffrion Ceiriog; and sparingly about midway between Ruthin and Llangollen. Waring. VERBASCUM Lychnitis. The variety B, or y,

(V. thapsoides Huds.) has frequently

been produced in my garden by the farina of V. Thapsus falling upon the stigma of V. Lychnitis. This plant is strictly an hybrid, the seeds, though good in appearance, never vegetating. Mr. Griffith.

RHAMNUS catharticus. Garreg wen rocks by Garn.

VINCA major. On the Common near Rhyd y cilgwyn bridge, between Denbigh and Ruthin.

GENTIANA campestris. In the Coppice, but usually called Coppy, between Garn and Denbigh, and very common in most mountainous pastures in Wales.

SCANDIX odorata. About New Hall, and Chirk Castle. Mr. Griffith. Plentifully in the Vale near the Abbey of Valle Crucis, and among the ruins. Waring.

SAMBUCUS Ebulus. Between Llansannan Church and the river. Mr. Griffith.

VACCINIUM uliginosum. In the higher and moist parts of Gwydir Woods near Llanrwst. Aikin.

 Oxycoccos. Bogs, very common. Mr. Griffith.
 CHLORA perfoliata. About Abergele. Id.
 POLYGONUM Bistorta. In a trench without the Western wall of the Castle of Ruthin. Waring.
 PARIS quadrifolia. Among the thickets on the North side of Garreg wen rocks, and on

the opposite side of the rivulet. Mr. Griffith.

I.

Id.

Id.

DENBIGHSHIRE.

ANDROMEDA polifolia. In a morassy flat about a	
mile N. E. of Llyn Aled, called Gors	
y Casseg: in another place about the	
same distance West of the Pool; and	
South end of a large boggy flat, called	
Waen y dalfa, also near Llyn Aled.	
	Tuite 1
SAPONARIA officinalis. Denbigh Castle, between	Frifith.
the Church and the Town; also on the	
left of the road as you enter the Town of	
Ruthin from Denbigh.	Id.
SEDUM dasyphyllum. ZEast side of Denbigh	10.
album. Castle.	7.7
rupestre. On a wall South of Gwydir	Id.
Chapel near Llanrwst.	
	Id.
SPERGULA subulata. In dry pastures above Deû-	
nant, between Llanfannan and Denbigh;	
and on the first Common on the ascent	
of the hill from Nant Glyn towards	
Henllan near Denbigh.	Id.
PRUNUS Padus. About Chirk.	Id.
PYRUS torminalis. Very common about Garn,	
attaining the height of 20 or 25 feet.	Id.
* hybrida. On the walls of Castell Dinas	
Brân. Mr. W	aring.

* " The plant on the walls of Castell Dinas Brân, I consider as 2.mere variety of *Pyrus Aria*, not a distinct species, and certainly not *P. bybrida*, of Linnæus; the leaves being only lobated, and not

Rosa villosa. Very common all over Wales. ---- tomentosa. Mr. Griffith. RUBUS saxatilis. Garreg wen Rocks near Garn, abundantly. Id. TORMENTILLA reptans. Sandy places about Llanrwst. Bingley. GEUM rivale. Wood below Garn. Mr. Griffith. Among the rocks of the Waterfall called Rhaiadr Mawr between Conway and Llanrwst. Bingley. PAPAVER cambricum. Near Pont Meredith. Mr. Griffith. RANUNCULUS gramineus. Said to have been found in the neighbourhood of Llanrwst, but late Botanists have sought for it in vain. Bingley. (Dr. Withering speaks as if he had seen specimens brought from North Wales by Mr. Pritchard.) parviflorus. About Garn. Mr. Griffith. TROLLIUS europaus. Frequent in this County. Waring. HELLEBORUS viridis. Wood at Plasnewydd near Garn.

Mr. Griffith.

pinnatifid. Mr. Waring about fourteen years ago shewed me specimens, gathered from the same place, in which the leaves were completely pinnatifid, and I have engrafted upon *P. aucuparia* scions of this, the leaves of which become more pinnatifid every year, but I have not yet been able to obtain seeds from my plants." *Mr. Griffitb.*

HELLEBORUS fætidus. In Park Pierce, and the Crest	
The second secon	Griffith.
TEUCRIUM Chamædrys. In the N. W. hedge of	
the bowling green on Ruthin Castle,	
where it has grown many years.	Id.
MENTHA acutifolia. By the roadside between	
Cotton Hall, and Whitchurch, and	
Denbigh.	Id.
gentilis. Pools and brooks betwixt	
Mold and Llanrwst.	Sole.
GALEOPSIS versicolor. In Cornfields about Chirk;	
and in many small inclosures by the road-	
side leading from Chirk to Llangollen.	
Mr. (Griffith.
THYMUS Nepeta. Denbigh Castle; and Graig	
near Denbigh.	Id.
LATHRÆA squamaria. In shady places close to	
the rivulet, on the West side of Garreg wen Rocks, near Garn.	~ 4
ANTIRRHINUM Cymbalaria. Brynkynallt, on	Id.
.5 1. 1 11	, ,
minus. Cornfields near Abergele. Mr. G	aring.
SUBULARIA aquatica. Llyn Aled.	Id.
LEPIDIUM latifolium. In Denbigh Castle, near	LU.
the Gateway that leads down to High-	
gate.	Id.
THLASPI alpestre. By the side of a Dingle,	
called Nant Bwlch yr Hiarn, about a	
mile from Llanrwst bridge, and not more	

DENBIGHSHIRE.

than twenty yards from the Turnpike road leading to Conway. Mr. Griffith. IBERIS nudicaulis. By the side of the hill by the road leading from Corwen to Llangollen about half a mile from Corwen; and on the banks of the road about midway between Cerig y Druidion and Denbigh. Id. CARDAMINE bellidifolia. Various places in Denbighshire. Merrett. Not to be found near Denbigh; but in the place where it is represented to have grown there is abundance of a stunted variety of Turritis hirsuta, not Arabis stricta, as stated in Withering. Id. ERYSIMUM cheiranthoides. Near Tan y Llan in the Parish of Llanyfydd, about five miles N.W. of Denbigh. Mr. Griffith. By the roadside between Talycafn and Bettws. Rev. H. Davies. TURRITIS birsuta. On Garreg wen Rocks near Garn. Mr. Griffith. ERODIUM maritimum. Between Abergele and the sea. Id. GERANIUM pratense. On the bank at Clomendy in Llanveras; and in the Churchyard of Llanveras. Waring. rotundifolium. Rough and sterile places near Garn. Mr. Griffith,

GERANIUM sanguineum. Cefn Rocks near Garn.	
Mr. (Griffith.
OROBUS sylvaticus. About Voylas; and Cerig y	
Druidion.	Id.
VICIA sylvatica. On the S. W. side of Garreg	
wen Rocks, close to the wall that divides	
the Garn demesne from that of Gallt-	
faynan.	Id.
sativa. y. Rocks in the front of Garn.	Id.
ASTRAGALUS glycyphyllos. Hills about Yspytty.	II.
TRIFOLIUM ornithopodioides. On Henllan Hill	
near Garn.	Id.
scabrum. Hill of Denbigh Castle.	D. T.
glomeratum. Rocks about Garn.	
Mr. G	riffith.
HYPERICUM androsæmum. Merllyn Dingle near	
Garn, between the Wash House and the	
Mill.	Id.
dubium. About Garn, but not very	
common,	Id.
HIERACIUM paludosum. Near Garthewin.	
Rev. H. L	avies.
CARDUUS tenuiflorus. Cefn Rocks; and about	
Denbigh Castle. Mr. G	Friffith.
GNAPHALIUM rectum. On Windy Bank Hill,	
near Denbigh.	Id.
INULA Helenium. Road side between St. Asaph	
and Denbigh,	Id.

CENTAUREA solstitialis. I have observed this latel	v
in a field by the Green House at Garn	
but suppose it must have been introduced	
with the seeds of Saintfoin or Lucern	
with which that field has been sown. Mr.	
CAREX curta. Near Wrexham.	Ray.
pendula. Near the rivulet on the West side	
	Griffith.
strigosa. In Garn Dingle, in a wet place	
at the bottom of the Glade.	Id.
Pseudo-cyperus. In the Pool in a field,	
called Marlas pitia bâch, near Garn.	Id.
binervis. On all the Moors of Wales.	Id.
LITTORELLA lacustris. Shores of all Alpine lakes	. Id.
QUERCUS sessiliflora. Very common in Wales.	Id.
TAXUS baccata. Cefn Rocks.	Id.
EQUISETUM sylvaticum. By the roadside in the	
Township of Llai.	Varing.
byemale. On the West side the	0.
brook that runs from Henllan Mill into	
the river Elwy, about 300 yards from	
Trap Bridge, less than a mile from Garn.	
Mr. G	riff. b.
OSMUNDA lunaria. Old pastures West of Chirk	20
	Id.
regalis. In a boggy bottom between	
Waendwyssog and Rhydgaled, in the	
Parish of Henllan.	Id.

DENBIGHSHIRE.

LYCOPODIUM alpinum. Moors, very common.	
	Friffith.
POLYPODIUM Oreopteris. Very common in moun-	JU .
tainous pastures.	Id.
PTERIS crispa. About Cerig y Druidion.	Id.
ASPLENIUM septentrionale. On old Banks of Llan	
Dethylae, about a mile from Llanrwst.	
Dr. Rich	ardson.
SCOLOPENDRIUM vulgare. var. 2. With. Near	
a petrifying spring by the side of a rivulet	
at the bottom of Garn Dingle, three miles	
from Denbigh. Mr. (Friffith.
CYATHEA regia. Foot of the walls of Castell Di-	30
nas Brân. Rev. H. L	avies.
PHASCUM rectum. Among trees in the front of	
Garn House, very near the road leading	
to Henllan. Mr. G	riffith.
SPLACHNUM spharicum. (gracile?) Cowdung	
about Llyn Aled.	Id.
GRIMMIA recurvata. Garn Dingle.	Id.
calcarea. Garreg wen Rocks by Garn.	Id.
bomomalla. In Garn Dingle; and	
about peat holes near Llyn Aled.	Id.
heteromalla. Sides of Turbaries near	
Llyn-Aled.	I.J.
pusilla. Stones and roots of trees in	
damp situations in Garn Dingle, particu-	
larly on sand stones by the petrifying	
spring.	LJ.

 small rivers about Garn: in the river Clwyd, not ro yards above high water mark. Mr. Griffin. TORTULA tortuosa. On a bushy hill, half a mile from Garn, called Coed Mowr. Id. — cuneifolia. Between Garn and Henllan. Id. — cuneifolia. Between Garn and Henllan. Id. — cuneifolia. Between Garn and Henllan. Id. DICRANUM aciculare. Rocks on the side of the hill about 50 yards above Garthmeilio, the seat of R. W. Wynne, Esq. Id. — β. In the brook after crossing Pont y Alwen between Denbigh and Cerig y Druidion. Id. — strictum. Near Llyn Aled Lake. Id. * viridissimum. On trees near Eglwyfach. Id. POLYTRICHUM urnigerum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania. Id. NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. var. 2. With. On Turbaries about Llyn Aled. Id. — trichodes. Moors about Llyn Aled, abundantly. Id. 	TRICHOSTOMUM fontinalioides. Hedw. In many	,
Clwyd, not io yards above high water mark. Mr. Griffit. TORTULA tortuosa. On a bushy hill, half a mile from Garn, called Coed Mowr. Id. — cuneifolia. Between Garn and Henllan. Id. DICRANUM aciculare. Rocks on the side of the hill about 50 yards above Garthmeilio, the seat of R. W. Wynne, Esq. Id. — β . In the brook after crossing Pont y Alwen between Denbigh and Cerig y Druidion. Id. — strictum. Near Llyn Aled Lake. Id. * viridissimum. On trees near Eglwyfach. Id. POLYTRICHUM urnigerum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania, Id. NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. var. 2. With. On Turbaries about Llyn Aled. Id. — trichodes. Moors about Llyn Aled, abundantly.		
mark. Mr. Griffith TORTULA tortuosa. On a bushy hill, half a mile from Garn, called Coed Mowr. Id. from Garn, called Coed Mowr. Id. — cuneifolia. Between Garn and Henllan. Id. DICRANUM aciculare. Rocks on the side of the Id. hill about 50 yards above Garthmeilio, the Id. the seat of R. W. Wynne, Esq. Id. Id. mark. Ø. In the brook after Id. crossing Pont y Alwen between Denbigh and Cerig y Druidion. Id. mark. Id. Id. Id. mark. Near Llyn Aled Lake. Id. mark. Viridissimum. On trees near Eglwyfach. the viridissimum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania. Id. NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. war. 2. With. On Turbaries about Llyn Aled. Id. marcureum. dar. 2. With. On Turbaries about Id. Id.		
 TORTULA iortuosa. On a bushy hill, half a mile from Garn, called Coed Mowr. Id. <i>cuneifolia</i>. Between Garn and Henllan. Id. DICRANUM aciculare. Rocks on the side of the hill about 50 yards above Garthmeilio, the seat of R. W. Wynne, Esq. Id. <i>β</i>. In the brook after crossing Pont y Alwen between Denbigh and Cerig y Druidion. Id. <i>strictum</i>. Near Llyn Aled Lake. Id. <i>strictum</i>. Near Llyn Aled Lake. Id. <i>viridissimum</i>. On trees near Eglwyfach. Id. POLYTRICHUM urnigerum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania. Id. NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. var. 2. With. On Turbaries about Llyn Aled. Id. <i>trichodes</i>. Moors about Llyn Aled, abundantly. 		
from Garn, called Coed Mowr.Id.cuneifolia. Between Garn and Henllan.Id.DICRANUM aciculare.Rocks on the side of the hill about 50 yards above Garthmeilio, the seat of R. W. Wynne, Esq.Id	TORTULA tortuosa. On a bushy hill, half a mile	
 cuneifolia. Between Garn and Henllan. Id. DICRANUM aciculare. Rocks on the side of the hill about 50 yards above Garthmeilio, the seat of R. W. Wynne, Esq. Id. 6. In the brook after crossing Pont y Alwen between Denbigh and Cerig y Druidion. Id. strictum. Near Llyn Aled Lake. Id. strictum. Near Llyn Aled Lake. Id. * viridissimum. On trees near Eglwyfach. Id. POLYTRICHUM urnigerum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania: Id. NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. var. 2. With. On Turbaries about Llyn Aled. Id. 		
 DICRANUM aciculare. Rocks on the side of the hill about 50 yards above Garthmeilio, the seat of R. W. Wynne, Esq. Id. ————————————————————————————————————	cuneifolia. Between Garn and Henllan.	
 hill about 50 yards above Garthmeilio, the seat of R. W. Wynne, Esq. Id. 	DICRANUM aciculare. Rocks on the side of the	
the seat of R. W. Wynne, Esq. Id. ————————————————————————————————————		
crossing Pont y Alwen between Denbigh and Cerig y Druidion. Id. 		
and Cerig y Druidion. Id. ————————————————————————————————————		
 strictum. Near Llyn Aled Lake. Id. * viridissimum. On trees near Eglwyfach. POLYTRICHUM urnigerum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania; NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. war. 2. With. On Turbaries about Llyn Aled. trichodes. Moors about Llyn Aled, abundantly. 	crossing Pont y Alwen between Denbigh	
 * viridissimum. On trees near Eglwyfach. POLYTRICHUM urnigerum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania: NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. war. 2. With. On Turbaries about Llyn Aled. Id. 		Id.
Eglwyfach. Id. POLYTRICHUM urnigerum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania: Id. NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. war. 2. With. On Turbaries about Llyn Aled. Id. trichodes. Moors about Llyn Aled, abundantly.	strictum. Near Llyn Aled Lake.	Id.
 POLYTRICHUM urnigerum. Near the roadside between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania; NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. var. 2. IVith. On Turbaries about Llyn Aled. Id. trichodes. Moors about Llyn Aled, abundantly. 		
between Denbigh and Voylas, by the rivulet before you arrive at a place called Pensylvania: Id. NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. war. 2. With. On Turbaries about Llyn Aled. Id. trichodes. Moors about Llyn Aled, abundantly.		Id.
rivulet before you arrive at a place called Pensylvania: NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. war. 2. With. On Turbaries about Llyn Aled. Id. trichodes. Moors about Llyn Aled, abundantly.	POLYTRICHUM urnigerum. Near the roadside	
Pensylvania: Id. NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. war. 2. With. On Turbaries about Llyn Aled. Id. trichodes. Moors about Llyn Aled, abundantly.	between Denbigh and Voylas, by the	
NECKERA crispa. Garreg wen Rocks, abundantly. Id. BRYUM aureum. war. 2. With. On Turbaries about Llyn Aled. <i>trichodes.</i> Moors about Llyn Aled, abundantly.	rivulet before you arrive at a place called	
Llyn Aled. <i>trichodes.</i> Moors about Llyn Aled, abundantly.		Id.
Llyn Aled. Id. trichodes. Moors about Llyn Aled, abundantly.	BEXUERA crispa. Garreg wen Rocks, abundantly.	Id.
abundantly.	Thun Al. 1	
abundantly.		Id.
Id.	Light Aled,	
	abundantiy.	Id.

* Mr. Griffith communicated this plant to Mr. Dickson and Mr. Davies in 1800, so that he ought to be considered the first difcoverer.

M

VOL. I.

BRYUM alpinum. (as figured by Dill. 50. 64.c.)	
On a small dripping rock by Hendre	
Dwyfan bridge, near Cerig y Druidion.	
Mr. Gr	iffith.
HYPNUM undulatum. Garn Dingle, sparingly:	D
1	Id.
loreum. Woods about Garn, very com-	
	Id.
rugosum. Boggy parts of the hills about	
T 1 A 1 1	Id.
JUNGERMANNIA curvifolia. Near Llyn Lum-	
	Id.
RICCIA fruticulosa. Mossy stems of trees be-	
	D. T.
LICHEN confluens. Garreg wen, on limestone	
rocks. Mr. Gr	iffith.
pilularis. Stones upon Bettws mountain.	
	Id,
concentricus. Wall between Bettws	
mountain and Garthewin.	Id.
corneus. Oaks in Garn Dingle.	Id.
Acharii. On the shore of Llyn Aled	
lake, on stones of granite, which are	
	Id.
ferrugineus. Rocks near Llanufydd.	Id.
tricolor. Garreg wen rocks near Garn	Id.
gelidus. At Gallt y ddôl Garn, between	
	Id.

LICHEN aquilus. Rocks in Pengwern Frith	,
above the road leading from Llansanna	n
to Llanufydd. Mr.	Griffith.
	Id.
squamatus. On a wall about a mile from	n
Cerig y Druidión, on the roadside leading	5
to Denbigh.	Id.
carnosus. Rocks on the side of the hil	1
about 50 yards above Garthmeilio.	<i>I.</i> ?.
marginalis. Fir plantation near Cefr.	1
House.	Id.
muscicola. About Garthmeilio, abun-	-
- dantly.	Id.
pulmonarius. Garn Dingle.	Id.
glomuliferus. About Cenioge House,	,
between Llanrwst and Corwen: also be-	
tween Llanrwst and Capel Cerig.	Id,
scrobiculatus. Garn Dingle; and woods	
about Garthewin.	Id.
plumbeus. Garn Dingle; and about	
Llanrwst.	Id.
miniatus. In a tower of Denbigh Castle	
above the Goblin well: also on Garreg	
wen rocks near Garn.	Id.
saccatus. Garreg wen rocks near Garn.	Id.
resupinatus. Garn Dingle.	Id.
granulatus. Garreg wen rocks.	Id.
N 2	

ī

DENBIGHSHIRE.

LICHEN cochleatus.] About Garn, very common. 	
sinuatus. S Mr. G	riffith.
fascicularis. Ash trees in bleak situ-	
ations about Nant Glyn.	Id.
fluviatilis. On stones in the river Elwy,	
about half way from the ford, opposite	
the cave to Pont-newydd, four miles from	
Denbigh.	Id.
Fucus ligulatus. } Near Abergele.	TJ
	Id.
CONFERVA fluviatilis. Rivulet in Garn Dingle.	Id.

FROM Mr. Pilkington's "View of the present state of Derbyshire" we have extracted, not merely the habitats followed by his own name, but those also which are inserted upon the authority of Mr. Whately, Mr. Coke, Mr. Hallows, Mr. Hadley, and Dr. Johnson, under which is comprised by far the greater part of the information we have been able to procure respecting this most beautiful and interesting County, the opposite extremes of which, so entirely unlike each other, and comprehending every gradation from the most fertile cultivated fields to the wildest mountains and bleakest moors, are highly favorable to the views of the Naturalist.

CIRCEA alpina. Matlock. Mr. J. Woods, jun. side near Matlock Bath. Smith. SALVIA pratensis. Crich. Mr. Hallows. CROCUS sativus. Holmes: and Siddals near Derby: Breadfall; and Duffield, near the Banks of the River Derwent. Pilkington AVENA pubescens. Common in the County. 1a. ELYMUS europæus. Rocks opposite Matlock Baths. Mr. Woodward. DIPSACUS pilosus. At Matlock Bath. Smith. Derby: betwixt Derby and Spondon. Dr. Johnson. GALIUM pusillum. At Matlock, and elsewhere in the County. plentiful. Mr. Woodward. Middleton Dale. Mr. Coke. Dovedale. Mr. S. Woods. SANGUISORBA officinalis. Common in moist pastures. Pilkington. About Matlock and Buxton. Mr. Woodward. POLEMONIUM caruleum. Matlock: Alfreton Brook. Mr. Coke. Lover's Leap, near Buxton. Mr. Wood. By the side of the Turnpike road in Bakewell Meadows. Mr. Whately. Banks of the Wye between Buxton and Bakewell. Mr. O. Sims, CAMPANULA patula. Frequent. Pilkington. (Surely this must be a mistake, for so rare

a plant could hardly have escaped the notice of other Botanists.) CAMPANULA latifolia. Between Ashbourn and Dovedale. Dr. Johnson. Little Chester, near Derby : between Rowsley and Darley. Pilkington. Near Matlock on the Ashbourn Road. D. T. DATURA Stramonium. Derby; and Pinxton. Pilkington. VIOLA lutea. Hills above Dovedale. Mr. Woodward. Lea: Wirksworth: Chapel le Frith: Tideswell Moor: Chatsworth. Pilkington. Abundant on the moors round Buxton. D. T.RHAMNUS catharticus. Not uncommon in hedges and woods. Pilkington. VINCA major. Pleasly Park. Mr. Coke. HERNIARIA birsuta. Duffield. Mr. Hadley. CHENOPODIUM olidum. Dethick. Mr. Hallows. GENTIANA pneumonanthe. Egginton Heath. Pilkington. ----- campestris. Mountainous Pastures near Buxton, in the same luxuriance as delineated in Flora Danica, t. 367. Sir T. G. Cullum. Chinley Hill, near Chapel le Frith. Mr. O. Sims. CAUCALIS latifolia. Dovedale. -Dr. Johnson. HERACLEUM sphondylium. B. Between Ashbourn and Okeover. Dr. Stokes. Hedges near Bakewell. D. T.

SCANDIX odorata. Banks of the Derwent above Chatsworth. Mr. Whately. About Buxton. Rev. W. Wood. In the little inclosed Meadows about Poole's Hole at Buxton. Sir T. G. Cullum. SAMBUCUS Ebulus. S. Normanton: Bakewell: Dethick: Wirksworth: Alport near Youlegrave; and Boulton. Pilkington. Behind the White Hart, Buxton. Rev. W. Wood.

LINUM usitatissimum. Pinxton. Mr. Coke. MYOSURUS minimus. Cornfields near Derby. Mr. Whately. ORNITHOGALUM luteum. Park Hall Woods. Rev. Mr. Pashley. In a meadow adjoin-

ing the Copper Mills, Derby. Mr. Whately. CONVALLARIA multiflora. Pleasly Park. Mr. Hallows. ACORUS Calamus. Cutthorpe near Chesterfield. Id. RUMEX sanguineus. Dethick. Id.

COLCHICUM autumnale. Meadows at Marston Montgomery: pastures between Duffield and Burley: very abundantly in Breadsal long-meadow, and near Derby. Pilkington.

ALISMA ranunculoides. Between Derby and Burton. Mr. Whately.

EPILOBIUM angustifolium. Matlock: Darley. Mr. Coke. On high rocks by Sparry Pit near Chapel le Frith. Mr. O. Sims.

CHLORA perfoliata. Pleasly Park. Mr. Coke. Whitewell. Pilkington. VACCINIUM Vitis idea. Plentiful among the heath in the mountainous parts. Ray. East Moor. Rev. W. Wood. On High Moor near Hayfield. Mr. O. Sims. Oxycoccos. On the Moors. Pilkington. On High Moor near Hayfield : on Kinder Scout; and on Mosses near it. Mr. O. Sims. ERICA multiflora. (vagans?) Ashover. Pilkington. (If this habitat be correct, it is a very interesting addition to the British Flora: at all events it would be worth the while of any Botanist to ascertain the plant intended.) DAPHNE Mezereum. Matlock, Chee Tor. Mr. Coke. POLYGONUM Bistoria. S. Normanton : Crich : in a meadow by Mr. Bennett's, Mackworth. Pilkington. At Ashbourn. Mr. J. Woods, jun. PARIS quadrifolia. Pinxton: Newton wood. Mr. Coke. Love-lane near Derby. Mr. Whately. In Longford long-lane, in a pit near Dalbury Lees. Pilkington. Rocks opposite Matlock Bath. D. T. ARBUTUS Uva Ursi. On Kinder Scout near Hayfield. Mr. L. Howard.

CHRYSOSPLENIUM alternifolium. Dethick. Mr.
Hallogue Chas T
SAXIFRAGA aizoides. I received a specimen of
this and S. caspitosa from Derbyshire.
Mr. L. Howard.
cæspitosa. Castleton Hill. Mr. Coke.
Middleum D. 1. Mr. Coke.
Middleton Dale. Mr. Whately. Dove-
dale. Mr. Woodward.
SAPONARIA officinalis. Elmeton. Mr. Hallows.
Duffield. Pillington
DIANTHUS caryophyllus. B. Near Edensor Mr. Col.
- deltoides. Hills between Bakewell
and Chatsworth. Mr. Whately. On rocks
in Litton Dale near Tideswell. Mr. O. Sims.
SILENE nutans. Middleton Dale. Mr. Coke.
Rocks in Dovedale. Mr. Woodward,
ARENARIA verna. Matlock. Mr. Woodward,
About the mouther of The woodward.
About the mouths of Lead-mines. Dr.
Smith. Between Bakewell and Castle- ton.
What when the woods in the
SERVER dassethell D: SERVER dassethell D:
Pinxton.
rupestre. Rocks near Castleton. Mr. O. Sims.
KASIIUM tomentosum. Kedleston. Mr. Hadlen
TRONOS Padus. Chee Tor. Matlock Mr. Coke
Matlock. Mr. Coke. Mountain-
ous parts of the County, from the

C

fissures of the precipices without any ap-Mr. Woodward. pearance of soil. PYRUS domestica. Crich woods : Matlock. Mr. Coke. ROSA villosa. Pastures at Rough Heanor. Pilkington. In several Hedges about Hather-Martin sage. RUBUS sanatilis. Hills opposite Matlock Bath. Mr. Woodward. ----- chamæmorus. On Ax-edge near Buxton. Mr. Lambert. N. E. side of Kinder Scout near the summit. Mr. O. Sims. GEUM rivale. Chee Tor: Hassop: Pinxton. Mr. Coke. About Hayfield. Mr. L. Howard. Dovedale. Mr. J. Woods, jun. Beard's wood near Low Leighton. Mr. O. Sims. Dr. Smith. THALICTRUM minus. In the entrance of the cave at Castleton: Between Matlock and Castleton. Mr. J. Woods, jun. Litton Dale; and Monson Dale. Mr. O. Sims. ADONIS autumnalis. In fields, but not very com-Dr. Johnson. mon. RANUNCULUS Lingua. S. Normanton. Mr. Coke. TROLLIUS europæus. Litton Dale near Tideswell. Mr. O. Sims. HELLEBORUS viridis. Matlock : Codnor Castle. Mr. Coke

HELLEBORUS fætidus. Cromford Moor. Mr.
Coke. Dethick. Mr. Hallows.
Ajuca alpina. On the mountain that leads from
Matlock into the Town of Castleton, on
the left hand side, immediately adjoining
the Road. D. T.
MENTHA sylvestris. Near Matlock. Mr. J.
Woods, jun. Pinxton, Mr. Colo
birsuta. ζ. Common. Pilkington.
piperita. South Normanton. Pilking-
ton. Rivulet in Bonsall Dale near Mat-
lock. Smith. Pingett wood, Low Leigh-
ton. Mr. O. Sims.
Pulegium. Pinxton. Mr. Coke. Ock-
brooke, Radbourn, and Langley Com-
mons. Pilkington.
GALEOPSIS versicolor. About Matlock Bath, and
Driffield. Sir T. G. Cullum.
STACHYs germanica. Pinxton. Mr. Coke.
LEONURUS Cardiaca. Handley. Mr. Coke.
Mackworth. Pilkington.
THYMUS Nepeta. S. Normanton. Id.
Scutellaria minor. Pinxton. Id.
MELAMPYRUM sylvaticum. Crich: Matlock.
Mr. Coke. (We have inserted this, be-
cause it stands so in Pilkington, but at the
same time as he has not noticed M. pra-
tense, we feel little doubt of that being
the plant intended.

LATHREA squamaria. Pleasly Park. Mr. Hallows. ANTIRRHINUM Cymbalaria. Wingerworth : Glapwell: S. Normanton. Mr. Coke. On an old wall at Pleasly, between Chesterfield and Mansfield. Rev. W. Wood. minus. Pinxton. Mr. Coke. DRABA muralis. Frequent in this County. Hudson. Elden Hole, by Castleton. Rev. W. Wood. ALYSSUM sativum. Normanton. Mr. Coke. COCHLEARIA officinalis. y. Mountains round Castleton. D.T. ----- anglica. Hills near Castleton. Mr. O. Sims. On Kinder Scout. Mr. L. Howard. THLASPI alpestre. Matlock, plentifully. Mr. Woodward. CARDAMINE impatiens. Woods on the rocks opposite the Bath at Matlock, plentifully. Mr. Woodward. Poole's Hole near Buxton. Mr. J. Martin, amara. Siddals near Derby: Rough Heanor. Pilkington. SISYMBRIUM Irio. Wingfield Manor. Id. ERYSIMUM cheiranthoides. Near Ashbourne. Martyn. HESPERIS inodora. Banks of the Derwent, near Matlock. D. T. ARABIS hispida. Middleton Dale. Mr. Coke. TURRITIS glabra. Between Ashbourne and Okeover. Dr. Stokes. ERODIUM moschatum. Alfreton. Mr. Coke.

GERANIUM pratense. Frequent in Meadows. Pilkington. Most abundantly about the Derbyshire side of Burton Bridge. Waring. Dovedale. Mr. S. Woods. pyrenaicum. In Litton, and Monsal Dales near Tideswell. Mr. O. Sims. rotundifolium. Derby. Pilkingtons sanguineum. Near Buxton. Mr. Woodward. Litton, and Monsal Dales near Tideswell. Mr. O. Sims. * FUMARIA lutea. Old walls near Castleton. Mr. L. Howard, LATHYRUS Aphaca. Pinxton: Derby. Pilkington. Nissolia. Between Breason and Risley. Pilkington. Carnfield, near Alfreton. Dr. Johnson. birsutus. S. Normanton. Mr. Coke. palustris. Pinxton. Id. VICIA sybuatica. Hedges behind Matlock Bath. Dr. Smith. ----- lutea. Pinxton; and Derby. Pilkington. LACTUCA scariola. Near the entrance into Peake's Hole. Mr. J. Martin. + HIERACIUM dubium. Coxbench Wood. Dr. Johnson.

* See Note p. 117.

† An Hieracium, distinct I believe from every other British species, but most nearly allied to *H. murorum*, grows on the walls of Custleton Castle. It was first mentioned to me by Rev. Mr.

HIERACIUM paludosum. Meadows near the
Town of Castleton. Sir T. G. Cullum.
CARDUUS eriophorus. In dry pastures near Low
Leighton. Mr. O. Sims. About the
Zig-Zag walk at Mark 1
zig-zag walk at Matlock. Dr. Smith. Thorp. Mr. S. W.
GNAPHALIUM gallicum. Heaths. Mr. Woodward.
dioicum. Hills between Havfield
and Kinder Scout. Mr O Since
DENECTO saracenicus. Between Chesterfield and
Sheffield. Par III THE
ORCHIS usiulata. Alfreton: S. Normanton D.U.
SATYRIUM bircinum. Crich. Mr. Coke.
wir. Coke.
neighbourhood of Chapel le Frith.
OPHRYS cordata On the line of Mr. L. Howard.
OPHRYS cordata. On the high Moor between
Chatsworth and Sheffield. Dr. Smith.
On Kinder Scout. Mr. O. Sims.
- apyera. In a field opposite the old Bath
at Watlock.
muscifera. 1 never found this plant in
wereysine, but, from the great numbers
which I have seen exposed for sale as
Matlock, it cannot be uncommon in the
neighbourhood.
Jewton of Jesus College Cambridge D. T.

llege, Cambridge, and I afterwards gathered it myself; having given away my only specimen, I cannot now state any particulars respecting it. D. T.

ñ

SERAPIAS grandiflora. Newton Wood. Mr. Coke. CAREX curta. Chee Tor. Id. EMPETRUM nigrum. On the Moors about Buxton. Ray. Abundant near Chapel le Frith. D.T. TAXUS baccata. Rocks at Matlock and Dovedale; and in a scattered manner in other parts of the County. Pilkington. EQUISETUM fluviatile. Newton wood. Id. OSMUNDA lunaria. Dethick. Mr. Coke. Chinley Hill, near Chapel le Frith. Mr. O. Sims. Heights of Abraham behind Matlock Bath. D. T. --- B. Buxton, on the road to Manchester. Rev. W. Wood. LYCOPODIUM selaginoides. 7 On Kinder Scout. inundatum. Mr. O. Sims. alpinum. Mountains near the Derwent. Mr. J. Martin. ASPLENIUM viride. Coxbench Wood. Dr. Johnson. POLYPODIUM Dryopteris. About the rocks near Pleasly Forges. Deering. Chinley Hill near Chapel le Frith. Mr. L. Howard. ---- calcareum. About Matlock Bath. Dr. Smith. PTERIS crispa. Chinley Hills near Chapel le Frith. Mr. L. Howard. TRICHOSTOMUM fontinalioides. River at Matlock Bath. Mr. Teesdale.

TRICHOSTOMUM fontinalioides. B. River at Bux-
ton. Mr. Bradbury.
DICRANUM flagellare. Cromford Moor, near Mat-
lock. Smith
NECKERA crispa. Rocks at Buxton. Mr. Bradbury.
HYPNUM Teesdalii. Woods on the S. E. Side
of the river at Matlock. Mr. Teesdale
recognitum. Matlock. Dr. Smith.
LICHEN physodes. In fruit on Cromford Moor. Id.
fragilis. Near the high rocks on Cromford
Moor. Id.
globiferus. Abundant on Cromford Moor. Id.
senguinarius. Granite rocks on Cromford
Moor. Id.
earth covered with decayed Moss. Dickson.
corallinus. Coxbench, frequent. Pilkington.
cæruleo-nigricans. Dovedale. Dr. Johnson.
miniatus. Dovedale. Dr. Johnson. Plen-
tiful under hanging rocks above Sexton's
Bath, Matlock. Dr. Smith.
omphalodes. Abundant on the high stony
rocks of Chamber 1 DA
Id. Id. Id.
resupinatus. Plentiful near Broad-bottom
Bridge, Mr. Bradbury.
all o all (GUUTY.

٤)

DEVONSHIRE.

THE County of Devonshire, the second in the Kingdom in point of extent, has long been known to comprize several plants, which are scarcely to be found in any other part of England, especially Scirpus holoschænus, Polycarpon tetraphyllum, and Oxalis corniculata. To these is now to be joined Cynoglossum omphalodes, which we have added to the British List entirely upon the authority of Mr. Polwhele, who does not appear to entertain any doubt of its right to a place in our Flora, from its having been found among the rocks at Teignmouth, a spot so wild that it is by no means probable it should have been planted there intentionally, or that the seeds should by any accident have escaped out of a garden. From this Gentleman's history of the county we have derived that great proportion of our present Catalogue, which stands either

under his name, or those of Dr. Wavell, Mr. Weston, Mr. Yonge, and Mr. Cornish. Bishop Gibson also, in his edition of Camden, has inserted in this, as in many other instances, a list of the indigenous plants, and Mr. Pike of Biddeford, has published a catalogue of those found in his own neighbourhood; but many of these, as Mr. Polwhele observes, are not peculiar to that Town, or its environs, and, besides this, he leaves room for suspecting his accuracy, by having inserted the Coix Lachryma Jobi, which there is no reason to believe really a native of this Kingdom. In spite of all this, however, Devonshire is by no means one of those Counties whose Botany appears to have been most successfully investigated, and we have no doubt of seeing at some future time a much. more copious catalogue of its rare vegetable productions.

VERONICA montana. In a lane near Torrington, leading towards Holdsworthy. Dr. Wavell.
PINGUICULA lusitanica. In boggy meadows midway between Oakhampton and Launceston, betwixt a great wood and the river. Ray. Haldon goyle. Polwbele.

IRIS fatidissima. Very common in the S. W. of this county, particularly about Teignmouth. Withering. On the banks of North Yeo. Dr. Wavell. About Biddeford.

SCHOENUS albus. On a Common near Axminster. Mr. E. Forster, jun.

SCIRPUS Holoschanus. On Braunton Burroughs. Ray. It grows there plentifully, and covers nearly half an acre of ground.

Dr. Wavell.

ERIOPHORUM vaginatum. Bogs near the road leading from Clovelly to Kilkhampton.

Dr. Wavell. On Dartmoor. Mr. E. Forster, jun.

MILIUM lendigerum. Near Instow. Curtis. AGROSTIS setacea. Piddletown Heath. Curtis. Common in this and other Western Counties. Earl of Gainsborough.

MELICA nutans. In a wood near Dolton. Dr. Wavell. POA bulbosa. Plentiful on the Den at Teignmouth.

Rev. Dr. Beeke. ELYMUS arenarius. Near Exmouth. Polwhele. POLYCARPON tetraphyllum. At Lymston, near Exeter. Mr. Newbery. Exmouth. Rev. J. Lightfoot. On the walls of Mr. Smith's house, at Lymston: Mr. Weston. DIRSACUS Fullonum. Very plentiful in the South-

hams, Mr. Cornish. Common in the

North of Devon. Dr. Wavell. Meadows near Alphington; and in George'sclyst. Polwhele.

* RUBIA peregrina. On Chudleigh rock, ten miles beyond Exeter. Sir T. Cullum. Common in the hedges near Dawlish. Maton. Very common in the hedges of the road from Barnstaple to Biddeford; and also near Braunton; and various other places. Dr. Wavell. About 'Feignmouth and Exeter. L. W. D. Near the Devil's Point at Stonehouse, near Plymouth. Mr. E. Forster, jun. Near Exmouth. Hudson. Cliffs at Sidmouth. D. T. EXACUM filiforme. Here and there in the County. Hudson. More frequent in Devon than in other counties. Mr. Yonge. SANGUISORBA officinalis. In meadows, not common. Dr. Wavell. RUPPIA maritima. Ditches in Braunton marsh. Id.

ANCHUSA sempervirens. Near Sidmouth. Wither-

* I cannot help thinking that the *Rubia* found at Dover, and that which abounds in the Western Counties, are distinct species. In the former the leaves are elliptical and sub-quaternate, in the latter, lanceolate and quinate, nor have I ever seen them grow by fours, except occasionally in two or three of the end whorls: the former is procumbent on the beach and chalky cliffs, the latter climbs over hedges and bushes. *L.W.D.*

ing. Kenton; and Maidenhead. Rev. Dr. Beeke. About Barnstaple. Dr. Maton. On Durnsford Hill, between Exeter and Moreton Hampstead.

Mr. E. Forster, jun.

CYNOGLOSSUM omphalodes. Found by Mrs. Taylor at Teignmouth, among the rocks.

Polyubele.

Ray.

PULMONARIA maritima. The Southams. Mr. Cornisb. SYMPHYTUM officinale. B. Meadows between

Lyme Regis and Sidmouth. D. T. CAMPANULA hederacea. Frequent in moist and shadowy places throughout the county. L. W. D. Lidford water-fall. Dr. Maton. Valley between Biddeford and Kilkhampton; and near Woodford Bridge.

LOBELIA urens. On the sandy sides of a small rill which flows by the Turnpike road on Shute Common, descending towards Axminster. (Kilmington Hill and Shute Common (vide Fl. Brit.) are the same; and on this Common the plant is confined to a spot about two hundred yards in length, and not spreading at all into similar situations). Rev. Dr. Beeke. Near Ottery St. Mary. Miss Burges. VERBASCUM Lychnitis. Hedges, not common.

Dr. Wavell,

VERBASCUM Blattaria. Frequent about Plymouth and Ashburton. Hudson. At East Teignmouth, near the New Chapel. Mr. Weston. Not uncommon in Devon. Rev. Dr. Beeke. CHIRONIA pulchella. Braunton Burroughs. Rev. Dr. Goodenough. RHAMNUS catharticus. Plentiful at Brixton; and not uncommon in other parts of Devon. Polwhele. ILLECEBRUM verticillatum. Not uncommon in this County. Withering. SALSOLA fruticosa. Sea Coast, Devon. Hudson. ERYNGIUM campestre. On a rock as you go down to the Ferry from Plymouth into Cornwall. Ray. About the Eastern point of the entrance into Hamoaze. Mr. Yonge. HERACLEUM sphondylium. B. Hedges near Holdsworthy. D.T. SIUM vericillatum. On the road to Moreton, near the two mile stone, on a Common at the left hand. Mr. Weston. (This hardly appears a probable station for so extremely rare a plant.) OENANTHE pimpinelloides. On the approach to Cleave House, in the ditch on the right hand about one-third of the way up the hill. . Mr. Weston. SAMBUCUS Ebulus. Staverton. Polaubela.

CORRIGIOLA littoralis. Found by Mr. Hudson on Slapham Sands beyond Dartmouth; and on the Start Point. Prof. Mariyr. LINUM usitatissimum. Cornfields and sandy pastures: on Dartmoor. Polzubele. ------ perenne. Hedges near Sowdown. Id. ------ angustifolium. Roadsides luxuriantly decorated with this plant near Dartington Bridge. Maton. By the shore near Plymouth. Mr. E. Forster, jun. DROSERA anglica. Plentiful on the swamps, or rather Goyles, of Haldon and Blackdown. Polubele. GALANTHUS nivalis. Near St. John's Chapel, and at Broadgate, Barnstaple. Id. SCILLA verna. Hartland Promontory; and Torquay. Maton. --- autumnalis. On Sir Robert Palk's high grounds at Torkay, on the left hand, plentifully. Mr. Weston. In great plenty on a hill that overlooks Bigbury Bay, near a point called by the Fishermen, Grenore Point. Mr. Yonge. Acorus Calamus. About Biddeford. Pike. JUNCUS acutus. Instow; and Braunton Burroughs, plentifully. Rev. Dr. Goodenough. CHLORA perfoliata. Braunton Burroughs. Polwhele. ERICA vagans. Heath between Axminster and Lyme. Miss Webster.

PYROLA rotundifolia. Biddeford; and the neigh-	
bourhood. Pike. I have never met with	
it at Biddeford, or in any part of Devon-	
shire. Dr. Wavel	7.
CHRYSOSPLENIUM alternifolium. Banks of the	
Yeo, near the Oil-mill. Polwhele	•
SAPONARIA officinalis. In great plenty near the	
sands at Shaldon. Mr. Cornish. Near	
Wilmington, between Axminster and	
Honiton. Mr. E. Forster, jun	•
DIANTHUS d. lioides. On the road from Kings-	
bridge to Teignmouth. Polwhele	•
SILENE anglica. Teingrove, on the rubbish dug	
from a canal. Rev. Dr. Beeke	
acaulis. On Dartmoor. Hudson	
OXALIS corniculata. In several places in the neigh-	
bourhood of Exeter. Mr. J. Turner.	
In a waste rubbishy spot at Dawlish; and	
since in several other places in the County.	
Mr. Martyn. In a lane near Topsham.	
Miss Burges. Plentiful near the rivulet	
that runs through Dawlish. Rev. Dr. Becke.	
SPERGULA subulata. Dry pastures on a gravelly	
soil. Polwhele.	
EUPHORBIA Peplis. In the loose sand near Ex-	
mouth, plentifully. Hudson. Northam	
Burroughs. Mr. Yonge.	
portlandica. On the Warren, near	
Exmouth, plentifully. Rev. Dr. Good-	

enough. At Braunton Sands on the Northern Coast. Rev. J. Lightfoot. MESPILUS germanica. In a hedge on the right hand side of the Biddeford Road, near the marsh, about six miles and a half from Barnstaple. Polyuhele. PYRUS Aria. Woods and Hedges at Goodleigh, &c. Id. the cliffs near Torkay : in great plenty at Hatzel, in the Parish of Ashburton. Id. * CISTUS poinfolius. At Babbicombe, near Newton Abbot. Rev. A. Neck. Plentiful on the rocks above Tor-quay. Rev. D. Beeke. TEUCRIUM scordium. Lord Courtenay's Park : Braunton Burroughs. Polywhele. MENTHA viridis. Near Exmouth. Hudson. ---- rotundifolia. Lord Lisburne's grounds. Mr. Weston. gentilis. Winscott. Polywhele. LEONURUS Cardiaca. Near the rope-walk, Biddeford. Id. + MELITTIS melissophyllum. Woods about

* It is said by a writer in the Gentleman's Magazine, who signs himself *Incompettus*, that the single yellow Rose grows spontaneously on Exmoor. See vol. 67. p. 197

[†] There can be little doubt but most if not all of these habitate belong to *M* grand flora; but as the two species were not separated when Mr. Polwhele published his work, we have thought it best to retain them as placed by him.

Totnes. Ray. In the Southams, particularly in Hempston wood. Mr. Cornisb. Dartington woods : Canonteign woods, in the road to Brampford-Speke, beyond Sir Stafford Northcoue's Pillars on the left hand, not far from the blacksmith's shop. Mr. W. ston. A small wood not far from Kingsweare, near Dartmouth. Miss Burges. Woods near Biddeford ; and near Hall. Dr. Wavell.

MELITTIS grandiflora. Common in most Coppies in every part of the County. Row. Dr. Beeke. By the road side at Hall, near Barnstäple. Curtis. A mile from Ashburton on the road to Plymouth in the hedges, in great plenty. D. T.

Scutellaria minor. Woolleigh, near Kilkhampton. Palcuhele.

BARTSIA viscos 1. Near Whistley Cross, Mamhead Mr. Weston. Fields at Instow. Dr. Wavell. Frequently about the springs which rise at the tops of the hills that overlook Bigbury Bay. Mr. Yonge. Wet places in the Parish of Kenton. (The Calyx never of any other colour than the leaves.) Rev. Dr. Reebe. ANTIRRHINUM Cymbalaria. Walls, not common.

Polyubele.

SIBTHORPIA europaa. Plentiful in this County.
Dr. Sibthorpe.
OROBANCHE ramosa. Dry pastures in Devon.
Hudson.
LEPIDIUM ruderale. Near Exeter. Magna Britannia.
COCHLEARIA officinalis. On the Coast; and at
Biddeford, near the Bridge. Polubele.
On a stone wall between Oakhampton
and Lifton. Mr. E. Forster, jun.
γ. Walls of Georgham
Church Yard. Polwhele.
danica. On Braunton Burroughs.
Polwhele. Near Dartmouth. Dr. Ma-
ton. About the village of Tor-quay,
but rather scarce. Rev. Dr. Beeke.
Plentiful about Ilfracombe. L. W. D.
CORONOPUS didyma. Topsham Road near Coa-
ver: near the Courtney Arms, under the
dwarf wall, Starcross. Mr. Weston. Very
plentiful as a weed in all the Gardens
about the Walls of Exeter. Rev. Dr. Beeke.
ISATIS tincloria. Cornfields, and borders of
cornfields, rare. Polwhele.
CRAMBE maritima. Cliffs near Teignmouth.
Dr. Maton. Frequent on the marly
Cliffs, but rare in pure sand. Rev. Dr.
Beeke. Sidmouth Cliffs, in inaccessible
places. D. T.

* ERYSIMUM præcox. On a hill about half a mile
north of Teignmouth: about Dawlish:
Kingsteignton: Honiton; and Bovey
Tracy. Rev. Dr. Beeke.
CHEIRANTHUS sinuatus. On the rocks adjoining
Braunton Burroughs. Polwhele,
ERODIUM moschatum. At Teignmouth. Wither-
ing. Bank under a Hedge near Upcott.
Polwhele. Near Plymouth. Mr. E. Forster, jur.
maritimum. On the Sea Coast at Teign-
mouth. Withering.
GERANIUM rotundifolium. Near Plymouth.
Mr. E. Forster, jun.
Mr. E. Forster, jun. LAVATERA arborea. At Teignmouth. Withering.
LAVATERA arborea. At Teignmouth. Withering. On the rock at the entrance of Torbay, plentifully. Mr. Weston.
LAVATERA arborea. At Teignmouth. Wathering. On the rock at the entrance of Torbay, plentifully. Mr. Weston, LATHYRUS Nissolia. Very common in the South-
LAVATERA arborea. At Teignmouth. Wathering. On the rock at the entrance of Torbay, plentifully. Mr. Weston. LATHYRUS Nissolia. Very common in the South- ern parts of Devon. Mr. Cornisb. Also
LAVATERA arborea. At Teignmouth. Wathering. On the rock at the entrance of Torbay, plentifully. Mr. Weston, LATHYRUS Nissolia. Very common in the South-
LAVATERA arborea. At Teignmouth. Withering. On the rock at the entrance of Torbay, plentifully. Mr. Weston. LATHYRUS Nissolia. Very common in the South- ern parts of Devon. Mr. Cornish. Also in every other part. Polwhele. About Teignmouth. Dr. Mariyz.
LAVATERA arborea. At Teignmouth. Withering. On the rock at the entrance of Torbay, plentifully. Mr. Weston. LATHYRUS Nissolia. Very common in the South- ern parts of Devon. Mr. Cornish. Also in every other part. Polwhele. About
LAVATERA arborea. At Teignmouth. Withering. On the rock at the entrance of Torbay, plentifully. Mr. Weston. LATHYRUS Nissolia. Very common in the South- ern parts of Devon. Mr. Cornish. Also in every other part. Polwhele. About Teignmouth. Dr. Martyn. Aphaca. Hedges near Chittlehamp- ton. Polwhele.
 LAVATERA arborea. At Teignmouth. Withering. On the rock at the entrance of Torbay, plentifully. Mr. Weston. LATHYRUS Nissolia. Very common in the Southern parts of Devon. Mr. Cornisb. Also in every other part. Polwhele. About Teignmouth. Dr. Mariyn. — Aphaca. Hedges near Chittlehamp-

* Dr. Beeke observes that he has only once found E. Barbarea in Devonshire, and then near a Garden; and Mr. Polwhele gives but one habitat for it.

dridge; and the neighbourhood. Polwhele. Cliffs at Ilfracombe. Mr. Sowerby. VICIA bithynica. On most of the red Cliffs, but near the Sea only. (It varies in the breadth of its leafits, but not in their number, nor in any other circumstance.) Rev. Dr. Beeke. ASTRAGALUS glycyphyllos. Lord Lisburne's Grounds. Polwhele. TRIFOLIUM suffocatum. Plentiful on the Den at Teignmouth. Rev. Dr. Beeke. Lorus diffusus. In three places remote from each other in the parishes of Bishopsteignton, and Kingsteignton : plentiful in each place, but only on the outside of dry banks. Id. MEDICAGO falcata. In all the hedges in the Southams. Polyubele. HYPERICUM Androsemum. Brixton. Id. * HIERACIUM umbellatum. (var. foliis penè ovatis, vix dentatis, caule humili, pene unifloro.) Near Upton Pyne. Dr. Maton. ANTHEMIS nobilis. On the sands at Slapton-lea, in great quantity. Polyubel.

* Some specimens exactly answering this description were found two years ago, at Rollesby, near Yarmouth, by Mr. Wigg, and have all the appearance of being a distinct species; but, unfortunately, from the habitat being destroyed a stop has been put to farther investigation. D. T.

OPHRYS apifera. Found by Mr. Beeke at Kingsteignton. Polaubele. CAREX extensa. On the marshy part of the west side of Braunton Burroughs. Rev. Dr. Goodenough. EQUISETUM fluviatile. Between Axminster and Honiton, common. Mr. E. Forster, jun. OSMUNDA regalis. Near Lifton. Id. POLYPODIUM Phegopteris. Here and there in this County. Hudson. ----- spinulosum. Hollow ways in Devon. Withering. ASPLENIUM marinum. Rocks by the sea side. Hudson. SCOLOPENDRIUM Ceterach. On a wall between Totnes and Newton Bushell. Mr. E. Forster, jun. HYMENOPHYLLUM tunbrigense. Moist rocks on Dartmoor. Hudson. PILULARIA globulifera. Wet places on Blackdown. Polaubile. GYMNOSTOMUM osmundaceum. In the road from Zele to South Tawton Church, four miles from Okehampton. Rev. Mr. Newberg. SPLACHNUM Turnerianum. In a moor to the right hand of the road about three miles from * Sidmouth towards Exeter. D. T.

* The words Sidmouth, and Excher are reversed in Flora Britannica.

JUNGERMANNIA purpurea. On moist peat earth	
on a mountain called Cowsand; and on	
the sides of hills in the forest of Dart-	
moor. Mr. Newber	30
julacea. On rocks on the sides	
and tops of hills in Dartmoor. Id.	
TARGIONIA hypophylla. On heaths, and by the	
sides of ditches near Dawlish, by Ex-	
mouth. Hudse	071.
LICHEN geographicus. Rocks in the Valley of	
Stones near Linton. D.	T.
ventosus. Upon rocks on the sides and	
tops of hills in Dartmoor. Mr. Newbe	ry.
furfuraceus. Plentiful on Wild Tor Rock,	
a large rock five miles from Chagford,	
and also on many of the smaller rocks of	
granite thereabouts. Id.	
glomuliferus.	
ash in the N. W. of the County. Id.	
plumbeus. County. Id.	
perlatus. Abundant in the County; but	
Mr. Sowerby and I found it in fruit only	
on a stone fence near Ilfracombe, on the	
road from Barnstaple. D.	T.
townefacture Rocks in the mountainous	
proboscideus. } Rocks in the mountainous part of Dartmoor, rare.	
Mr. Newler	- V .

LICHEN cochleatus. Roots of Trees. Mr. Slater.
On rocks and stones, abundantly. D. T.
sinuatus. Rocks about Ashburton. Hud-
son. On a stone wall, at Chagford, be-
tween the town and parsonage, but with-
out shields. Mr. Newbery.
aphthosus. Rocks on Dartmoor. Hudson,
sylvaticus. var. 2. With. On ash, syca-
more, and oak in the N. W. of Devon.
Mr. Newbery.
fallax. Rocks in this County. Mr. Slater.
horizontalis. On Durnsford Hill, be-
tween Exeter and Moreton Hampstead.
Mr. E. Forster. jun.
chrysophthalmus, Brought on branches
of apple trees from this County to Mr. Sowerby.
globiferus. Rocks in the mountainous
paschalis. > parts of Dartmoor.
paschalis. parts of Dartmoor. Mr. Newberg.
lanatus. Rocks on the sides of hills in
Dartmoor. Id.
exilis. On rocks whose surfaces lie
nearly even with the ground, on the sides
of hills the soil of which is peat earth, in
Dartmoor. Id.
vulpinus. Branches of trees and heath in
Dartmoor, and elsewhere. Hudson,
YOL, I. P

J Upon broken sand-
* LICHEN stictoceros. banks in the war-
t articulatus, var. fren opposite Ex-
J mouth. Mr. Brodie.
ULVA pavonia. Found by Mr. Stephens on the
rocks near Exmouth, plentifally. Gibson.
pluntaginea. Dawlish. Mr. Pigott.
coccinea. Rocks near Plymouth. Hudson.
dichotoma. Shores of Devon. Id.
narrow var. Dawlish. Mr. Pigott.
plumosa. Rocks near Exmouth. Hud-
son. At Ilfracombe. L. W. D.
filiformis. Rocks at Ilfracombe. L. W. D.
At Plymouth. D. T.
rubens. Rocks at Plymouth. Id.
Fucus Hypoglossum. Teignmouth. Mr. Pigott.
Plymouth. D. T.

* Mr. Brodie has been so obliging as to favor me with some specimens of this Lichen gathered at Exmouth, and I really can find in them no difference from the common *L. prunastri*. The black spots about the ends of the shoots, which led Dr. Smith to describe it as a new species, are, as was first pointed out to me by Mr. Edw. Forster, jun. common to that plant in all situations, though overlooked, till discovered by Mr. Brodie. D. T.

 \ddagger In this very extraordinary variety, as different from *L. articu*latus in habit as in place of growth, the stems are of the thickness of the little finger, and un formly inflated, not in that interrupted manner which gives the plant its common jointed appearance. My friend, Mr. Brodie, is of opinion that it is a distinct species, and, when the fruit is discovered, it will probably be found to be so. *D.T.*

Fucus ovalis. Rocks near Exmouth. Goodenough.
Dawlish. Mr. Pigott. Common all
along the Coast. D. T
dasyphyllus. Dawlish. Mr. Pigott
obtusus. Near Exmouth. Goodenough.
Here and there in the County. Hudson,
ligulatus. Among the rejectamenta of the
sea at Exmouth. Goodenough.
tamariscifolius.
tamariscifolius. } Rocks in this County. Hudson.
discors. Sidmouth. Stackheuse.
membranaceus. Found on a Limpet on the
shore at Sidmouth. Id.
bulbosus. On the shores of this County. Id.
——— laceratus. S. At Ilfracombe. Goodenough.
bifidus. At Stonehouse, near Plymouth.
Mr. E. Forster, jun.
ciliatus. 8. Abundantly on the rocks at
Ilfracombe. Goodenough.
s, & ζ. Exmouth. Sir T. Frankland.
crispus. Z. Common at Ilfracombe. Goodenough,
pygmæus. Ilfracombe : Sidmouth; and
elsewhere on the Coast. D. T
corneus. a, & B. Near Exmouth. Goodenough
J. Ilfracombe. Id
coronopifolius. Rocks in this County. Goodenough.
j. Louis and the second s

211

Fucus tomentosus. Found frequently at	Exmouth.
	Sir T. Frankland.
tuberculatus. Ilfracombe.	0
capillaris. Submersed rocks	, here and
there.	Hudson.
kaliformis. Exmouth. Goodence	ough. Daw-
lish. Mr. Pigott. About	Plymouth;
and Mount Edgecombe.	D. T.
flagelliformis. Dawlish.	Mr. Pigott.
CONFERVA byssoides. Among the reje	ctamenta of
the sea at Exmouth.	Goodenough.

SALICORNIA herbacea. β . γ . δ . In a little salt marsh, on the East side of Poole.

Mr. Stonestreet.

** fruticosa*. On the sea-shore all the way from Weymouth turnpike to Rhodipole. Mr. Yalden.
 Снака bispida. In the bog-pools on Wareham, and Poole, or Canford Heath. Pultency.

• Were this plant really found here it would hardly have escaped the notice of Dr. Pulteney, in his "Catalogue of the more rare plants of Dorsetshire." Indeed I have every reason to believe that we have only one species of Salicornia in this kingdom, and that what has been called S. fruticosa by British authors is in reality only a large variety of S. berbacea, which varies very much in size. D. T. VERONICA montana. By the river side under Hod hill. Pulteney.

PINGUICULA *lusitanica*. On the heath near Sandford bridge, not far from Wareham: on the heath near the three-mile stone

between Wimbourn and Poole: near the first fleet at Patnam: by the roadside about the midway between. Woodberry Hill and Wareham: in Purbeck, and many other places: on the heath between Wimbourne and Ringwood: by the roadside going from Corfe Mullein Schoolhouse to Poole: much more common than *P. vulgaris*.

Id.

Id.

UTRICULARIA minor. In a bog near the threemile stone between Wimbourne and Poole.

IRIS fatidissima. Abundant in the chalky parts, in shady lanes and under hedges. Pulleney. Near Broadway and Blandford. Mr. E. Forster, jun. About Weymouth. D. T.

SCHOENUS mariscus. At Weymouth, by the Fleet, in ditches communicating with the salt water. Pulteney.

 albus. Common on the bogs of Wareham, Canford, and Poole heaths. Id.
 CYPERUS longus. It does not appear that it has been found in Purbeck, as mentioned in

Ray's Synopsis, although diligent enquiry after it has not been wanting. Pultency. SCIRPUS acicularis. Boggy pools on the heaths, but not common. Id. ----- sylvaticus. By the river Stour, in many places near Blandford, as near White Cliff mill. Id. ALOPECURUS bulbosus. Salt marshes near Weymouth. Mr. Lambert. MILIUM lendigerum. Fields by Radipole, near Weymouth. Pulteney. Near Weymouth Turnpike Gate. Dr. Maton. * AGROSTIS maritima. With. Not uncommon on the sandy shores : on the North shore at Poole; at Swanage; and elsewhere. Pulteney. setacea. First sent to Mr. Curtis by Robert Squibb, from Piddletown Heath. Pulteney. On the higher Downs of this county. Withering. Sea coast near Weymouth. Duchess Dowager of Portland. AIRA canescens. On the sandy shores of Dorset : on the North shore near Poole. Pulteney.

* Dr. Pulteney agrees with Dr. Withering, in considering this plant a distinct species, and observes that its most distinguishing characteristic is the length of the vaginal leaf, which extends beyond the panicle. Dr. Smith, however, is of a different opinion, and does not appear to have noticed it in the Flora Britannica, even as a variety.

Pox procumbens. Not uncommon on the Coast:	
about Radipole, near Weymouth : about	
Poole Harbour, and elsewhere. Pu	lienev.
Festuca bromoides. On dry pastures, not uncom-	j.
mon; and sometimes on walls.	Id.
uniglumis. Found by Mr. Lightfoot	
in Portland.	Id.
BROMUS pinnatus. Common on the chalky	
Downs; frequently among the furze,	
and on dry banks in chalky lanes. Pul-	
teney. Cornfields near the Turnpike Gate,	
Weymouth. Dr. Maton. Near Broad-	
way, and Blandford. Mr. E. Forster	r. iun.
AVENA fatua. Near Weymouth. Rev. Dr. Gooder	
pubescens. On Hod Hill, and similar	0
	lteney.
ARUNDO epigejos. Wet ditches about Weymouth;	<i></i>
and in Purbeck : in several places in the	
	Id.
calamagrostis. On the borders of the	
wood, and in the hedges of the inclosures	
on the right hand, going between Fiddle-	
ford and Sturminster Newton. Pulteney.	
Road sides near Weymouth. Dr. M	laton.
ELYMUS arenarius. On the coast, not common.	
	teney,
HORDEUM maritimum. Not uncommon in the	
marshes near the sea, and on the cliffs	

.

216

Id.

TRITICUM loliaceum. On the beach at Swanage ; and at Poole, in the Harbour. Pulteney. On the Pier at Weymouth. Mr. E. Forster, jun. POLYCARPON tetraphyllum. Isle of Portland. Hudson. On the neck of the Isle of Portland, close to the shingly beach. Rev. Dr. Goodenough. On Chesil Bank. Mr. Lambert. DIPSACUS pilosus. Sparingly in Purbeck; and in several places in the Vale of Blackmoor. Pulteney. RUBIA peregrina. Hedges in Purbeck, but sparingly : hedges between Whitchurch and Milbourn St. Andrews. Pulteney. In Portland. Mr. Lambert. At Hod hill, on the side next the river, in the Parish of Stourpain, near Blandford : Wareham, on a mud wall; and at Somerpill, near a Chapel. Parkinson EXACUM filiforme. On Wareham Common, by the side of the road as you go to Corfe Castle. Rev. Dr. Goodenough. Common on the sandy parts of heaths, especially on the borders of pits, pools,

ditches, and bogs: about a mile from

Sherford Bridge, in the road to Ware-

ham : on Pool heath, and by the fleets :

in the way from Corfe Mullein to Poole :

i	Purbeck : heaths	between Wimbourne	
		t Christchurch; and	
	lorton, plentifully.		uliency.
		oole, and Wareham	
		where the soil has	
		utting turf: near the	
		way between Wim-	
	ourne and Poole.	j • • • • • • • • • • • •	Id.
Potamoge	TON gramineum.	In the river Stour,	2000
ar	id ditches communi	cating with it.	Id.
RUPPIA m	aritima. Ditches	on the coast: at	
P	pole.		Id.
ANCHUSA	sempervirens. Spari	ingly in some hedges	
	Damary farm, Bla		Id.
Asperugo	procumbens. Ch.	alky roads by the	
		forgotten the place.	
		Rev. G. C	Trabbe.
CAMPANUL	A patula. In the	woods above Corfe	
	ullein: and lanes r		
		Rev. Sir H. P	arker.
IMPATIENS	noli-tangere. In t	he grove at Dean's	
		apparently of natu-	
	growth.		lteney.
VIOLA birta	. Not uncommon	in woods and under	
	lges. Pulteney.		
	·	Mr. E. Forster	· 1 UT.
VERBASCUM	Blattaria. See	n in many places in	

orchards and hedges about the villages,	
as at Spetisbury.	ulteney
RHAMNUS catharticus. Not uncommon in the	
Chase, and the lanes adjoining : hedges	
about Blandford, and the neighbouring	
villages, Tarrant, Monkton, Crawford,	
Spetisbury, &c.	Id.
THESIUM linophyllum. Very common on the	
chalky, and especially the upland Downs :	
on Hod, and Hambledon hills: on the	
Race Down, Pimpern Down; and in	
Nutford Field, near Blandford.	Id.
CHENOPODIUM hybridum. In a corn-field near	
Blandford; and on waste grounds near	
Poole.	Id.
glaucum. I observed a large quan-	
tity of this two years successively on	
rubbish and dung near Blandford; but it	
was lost the third year.	Id.
SALSOLA fruticosa. Plentiful on Chesil Bank:	
a large quantity opposite Fleet House:	
abundantly at the Windmill Point, Poole :	
and elsewhere on the coast.	Id.
Cuscura europæa. On nettles about Blandford:	
on beans in Pimpern cornfield; and on	
vetches.	Id.
GENTIANA pneumonanthe. In Purbeck; and on	
the moist ground on heaths.	Id.

GENTIANA campestris. On the Downs near the Coast : on Pimpern Down, near Blandford. Pulteney. BUPLEURUM rotundifolium. Cornfields on Langton farm, near Blandford : between Spetisbury and Almer. Id. ----- tenuissimum. Fields near Radipole, Weymouth. Dr. Maton. SIUM latifolium. In the Stour, in many places: in the Froome and the Piddle, near White Cliff, by Blandford. Pulteney. SISON segetum. Langton cornfield, near Blandford. Id. OENANTHE pimpinelloides. In the lane, and in the copse adjoining, going from Sturminster Marshal field to Lytchet: about Weymouth, and Poole. Pulteney. Between Bincombe and Broadway. Mr. E. Forster, jun. PIMPINELLA magna. More common in the woods of Dorset, than in some of the midland Counties, yet sparingly wherever I have seen it. Pultener. SAMBUCUS Ebulus. In Spetisbury Town Street; and in hedges above the village near the Rings. Id. LINUM angustifolium. Purbeck: Portland; and Abbotsbury. Id.

DROSERA longifolia. anglica.

On Wareham, and other heaths: on a bog half a mile from Wareham towards Sandford Bridge.

Mrosurus minimus. On thatched houses : in old gravel pits : and among the corn, but not very common: Langton fields near Blandford.

FRITILLARIA Meleagris. I have been informed, on what I considered as good authority, that this plant grows in the wood on the left hand between Frittleford and Sturminster Newton.

TULIPA sylvestris. Some years ago it grew in the hollow way, near the Church at Melbury, near Shaftesbury. The roots were more than a foot deep in the earth. These plants answered exactly in the hairiness at the base of the stamens. I have seen it elsewhere, but always considered it as an outcast of the garden, and altered by neglect of culture.

ASPARAGUS officinalis. Very common on the Chesil Bank: on the shores and marshes at Poole Harbour, and especially about the North Haven Sandbanks, Pulteney.

Id.

Id.

Pulteney.

Id.

Near the Ferry, and extremity of Portland	
Island.	D. T.
CONVALLARIA multiflora. In the hanging woods	
	ulteney.
Acorus Calamus. In the river Stour, in two or	5
three places between Blandford and Stur-	
minster Newton.	Id.
JUNCUS acutus. In ditches in the sea marshes;	
and about the sandhills, not very uncom-	
mon.	Id.
RUMEX maritimus. On the sea coast in many	
places. At Poole; and at Wareham.	Id.
COLCHICUM autumnale. Plentifully in St. Mary	
Blandford Meadows on the East side of	
Mill Ditch; and in many other places in	
the meadows washed by the course of the	
Stour. Pulteney. About Spetisbury. Dr. 1	Maton.
ALISMA ranunculoides. Between Wareham and	
	ilteney.
CHLORA perfoliata. Not uncommon on the chalky	
Downs. On Hod, and Hambledon	
Hills: in the Isle of Branksea; and in	
Purbeck: in Nutford field, near Bland-	
ford.	Id.
DAPHNE Mezereum. In divers parts of Cran-	
	Id.
POLYGONUM Bistorta. Moist Meadows about	

Sturminster Marshall, and Shelborne,	
and Moreton. P	ulteney.
* PARIS quadrifolia, Wood at Turnworth, but	
sparingly: hanging woods at Ashcomb:	
Broadly and Morgan's wood, near	
Blandford.	II.
CHRYSOSPLENIUM alternifolium. On bogs with	
C. oppositifolium on heaths : in the hollow	
way about the Spring at Melbury, going	
to Shaftsbury,	Id.
SAXIFRAGA bypnoides. In the gardens and grounds	
at Stour Head, the seat of Sir R. Hoare,	
Bart. but I cannot ascertain it to be	
really a native.	Id.
SAPONARIA officinalis. Lane by the turnpike road-	
side near Milbourne St. Andrews ; and in	
many other places, in the hedges of	
Orchards about villages.	Id.
ARENARIA tenuifolia. On Pimpern Down, near	
the three mile stone : on the race ground.	Id.
SPERGULA subulata. On the more dry and sandy	
parts of the heaths of Wareham, Poole,	
and Canford.	Id.
EUPHORBIA peplis. Among sand at Bridport.	
Mr. J.	Sims
* Dr. Pultener charmen de la l	

* Dr. Pultency observes that he has seen an acre of ground in a wood thickly overspread with this plant; and the next year, at the same time, not a plant to be seen.

EUPHORBIA Portlandica. On the Eastern	
side of Portland, and on the Isth-	
mus, in divers places all the way to	
Abbotsbury: on the coast in Purbeck, in	
Langton Parish, Tineham, and at Lull-	
worth. Pulteney. Rocks in Portland,	
immediately under the Bow and Arrow	
Castle. Mr. B	ryer.
platyphylla. B. Plentifully in a field	
of Vetches on Langton Farm, near	
Blandford. Pult	eney.
Pyrus torminalis. Woods about Lytchet : Char-	
borough; and Henbury: Broadly Wood,	
near Blandford, and many others.	d.
Aria. Not unfrequent in the County:	
woods in Purbeck : many parts of Cran-	
bourne Chace, as about West Lodge, and	
about Hanley.	d.
TORMENTILLA reptans. In several places in Pur-	
beck: in the closes at New Bridge, be-	
tween Ringwood and Wimbourne, under	
Hod Hill.	d.
GEUM rivale. At Frome near Evershot, com-	
mon. Rev. G. Cra	bbe.
DELPHINIUM Consolida. In several fields near	
Blandford, and elsewhere, but sparingly.	
Pulie	ney.

.

ADONIS autumnalis. In several cornfields near Blandford: in a field opposite the first mile stone on the Dorchester Road: in Stour Paine field, on the borders of the camp-down. *Pulteney*.

RANUNCULUS parwiflorus. Common in cornfields, and on fallow lands, some of which near Blandford I have seen over-run with it.

HELLEBORUS viridis. Wood above Tolland Royal, to the Eastward, in considerable plenty; and in other parts of the Chace. Id MENTHA Pulegium. On the Heath below Woodbury Hill: in many places on the heathy

ground beyond Wimbourne. LEONURUS Cardiaca. About Dunghills; probably the outcast of gardens.

THYMUS Nepeta. Not uncommon in lanes, and borders of fields : Lane near Old Warren, and elsewhere about Blandford.

Scutellaria minor. Wareham Heath; and in Purbeck, sparingly. Id. MELAMPYRUM arvense. In Bere Field; observed

for two years, but afterwards lost. Id. LATHREA squamaria. In a very thick part of the Chace, near West Lodge, only two

100ts found. Pulleney. Near Chettle.

Rev. W. Chaffin.

VOL. I.

Q

225

Id.

Ich

Id.

Id.

ANTIRRHINUM Cymbalaria. Frequent on old	
Garden Walls. Pulte	ney.
spurium. Fields on the right hand	2
side of Stickland Lane; and in Langton	
and Pimpern cornfields : in the cornfields	
on the side of * Hambledon Hill. Id	-
minus. Bere Cornfields : Langton	
fields near_Blandford. Id	•
OROBANCHE minor. Among Vetches and clover	
in Pimpern field near Blandford. Id	•
ALYSSUM sativum. Flaxfields about Bridport,	
and Lyme. Hud	\$072.
incanum. In the year 1766 I found a	
considerable quantity of this plant near	
the rope walk at Weymouth, and on the	
spot where Gloucester Row and the	
royal Palace now stand. It was lost	
in three or four years after that period.	
Pulter	103.
Common on the cliffs,	
COCHLEARIA officinalis. as well as the sandy	
COCHLEARIA officinalis. 	,
Lobel. Purbeck Cliffs. Pulteney. Near	
Weymouth. Dr. Mate	otla
CRAMBE maritima. Not uncommon on the sandy	

* Dr. Pulteney found in this place a great quantity of A. spurium with flowers of the structure of the *Peloria*.

shores: On Chesil Bank; and abou	ıt
Weymouth: on the Purbeck coast; and	đ
at the North Haven, about Poole. Pul	
T 1 1 1	ithering.
CARDAMINE amara. Moist and marshy meadows	3
by the river Stour, especially those liable	
	Pulteney.
TURRITIS glabra. Old walls at Dorchester, very	
rare.	Id.
birsuta. Dry banks, and old walls, not	
common.	Id.
BRASSICA oleracea. Sea Cliffs at Purbeck; and,	14.
as I have been informed, on the Cliffs of	
Portland Isle.	Id.
ERODIUM maritimum. Not only on the sands, but	1a.
on the cliffs above, about Poole at the	
fleets, and elsewhere.	Id.
GERANIUM pratense. Not unfrequent by the sides	La.
of the Stour and the Frome.	T]
LAVATERA arborea. Recorded by Ray as a native	Id.
of Portland and Chesil Bank, where it is	
still found	Id.
PISUM maritimum. On the sand and naked pebbles	10.
of Chesil Bank, running from Portland to	
Abbotsbury: on the North Shore, and	
Factward of it at Deal 1 1	T 1
LATHYRUS Aphaca. Nutford cornfield, by White	Id.
Cliff: Langton farm, and in the coppice :	
Q 2	

in Spetisbury field, and other places; as about Weymonth. Pulteney. LATHYRUS Nissolia. Borders of Nutford cornfield, and in Cornfields about Weymouth. Id. --- sylvestris. In the Vale of Blackmoor : hedges by the road side near Milbourne St. Andrew's: at Encombe, the seat of W. M. Pitt, Esq. in the Vale near the sea. Pulteney.' Near Bridport. Mr E. Forster, jun. VICIA sylvatica. In several places in the County ; but particularly in a hedge going down Stoke Hill, from Bull-barrow. Pulteney. ----- lathyroides. Cornfield on the side of Hambledon Hill: in Purbeck; and about Wareham, but very sparingly. Id. ----- lutea. At Weymouth. Sir T. G. Cullum. * lavigata. Portland Island, among the shingles near Chesil Bank, and even flowering under the stones some depth. Sir T. G. Cullum. Lodmoor, near Weymouth. Hudson. ---- bithynica. In Purbeck. Hudson. In a field halfway between Weymouth and Portland Ferry, by the sea. Mr. Lambert.

* Dr. Pulteney remarks that Mr. Hudson took this plant for V. bybrida, till it was compared with the Linnwan Herbarium.

ASTRAGALUS glycyphyllos. Hedges in the borders	
of cornfields, but not common. I	ulteney
TRIFOLIUM ochroleucum. I have collected this in	
Dorset, but am uncertain where.	Id.
salt marshes near Poole, and elsewhere.	
Pulteney. About Weymouth. Sir T. G.	Cullum.
HYPERICUM Androsamum. Woods above Critchel	
House: about Henbury: in a wood	
above Grange in Purbeck ; and in Cran-	
bourne Chase.	ulteney.
Sonchus palustris. In some places by the Banks	
of the Stour : by ditches and rills on the	
coast, very rare.	Id.
LACTUCA saligna. Lanes about Cerne and Ever-	
shot, very rare.	Id.
Hroseris minima. Damary, and Langton corn-	
fields: Spetisbury fields.	Id.
CARDUUS tenuiflorus. Very common about Ware-	
ham; Poole; Christchurch; Weymouth,	
and elsewhere.	Id.
eriophorus. Here and there upon the	
higher Downs in the more Western part	
of the County; and about Revel's Hill.	Id.
pratensis. Boggy ground in Purbeck :	
at Merly, by the water side near the Tem-	
ple: more plentifully in the inclosures	

about New Bridge, between Wimbourne and Ringwood. Pulteney, SANTOLINA maritima. Near Poole. Hudson. Near Burton, by Bridport. Rev. J. Lightfoot. GNAPHALIUM rectum. Not uncommon on the heaths among furze and in the drier parts of woods. Pulteney. INULA Helenium. Vale of Blackmoor in many places, as Hasilbury, Sturminster Newton: St. Margaret's Marsh: a few roots in the parish of Langton, near Blandford. Id. ---- crithmoides. At Portland ; and about Weymouth : about Poole: in Purbeck, on the cliffs at Tineham. Pulteney. Chesil Bank. Dr. Malon. CINERARIA integrifolia. Top of Hod and Hambledon hills, but not in great plenty. Pulteney. PYRETHRUM maritimum. At Weymouth. Rev. Archd. Gooch. ANTHEMIS nobilis. About Lytchet: near Woodbury Hill: on the Common beyond the Wimbourne Turnpike going to Ringwood: and in many other places. Pultency. ORCHIS bifolia. B. In Meadows. Id. ustulata. On the chalky Downs, but not common: in Nutford field, near Blandford. Id. SATYRIUM viride. Dry pastures, but sparingly. Id.

OPHRYS Nidus avis. In the Chase, a little to the	
East of Rushmore Lodge : Little wood	
at Chettle. Rev. Mr.	Chaffin.
muscifera. Sent to me from the borders	
of the Chase, near Woodyats. P	ulteney.
apifera. On Hod and Hambledon Hills;	
and many other places on the Baulks in	
Nutford field, near Blandford.	Id.
SERAPIAS longifolia. Moist woods, not uncom-	
mon: Near Sturminster Newton: marshy	
Closes about New Bridge, between Wim-	
bourne and Ringwood.	Id.
grandiflora. In Purbeck, in the woods	
at Grange, and in the lane going up	
the hill above: in the Grove at the	
Down House, near Blandford; and in	
Little Wood at Chettle.	Id.
TYPHA angustifolia. Ditches communicating with	
the Stour near Sturminster Newton.	Id.
SPARGANJUM natans. In an old clay-pit on Corfe	
Castle Heath.	Id.
CAREX dioica. Bog in Purbeck.	Id.
pendula. By the Stour, between Blandford,	
and Durweston: by the Canal at En-	
combe, in Purbeck.	Id.
Pseudo-cyperus. Under hedges, in the	
ditches on the borders of heaths, but not	
common.	Id.

CAREX vesicaria. In Purbeck; and near New
Bridge, in the road to Ringwood. Pulteney.
ampullacea. Marshy boggy grounds, by
hedges and ditches. Id.
LITTORELLA lacustris. On Wareham Heath near
Sherford Bridge, and Sandford Bridge,
and between Wareham and Woodbury
Hill : near the road not far from the two
mile stone going from Wimbourne to
Poole. Id.
XANTHIUM strumarium. I found one plant only
by the river side, on a dunghill, at Ware-
ham, on the right hand of the bridge, in
the way to Stowborow. Id.
AMARANTHUS Blitum. On rubbish, and about
old Walls : at Weymouth, about the quay,
and elsewhere : at Poole. Id.
MYRIOPHYLLUM verticillatum. Boggy ditch near
Wareham. Id.
SALIX amygdalina. Common about Blandford. Id.
TAXUS baccata. Common in the Eastern part of
the Chase; and in a circuit of some miles
about Cranbourne. Id.
ATRIPLEX laciniata. On the sandy coast, but less
common than A. portulacoides. Id.
EQUISETUM sylvaticum. In the low, moist, and
shady parts of woods, but not common. Id.
fluviatile. Here and there on the

banks of the Stour, and about ponds, but most plentiful and of the finest growth in the crevices of the Cliffs on the Eastern side of Portland, and on some of the Cliffs in Purbeck; as at Encombe, and divers places from thence to Swanage. *Pulteney*. Near Bincombe. Mr. E. Forster, jun.

OSMUNDA lanarie. In the meadows near Sturminster Newton. Pulteney.

LYCOPODIUM immidatum. Common on the Heaths: between Woodbury Hill and Wareham: on Wareham Heath: between Sherford and Sandford Bridge: between Corfe Mullein and Poole; and between Wimbourne and Poole; and many other places. Id. ASPLENIUM marinum. Rocks and Cliffs in the Isle of Portland. Sir T. Cullum. In

Purbeck, in various places. Pulteney.

SCOLOPENDRIUM Ceterach. On the shady sides of old stone walls, and on bridges, not common. Id.

PILULARIA globulifera. By the road side, in

inundated places between Corfe Mullein and Poole, as about the eleventh mile stone: between Woodbury Hill and Wareham; and about Sandford Bridge near Wareham: about the saltpits at Poole, and at Lymington. Pulteney. PTEROGONIUM Smithii. Near Weymouth. Mr. Groult. LICHEN Jolithus. Stones at Maiden Castle, by Dorchester. D. T. vulpinus. Not uncommon in the woods, and on old pales, and on old apple trees in Orchards. Pulteney. In the close parts of the Chase and in various other woods: trees by the road side beyond Boy's lane plicatus. going to Shaftesbury: a-- articulatus. bout Rushmore: in the woods at Grange, in Purbeck: on trees in Milton Park, and in Clenston wood, near Blandford. Id. - Roccella. Rocks in Portland Island.

Earl of Dartmouth.

ULVA pavonia. Not uncommon on the rocks which are uncovered at low water: at Weymouth: Lullworth Cove: Swanage, and elsewhere. I have seen it in particular

spots in the month of June not larger than
the little finger nail, both growing and
dislodged in great quantities. Pulteney.
ULVA punctata. On the beach at Weymouth.
Mr. Stackhouse. A single specimen since
found by Dr. Maton.
rubens. Rocks in the Isle of Portland,
and near Poole. Iludson.
defracta. Not unfrequent at low water on
the Beach at Weymouth. Col. Velley.
elminthoides. In great abundance on the
Rocks off the Beal, at the extremity of
Portland, at very low water. Id.
Fucus sinuosus. B. Shores of the Isle of Portland. D. 7.
bypoglossum. At Weymouth, among the
of growing anong the
rejectamenta maris. Pulteney.
rejectamenta maris. Pulteney.
rejectamenta maris. Pulteney. ————————————————————————————————————
rejectamenta maris. Pulteney. β. A common parasitical plant on the stems and tips of Fuci at Poole. Stackhouse.

Fucus ligulatus. Among the rejectamenta at
Weymouth, and Portland Island. Goodenough.
fibrosus. At Weymouth, and elsewhere.
<i>Pulteney.</i> <i>discors.</i> Weymouth. Mr. Stackhouse.
abrotanifolius. In the sea at Weymouth.
Mr. E. Forster, jun. —— tamariscifolius. At Weymouth: Port-
Land D 1
selaginoides. Among the rejectamenta of
the sea at Weymouth, and the Isle of
Portland. Goodenough.
mucronatus. In the sea at Weymouth, in
great abundance. Mr. Bryer.
esculentus. Weymouth. Mr. Stackhouse.
alatus. β . Shores of the Isle of Portland. D. T.
ceranoides. Coast at Swanage, and, I be-
lieve, at Poole also. Pulteney.
bulbosus. Rocks about the Isle of Portland. Id.
phyllitis. Sparingly on the Coast. Pulteney.
Plentifully at Portland Head. Stackhouse.
ciliatus. S. In the Isle of Portland, rare.
Goodenough.
pinnalifidus. β. & δ. Weymouth. D. T.
pygmæus. Isle of Portland. Goodenough.
At Weymouth, on stones about the
quay, and under the look-out, common.
Pulteney.

Fucus crispus. E. Lyme Regis. D. T. Wey-
mouth. Mr. Stackhouse.
ζ. Lyme Regis. D. T.
mammillosus. d. Weymouth. Mr. Stackbouse.
corneus. y. In the Isle of Portland. Goodenough.
coronopifolius. Rocks on the coast. Pul-
teney. In the sea at Weymouth. Mr.
E. Forster. jun. Shores of the Isle of
Portland. D. T.
plumosus. Common on the Dorset shores.
Pulteney.
tomentosus. Portland, but rare. Id.
tuberculatus. Abundantly on the rocks at
the Bill of Portland below low water
mark. Mr. Bryer.
pinastroides. In great quantities at Wey-
mouth. Pulteney. Lyme Regis. Mr. Pigott.
narrow variety found at Weymouth by
Mr. Bryer. kaliformis. On the beach at Weymouth,
capillaris. At Weymouth. Pulteney. pedunculatus. Rocks about the Isle of
D 1 J
Fortland. Hudson. ——fruticulosus. On the stems of the larger
Trainet IV
Fuch at weymouth. $D. \mathcal{T}.$

CONFERVA corallina.	Very frequent near Wey-
mouth.	Mr. Stackhouse.
byssoides.	Among the rejectamenta at
Weymouth.	Goodenough.
parasitica.	Principally found on the
Fuci.	Pultency.

WE are not aware that any thing has hitherto been published respecting the Botany of Durham, and therefore feel a particular satisfaction in finding our Catalogue of this County equal to that of almost any other in the Kingdom; for which we are entirely indebted to the kindness of Mr. Harriman, and Mr. Winch, the latter of whom furnished not merely the habitats followed by his own name, but those also on the authority of Mr. Fenwick, Mr. Weighell, Mr. Waugh, and Mr. Thornhill. There is perhaps no part of England where the study of Botany, and especially the class Cryptogamia, has of late years been more sedulously cultivated; and we regret that in our list, copious at it is, we have, from the nature of our work, been obliged to omit many recent discoveries made among the Mosses by Mr. Winch, and among the Lichens by Mr. Harriman; a circumftance, however, that we the less lament, as, at the request of that most useful and respectable body, the Literary and Philosophical Society of Newcastle,

Ν

Mr. Winch has at this time in the Press a complete Catalogue of the Plants of Northumberland and Durham, where he will give a more full and accurate account than we could hope to do, of the additions made by himself and his friends to the British Flora.

CHARA bispida. B. Hell Kettles, near Darlington. Robson. ----- flexilis. At Cocken. Mr. Weighell. VERONICA montana. Dinsdale wood, near Darlington. Robson. Banks above Gateshead. Mr. Winch. VALERIANA dentata. Fulwell Hills : and Cocken. Id. SCHENUS Mariscus. Hell Kettles, near Darling-Robson. ton. compressus. Near Darlington. Robson. Bogs between Ryhope and the sea. Mr. Winch. Teesdale Forest. Rev. 7. Harriman. - rufus. Between Southwick and Hilton Castle. Mr. Winch. Marshes on the Wear above Sunderland. Mr. Weighell. albus. Beamish. Mr. Winch. monoicus. On Cronkley Fell. Rev. J. Harriman.

Scirpus pauciflorus. In a bog below Hilton Castle; and another below Willington Steam Mill. Mr. Winch. Near Darlington. Rev. J. Fenwick. - sylvaticus. By the Skern, near Darlington. Robson. Woods, and banks of rivers, near Newcastle. Mr. Winch. ERIOPHORUM vaginatum. Beamish Moor. Id. ----- polystachion. Bog at Hilton Castle. Mr. Winch. Whey Sike, Teesdale Forest. Rev. J. Harriman. * PANICUM viride. Willington, and Sunderland Ballast Hills. Mr. IV eighell.

* The claim which the plants found upon the various hills near Sunderland, formed of ballast from all parts of Europe, have to be considered as natives of this County is at best questionable. As many of them, however, may be naturalized there, and may thence have obtained the appearance of indigenous, we thought it best to incorporate those which were previously known to belong to our Flora; but besides these. Mr. Robson in his Catalogue of " Plantæ rariores agro Dunelmensi indigenæ,' printed some years since for the use of his friends, has included also the following, which not having at present been detected in any other place, we have not ventured to add to our Catalogue. - Phalaris paradoxa. Hyoseyamus albus. Convolvulus tric lor. Oenothera biennis. Reseda odorata. Euphorbia Tithyma'oides. Argemone Mexicana. Nigella arvensis. Ranun-ulus muricatus. Lejidium sativum. Alyssum incanum. Ornithopus compressus. Scorpiurus vermiculatus. Vicia Benghalensis. Irif lium M. indicum. Med cago prostrata. Anthem. s valentina. Centaunea galactites? Gannabis sativa. Calendula officinalis. VOL. I.

R

PHLEUM paniculatum. Sunderland Ballast Hills. PANICUM sanguinale. Mr. Weighell. crinitum. AGROSTIS Spica venti. Sunderland Ballast Hills. Mr. Weighell. Behind Beamish. Mr. Winch. MELICA nutans. Castle Eden Dean. Id. SESLERIA carulea. On both banks of the river Wear, and all the limestone hills between South Shields and Cleadon, plentiful. Mr. Winch. Banks of the Tees, near Egleston. Rev. J. Harriman. Poa procumbens. Quarry on the coast near Whitburn, along with Asplenium marinum. Mr. Winch. Houghton Bank. Mr J. Jansen. ---- humilis. Sea shore behind Sunderland North Mr. Winch. Pier. --- * glauca. Holy Island; and on the coast beteen Shields and Sunderland. Id. CYNOSURUS echinatus. St. Anthony's Ballast Hills. Mr. Thornhill and Mr. Waugh. FESTUCA vivipara. Near Egleston. Rev. J. Harriman. ----- bromoides. Walls above Gateshead.

Mr. Winch.

* I am not aware that this Grass is to be found in any author; but as it appears a very distinct species, and as I have received it under this name from my friend Mr. Winch, and from Mr. G. Don, as well as seen it so called in several British Herbaria, I have felt no scruple in introducing it by this appellation. D. T.

FESTUCA Inliacea. Near Lambton. Rev. J. Fenwick. AVENA fatua. Near Sunderland. Mr. Winch. Near Darlington. Mr. Brunton. ----- pubescens. Castle Eden Dean : Cleadon Hills; and pastures near New House, Weardale. Mr. Winch. ARUNDO epigejos. Castle Eden Dean. Mr. Winch. ELYMUS arenarius. Shore at Sunderland North Pier. Mr. Winch. The Snook, near Seaton. Rev. J. Dalton. Sand hills near Teesmouth. Mr. J. Jansen. South of Sunderland. Mr. Thornhill. HORDEUM maritimum. At Hartlepool, very frequent. Rev. J. Dalion. Coast near Seaton. Mr. Brunton. TRITICUM loliaceum. Coast near Seaton. Mr. Robson. In plenty upon the Cliff near Hartlepool. Rev. 7. Dalton. GALIUM Witheringii. Banks of Ponds at Lambton. R.v. J. Fenwick. ----- tricorne. Near Fulwell. Mr. R bson. Cornfields near the Tees Mouth. Mr. J. Jansen. boreale. Banks of the Tees at Whince Bridge: near Egleston. Rev. J. Harriman, POTAMOGETON fluitans. Ditches near Hell Kettles. R.v. J. Fenzuick. RUPPIA maritima. Near the Mouth of the Tees. R 2 Mr. Rolson.

PULMONARIA officinalis. Cliff wood, six miles West of Darlington. Mr. Robson. PRIMULA farinesa. Bogs above Hilton Castle : Whey Sike, Teesdale: near Painsher: Dalton; and Morton, common. Mr. Winch. About Darlington. Mr. Robson. Common near Egleston. Rev. J. Harriman. ANAGALLIS arvensis. y. Sunderland Ballast hills. Mr. Weighell. CAMPANULA latifolia. Woods near Beamish: Castle Eden Dean: every where near Middleton and Barnard Castle. Mr. Winch. VIOLA lutea. Cronkley Fell; and Teesdale Rev. J. Harriman. Forest. DATURA Stramonium. Sunderland Ballast hills. Mr. Winch. RHAMNUS catharticus. Ryhope Dene, edges of Mr. Weighell. rocks. RIBES alpinum. Woods near Darlington. Mr. Robson. Cliff wood. Mr. J. Jansen. _____ spicatum. By the Tees, between Piersbridge Mr. Robson. and Gainford. ---- petræum. Near Concliffe. Mr. Robson. Banks of the Tees, near Piersbridge. Mr. J. Jansen. Near Egleston. Rev. J. Harriman. Willington Ballast CHENOPODIUM hybridum. Hills. Mr. Winch. SALSOLA fruticosa.

GENTIANA verna. Teesdale Forest. Rev. J. Ha	rriman.
campestris. Cronkley Fell.	Id.
var. flore albo. At	
Whince Bridge.	Id.
ERYNGIUM campestre. Sunderland, and Willing-	
ton Ballast Hills. Mr.	Winch.
BUPLEURUM tenuissimum. By the Tees near	
Stockton. Mr.	Robson.
CAUCALIS daucoides. Fields near Whitburn.	
Mr. Winch. Hills near Fulwell. Mr. W	veighell.
DAUCUS maritimus. Sea Coast South of Sunder-	
land. Mr.	Winch.
OENANTHE pimpinelloides. Shore of the Wear	
between Sunderland and Hilton.	Id.
SCANDIX odorata. Near Harperly: Cawsey	
Wood : near Lanchester Ford : Beamish ;	
and banks of the Wear above Stanhope.	
Mr. Winch. About Egleston, common.	
Rev. J. Ha	rriman.
IMPERATORIA Ostruthium. At Newbiggen near	
Middleton; and in Teesdale Forest.	
Rev. J. Harriman. Banks of Dardree	
Burn, between Dardree Shield and the	
	Winch.
CARUM Carui. Willington Ballast Hills; and	
Fields at Beamish.	Id.
PIMPINELLA magna. Fields near Castle Eden	
Dean; and in Chester Lane.	Id.

SAMBUCUS Ebulus. Lane between Cawsey Hall, and Beamish Burn. Mr. Winch. LINUM usitatissimum. Sunderland Ballast Hills. Id. perenne. Baydales near Darlington. Mr. Robson. DROSERA longifolia. Whey Sike, Teesdale Forest. R v. J. Harriman. MYOSURUS minimus. Fields near Darlington. Mr. Brunton. GALANTHUS nivalis. Banks about Blackwell and Concliff. in situations which do not admit of its being the outcast of a garden. Mr. Robson. ALLIUM oleraceum. Baydales, near Darlington. Mr. Robson. Near Middleton. Reg. J. Harriman. Meadows near Silksworth. Mr. Weighell. ORNITHOGALUM luteum. Banks of the Tees near Greta Bridge. Mr. Robson. Near Barnard Castle; and Whorlton; and shores of the Tees near Egleston, and near Bishop Auckland. R.v. 7. Harriman. CONVALLARIA multiflora. In the lane between Stockton and Stainton. M. Robson. JUNCUS triglumis. Mell Fell. Mr. J. Jonsen. Moor beyond Cronkley Fell. Rev. J. Harriman. FRANKENIA lævis. Sunderland Ballast Hills. Mr. IT eighell. RUMUX sanguineus. Near Houghten le Spring Mr. Winch. ----- maritimus. Ballast on the Wear. Id.

TOFIELDIA palustris. Cronkley Fell; and Whey
Sike, Teesdale Forest. Rev. J. Harriman.
COLCHICUM autumnale. Field opposite Egleston.
Rev. J. Harriman. Near Darlington.
Mr. Robson.
ALISMA ranunculoides. In Hill-close-carr, two
miles west from Darlington. Robson.
Bogs at Cleadon. Mr. Weighell.
TRIENTALIS europæa. Allum Works near Carle-
ton Gisborough. Mr. Weighell. Wask-
erly Park, near Wolsingham. Mr. Thornhill.
EPILOBIUM angustifolium. Banks of the Tees
in Teesdale. Rev. J. Harriman. Rocks
above Feldon Lead Mill, two miles West
of Edmondbyers. Mr. Winch.
VACCINIUM uliginosum. Meldon Fell; and
Moor beyond Cronkley Fell. Rev. J. Harriman.
Vitis idaa. Cronkley Fell; and near
Egleston. Id.
Oxycoccos. Cronkley Fell. Rev.
J. Harriman. Beamish Moor. Mr. Winch.
ERICA vulgaris. var. flore albo. Cronkley Fell.
Rev. J. Harriman.
POLYGONUM minus. Morden Cars near Darling-
ton. Mr. Winch. Bistorta. Wood below Gateshead.
Mr. Winch. Near Egleston. Rev. J. Harriman.
viviparum. Frequent in meadows

near Middleton, and Burtryford, Wear-
dale. Rev. J. Harriman.
PARIS quadrifolia. Near Egleston. R v. J. Harri-
man. In deep shades near the Gunner's
Pool. Eden Dean. Mr. Winch.
ARBUTUS Uv. ursi. Cronkley Fell: Force Gaith
Scar, Teesdale Forest ; and near Caldron
Snout. Rev. J. Harriman.
PYROLA rotundifolia. North Branch of Castle
Eden Dean, plentiful. Mr. Winch.
Birch Carr. M. J. Jansen.
minor. Cocken Wood. Mr. Weighell.
Teesdale Forest. Rev. J. Harriman.
CHRYSOSPLENIUM alternifolium. In Baydales,
near Darlington. Mr. Robson. Near
Egleston. Rev. J. Harriman. Near
Croft Bridge. Rev. J. Dalton. Banks
of the Wear opposite Cocken. Mr. Thornhill.
SAXIFRAGAstellaris. Fgleston Moor. Rev. J. Harriman.
aizoides. Banks of the Whey Sike,
Middleton Forest. Id.
SCLERANTHUS perennis. Fields above Gateshead.
Mr. Thornhill.
SAPONARIA officinalis. Chester le Street new
bridge. Mr. Winch. Near Darlington.
Rev. J. Harriman.
SILENE anglica. Sunderland Ballast Hills. Mr. Winch.
conica. Hebburn Ballast. Mr. Waugh.

SILENE noctiflora. Sunderland Ballast. Hills. Mr. Weighell. In a field South of Cleadon Mr. Winch. Mill. STELLARIA Nemorum. Beamish, and Ravensworth woods. Mr. Winch. In Baydales; and Cliff wood, near Darlington. Mr. Robson. Cocken woods. Mr. Weighell. ARENARIA verna. On Cronkley Fell: about old lead mines in Weardale: near Cal-Rev. J. Harriman. dron Snout. SEDUM villosum. Sikes near Burley Ford, Weardale. Mr. Winch. Tcesdale Forest. Rev. 7. Harriman. Cronkley Fell. Mr. J. Jansen. ------ album. Abby Bridge near Barnard Castle. Mr. Robson. ----- rupestre. Walls about Darlington. Id. CERASTIUM tetrandrum. Sca shore near South Shields, and Sunderland, and Whitburn sands. Mr. Winch. PRUNUS Padus. Woods by the Tees. Mr. Robson. Woods near Beamish. Mr. Winch. PYRUS Aria. Wood near Whince Bridge. Rev. J. Harriman. Rosa villosa. Common on the Banks of the Darwent. Mr. Winch. Near Darlington. Mr. J. Jansen. ---- tomentosa. Hamsterly. Mr. Winch.

RUBUS saxatilis. North Branch of Castle Eden Dean. Mr. Winch. ------ chamamorus. Cronkley Fell: Knoutberry Fell, near Egleston. Rev. J. Harriman. POTENTILLA fruticosa. On the banks of the Tees between Egleston and Widdy Bank in 'the Forest. Rev. 7. Harriman. At Whince Bridge. Mr. Winch. Mr. Lawson observed thousands of these plants by Mickle Force in Teesdale. Ray. aurea. Whince Bridge, Teesdale. Rev. J. Harriman. GEUM rivale. Every wood near Newcastle. Mr. Winch. - — β. Castle Eden Dean. Mr. Winch. Near Darlington, frequent. Mr. Robson. DRYAS octopetala. Cronkley Fell. Rev. J. Harriman. PAPAVER hybridum. About Durham. Mr. Robson. Near Fulwell, and Sunderland. Mr. Winch. ----- somniferum. Willington Ballast Hills. Id. CISTUS marifolius. Cronkley Fell. Rev. J. Harriman. DELPHINIUM Consolida. Hebburn Ballast Hills. Mr. Winch. THALICTRUM alpinum. Cronkley Fell; and Caldron Snout. Rev. J. Harriman. minus. Sea Shore between Shields and Sunderland, plentiful. Mr. Winch. Common near Hartlepool. Rev. J. Dalton.

THALICTRUM majus. Baydales, near Darlington.
Mr. Robson. Hedges leading to Barnard
castle. Mr. Winch. Tees Banks op-
posite Egleston. Rev. J. Harriman.
RANUNCULUS Lingua.
parvislorus. } Near Darlington. Robson.
TROLLIUS europaus. Woods in Weardale: Beam-
ish; and bogs near Morton. Mr. Winch.
Near Egleston. Rev. J. Harriman.
HELLEBORUS viridis. Banks of the Tees near
Whorlton. Id.
fætidus. Banks of the Tees below
Winston Bridge. Rev. J. Harriman.
Woods in Weardale. Mr. Winch.
Ajuga alpina. It is erroneously stated in English
Botany that this plant has been found in
Durham by Mr. Robson, for the speci-
men he sent to Sowerby was from his
own garden. Id.
TEUCRIUM Chamadrys. On a Bank near the
Gateshead Engine. Id.
MENTHA Pulegium. Pond between Whinston
and Gainforth. Id.
BARTSIA alpina. Banks of the Whey Syke, Tees-
dale Forest: near Middleton. Rev. J. Harriman.
MELAMPYRUM sylvaticum. Banks of the Tees
above Middleton: Egleston wood,

Rev. J. Harriman. By Whince Bridge,
Teesdale. Mr. Brunton.
LATHRÆA squamaria. In a wood near Gainford.
Mr. Robson. Egleston wood. Rev.
J. Harriman. Cocken Woods. Mr. Weighell.
ANTIRRHINUM spurium. Sunderland Ballast Hills. Id.
repens. Hebburn ballast. Mr. Winch.
minus. Willington Ballast Quay:
Fulwell hills near Sunderland. Id.
DRABA incana. Cronkley Fell. Rev. J. Harriman.
Walls in Teesdale. Mr. Brunton.
ALYSSUM sativum. Willington Ballast. Mr. Winch.
LEPIDIUM latifolium. Near Seaton; and by the
river Were, near Durham Abbey. R bson.
At Hartlepool. Rev. W. Wood.
ruderale. St. Anthony's Ballast Hills.
Mr. Winch.
THLASPI arvense. Lanes about Gateshead. Id.
campestre. β. On Willington Ballast
Hills. Id.
alpestre. Near Whince Bridge. Rev. J. Harriman.
COCHLEARIA officinalis. Marsden Rocks, and sea-
shore. Mr. Winch.
shore. Mr. Winch. ———— y. Coalcleugh. Id.
y. Coalcleugh. Id.

Robson. Banks of the Were between Framwell Gate and New Bridges. Mr. Winch. CARDAMINE amara. Woods near Egleston, and Beamish. Id. SISYMBRIUM sylvestre. Shores of the Tyne near Dunstan; and Ballast Hills. Id. ERYSIMUM cheiranthoides. Sunderland Ballast Id. Hills. TURRITIS hirsuta. Sea Banks below Whitburn: Castle Eden Dean. Mr. Winch. Banks of the Tees opposite Egleston: walls above Middleton. Rev. J. Harriman South side of Bolden, Tunstall, and Cleadon Hills. Mr. Waugh and Mr. Thornhill. GERANIUM phaum. Woods about Darlington. Mr. Robson. bridge at Durham. Robson. Castle Eden, Walbottle, Scott's wood, and Denton Deans. Mr. Winch. --- pratense. Road side near Walbottle Dean : banks of the Wear at the new bridge; and banks of the Team. Id. ---- pyrenaicum. Hebburn Ballast. Id. ----- sanguineum. Foot of Castle Eden Dean, plentiful. Mr. Winch. Very common at Hartlepool. R.v. J. Dalton. LATHYRUS Aphaca. 7 Sunderland Ballast Hills. birsulus. Mr. Winch.

DURHAM:

VICIA sylvatica. Castle Eden Dean. Mr. Winch. Woods near Barnard Castle, Whorlton, and Egleston. Rev. J. Harriman. lathyroides. Sea shore South of Shields; and salt meadow below Gateshead. Mr. Winch. ASTRAGALUS glycyphyllos. Baydales. Mr. Robson. Abbey Bridge near Barnard castle. Rev. J. Harriman. Below Low Conniscliff, and in other places by the Tees. Robson. ----- bypoglottis. Near Seaton. Robson. St. Anthony's Ballast Hills; and seashore below Shields. Mr. Winch. TRIFOLIUM officinale. var. flor. albis. Willington Ballast; and on the Ballast Hills below Gateshead. Id. ochroleucum. Sunderland Ballast Hills. Mr. Weighell. - maritimum. Willington Ballast. Mr. Winch. Sunderland Ballast Hills. Mr. Weighell. _____ scabrum. Bishop Wearmouth Paddock. Id. ----- glomeratum. Sunderland Ballast Hills. Id. MEDICAGO falcata. Willington Ballast Hills. Mr. Winch. HYPERICUM and iosamum. Near Darlington. Mr. Robson. ----- dubium. Banks of the Tees near Middleton. Rev. J. Harriman. HIERACIUM paludosum. Near Barnard Castle,

Rev. J. Harriman. Castle Eden Dean. Mr. Winch-HIERACIUM prenanthoides. 3 Willington Ballast CREPIS fætida. 3 Hills. CREPIS fatida. Id. biennis. Sunderland Ballast Hills. Mr. Winch. Coast South of Sunderland. Mr. Waugh. CARDUUS tenuiflorus. Common about Newcastle. Mr. Winch. eriophorus. Hedge and quarry at Fulwell. Mr. Waugh. beterophyllus. Near Burtry-ford, Weardale. Mr. Winch. ---- pratensis. Castle Eden Dean : Teesdale Forest : South Shore of the Tyne, opposite Scott's wood. Id. GNAPHALIUM dioicum. Moor above Beamish : Gateshead Fell. Mr. Winch. Common in Teesdale Forest. Rev. J. Harriman. rectum. Woods and fields about Newcastle. Mr. Winch. Frequent in the County. Mr. Robson. ERIGERON canadense. Sunderland Ballast Hills. Mr Weighell. PYRETHRUM maritimum. At Whitburn. Mr. Robson. Sea Banks, and in a quarry South of Marsden Rocks. Mr. Winch. ANTHEMIS maritima. The habitat assigned to

this plant at Sunderland in Flora Britannica is an error of Mr. Robson's. Mr. Winch. ANTHEMIS nobilis. On Lints Green in front of Lints Green Hall between the house and the turnpike, plentiful. Mr. Waugh and Mr. Thornhill. tinctoria. On a Bank near the river Tees, not far from Sogburn. Ray. Not now to be found there. Mr. Robson. Sunderland Ballast Hills. Mr. Weighell. A single plant found by the roadside near St. Anthony's Colliery by Mr. R. Callender. CENTAUREA nigra. B. Near the Magazine at Walker : Walbottle Dean. Mr. Winch. ORCHIS ustulata. Near Darlington. Mr. J. Jansen. South side of Bolden Hills: sea banks at Ryhope : East side of Cleadon Hills. Mr. Thornhill and Mr. Waugh. SATYRIUM viride. Beamish woods, and the fields adjoining; and meadows near Moreton. Mr. Winch. Many places near Sunderland. Mr. Waugh. Near Middleton. Rev. J. Harriman. ---- albidum. Near Whince Bridge: and the Whey Sike, Teesdale Forest. Id. OPHRYS Nidus a-vis. Cocken, in the wood at the West side of Whartonshaugh; and in the dingle above the old windmill on the right hand side. Mr. Winch.

OPHRYS cordata. Gold Hill near Muggleswick;
and among the Ling in Charnberry, near
Egleston. · Rev. J. Harriman.
South branch of Castle Eden Dean ; and
together with the Cypripedium in the
North Branch; also above the Gunner's
Pool. Mr. Winch.
apifera. Rooker's Gill, near Fulwell.
Mr. Weighell. Fields on the Coast,
three miles South of Sunderland. Mr.
Thornhill. Near Ryhope Turnpike
Gate. Rev. J. Fenwick.
MALAXIS paludosa. Middleton in Teesdale. Mr.
J. Jansen. In Charnberry, near Egles-
ton, among Bog Moss. Rev. J. Harriman.
CYPRIPEDIUM Calceolus. The North Branch of
Castle Eden Dean. Robson.
SERAPIAS longifolia. Buttersby and Cocken Woods.
Mr. Weighell. Castle Eden Dean, on a
Bank opposite the old Soldier's House.
Mr. Winch.
TYPHA angustifolia. Corn Mill Dam near Team. Id.
SPARGANIUM natans. In the Team, plentiful. Id.
CAREX dioica. Teesdale Forest : bogs at Hilton ;
and on the moor near Beamish. Mr.
Winch. Near Darlington. Robson.
curta. Waskerley Park. Mr. Winch.
VOL. I. 6 S

Near Calberry Lead Mine near Middleton. Rev. J. Harriman. Weardale. Rev. J. Fenwick. CAREX pendula. Denton Dean, and Cocken Woods. Mr. Winch. By the Tees, near Croft. Mr. Robson. Gibside Woods. M. Thornhill. ----- capillaris. Whey Sike: Cronkley Fell; and Widdy Bank. Rev. J. Harriman. ----- fulva. Bogs below Hilton Castle. Mr. Winch. ----- extensa. Marsh at the foot of Hilton Burn to the East. Mr. Thornhill. Bog at Spencer's Quarry near Medomsley. Mr. Winch. Hartlepool. Rev. J. Dalton. ----- binervis. Heath west of Woodlands near Butsfield. Mr. Winch. distans. Marsh below Hilton Burn. Id. ----- rigida. Teesdale Moor near Caldron Rev. J. Harriman. Snout. ----- vesicaria. Bogs to the South of Castle Eden ; and ditches near Darlington. Mr. Winch. ampullacea. Bogs to the South of Castle Eden; and behind Willington Ballast Id. Hills. URTICA pilulifera. Ballast Hills, Sunderland. Id. SALIX Croweana. fam. In Weardale. Mr. Rev. J. Harriman. Winch. Teesdale. amygdalina. Hedges near Gilsland; and in Heaton Dean. Mr. Winch.

SALIX pentandra. Pockerley near 1	
Mr. Wnch. Near West Bold	
Cleadon. Mr. Thornhill an	d Mr. Wrough.
vitellina. Banks of the Team.	Mr. Winch.
rosmarinifolia. Banks of the I	Darwent,
near Ebchester.	Mr. Thornhill.
EMPETRUM nigrum. On Beamish mod	
every moor near Newcastle.	Mr. Winch.
RHODIOLA rosea. Maes Beck.	Mr. J. Jansen.
TAXUS baccata. Woods about Egleston.	
J. Harriman. Shores of the W	
low Hilton Castle; woods abov	e Der-
went Bridge, and Castle Eden	
plentiful, and undoubtedly wild.	
ATRIPLEX laciniata. Near Marsden Rock	
Winch Hartlepool.	
pedunculata. Hartlepool.	Mr. Robson.
EQUISETUM sylvaticum. Woods about	
castle.	Mr. Winch.
byemale. Woods above D	
Bridge. Mr. Winch. Castle Eden	
OSMUNDA lunaria. Fields near Marsden 1	
and moor above Beamish.	Mr. Winch.
LYCOPODIUM selaginoides. Near Middle	
Teesdale. Mr. Robson. Gateshead	
Mr. Winch. Cronkley Fell; and	
	J. Harriman.

259

\$ 2

LYCOPODIUM alpinum. Cronkley Fell; and near Egleston. Rev. J. Harriman. POLYPODIUM Phegopteris. Cawsey Wood. Mr. Winch. ----- D: yopteris. Cawsey Wood; and Walbottle Dean. Id. ----- Oreopteris. Cawsey wood: East side of Gateshead; and Beamish Moor. Mr. Winch: Woods about Darlington. Mr. Robson. ASPLENIUM viride. Near Middleton. Mr. J. Jansen. Cronkley Scar: near Egleston. Rev. J. Horriman. marinum. Rocks near Marsden Rocks. Mr. Winch. Black Hall Dean, West of Hartlepool. Rev. J. Dalton. Near Southwick. Mr. Brunton. PTERIS crispa. Rocks at Cocken. Mr. Wighell. Walls near Cronkley Fell: Teesdale Rev. J. Harriman. Forest. CYATHEN dentata. Cronkley Fell. Id. GYMNOSTOMUM fasciculare. Stopes in a rivulet Mr. Thornhill. near Edmondbyers. ANICTANGIUM ciliatum. Waskerley. Mr. Winch. TRICHOSTOMUM capillaceum. Bolden Hills; and Painsher wood. Id. ----- polyphyllum. Howden Burn. Mr. Thornhill. TORTULA rigida. Walls in Coppice Lane near Mr. Winch. Beamish Hall.

TORTULA tortuosa. Near Whitburn : Stanhope ;	
and Painsher. Mr.	Winch.
DICRANUM aciculare. Cawsey Wood.	Id.
B. Musc. Hib. Ravens-	
worth.	Id.
pellucidum. Cawsey Wood, in a rivulet;	
and Ravensworth Wood.	Id.
squarrosum. Beamish Moor.	Id.
POLYTRICHUM urnigerum. Gateshead Fell.	
NECKERA crispa. Cawsey wood: Castle Eden	
Dean: Woods about Derwent Bridge;	
and near Painsher. Mr. Winch. Woods	
near Sunderland, and Darlington. Mr. W	eighell.
curtipendula. South Bank of the	
Derwent, near Crook Oak. Mr. The	rnhill.
heteromalla. Near Kibblesworth. Mr.	
Thornhill. Ravensworth woods. Mr. 1	Vinch
BARTRAMIA Halleriana. Redwell Hill near	
Lanchester.	Id.
* pomiformis. B. Edmondbyers. Mr. Thom	
arcuata. North West branch of a	1 16138666
rivulet near Edmondbyers.	11
BRYTIM trichadas Cibaila NA	
punctatum. Bears fruit in woods about	Id.
Normani	7 2
HYPNIM pulchellum Common 1	
2 Childrey Wood.	Id.

* See König's Annals of Botany, I. p. 527.

HYPNUM myosuroides. Cawsey and Ravensw	orth
	Mr. Winch.
trichomanoides.	
bucens. Woodsabout Newc	astle. Id.
undulatum. Gateshead Fell:	and
woods about Newcastle.	Id.
7 cesdalii. Cawsey wood.	Id.
crenulatum. ¿Cawsey and Ravensw	orth
tenellum. S Woods.	Id.
plumosum. Trees in woods about N	Jew-
castle.	Id.
loreum. Cawsey and Ravensw	rorth
Woods: on the low parts of Gates	head
Fell; and near Lanchester.	Id.
palustre. Beamish Moor: and N	Wal-
bottle Dean.	Id.
dubium? Woods about Newcastle	e. Id.
cupressiforme. B. Musc. Hib. Beau	mish
Moor.	Id.
JUNGERMANNIA lanceolata. Woods near N	New-
castle.	Id.
emarginata. Gateshead Fell. Mr	. Thornhill.
purpurea. Gibside and Ravensw	orth
Woods.	Id.
reptans. Cawsey wood.	Mr. Winch.
LICHEN Jolithus. Middleton in Teesdale.	
Rev. J.	Harriman.
	head
Fell, on stone.	Mr. Winch.

LICHEN	s sanguinarius. pilularis. confluens.
	pilularis. > Gateshead Fell. Mr. Winch.
	confluens.
	caruleo-nigricans. Cleadon and Fulwell
	Hills. Mr. Thornhill. Moors in Tees-
	dale. Rev. J. Harriman.
	Oederi. Teesdale. Id.
	geographicus. Rocks at the Force of Tees;
	and on Cronkley Fell. Id.
	flavo-virescens. Moors near Middleton. Id.
	rupicola. Teesdale Forest. Id.
	calvus. Marble Rock, near Middleton. Id.
	baomyces. Gateshead Fell; and near
	Middleton. Rev. J. Harriman. Bea-
	mish Fell. Mr. Winch.
	escharoides. Near Muggleswick. Mr. Thornhill.
	polytropus. Teesdale. Rev. J. Harriman.
	luteo-albus.
	hymenius. } Near Egleston. Id.
terrarian gi terrarian gi	atro-rufus. Teesdale.'
	sponsus.
	salicinus. Trees at Cocken. Mr. Winch.
	WIT. W inch.
	chloroleucus. Fulwell Hills: Teesdale.
	chloroleucus. Fulwell Hills : Teesdale. Rev. J. Harriman.
	chloroleucus. Fulwell Hills : Teesdale. Rev. J. Harriman. inclusus. Teesdale Forest. Rev. J. Har-
	chloroleucus. Fulwell Hills: Teesdale. Rev. J. Harriman. inclusus. Teesdale Forest. Rev. J. Har- riman. Ravensworth Wood on Trees;
	chloroleucus. Fulwell Hills : Teesdale. Rev. J. Harriman. inclusus. Teesdale Forest. Rev. J. Har-

LICHEN ventosus. Teesdale Forest; and Cronkley
Fell Rev. J. Harriman. Walls near
Newbiggin, and Lanchester. Mr. Winch.
coccineus. On walls near Lanchester.
Mr. Winch. Rocks at the Force of
Tees, and Cronkley Fell. Rev. J. Harriman.
corallinus. Teesdale Forest; and Cronk-
ley Fell. Id.
Westringii.
Westringii. } Near Egleston. Id.
abietinus. Ach. Trees in Sharnberry Gill. Id.
subimbricatus. Cronkley Fell. Id.
concentricus. Cawsey Wood, on Stones.
Mr. Winch.
frustulosus. Muggleswick Fell. Mr. Thornhill.
gibbosus. Teesdale. Rev. J. Harriman.
corneus. Egleston Wood. Id.
athroocarpus. Ach. On Cronkley Fell. Id.
Acharii. In the bed of the Tees below
Cronkley Fell. Rev. J. Harriman.
Stones in Horsley hope Bourn. Mr. Thornhill.
Rev. J. Harriman.
exanthematicus. Banks of the Tees near
Mr. Morritt's Bridge. Id.
4
smaragdulus. (Endocarpon smaragdulum.
smaragdulus. (Endocarpon smaragdulum.

T

J. Harriman. Rocks below Wycliffe.

Mr. Thornhill. LICHEN quadricolor. Marble Rock at Middleton Burn. Rev. J. Harriman. gelidus. Near the Force of Tees; and Whince Bridge. Id. ---- luridus. Moors in Teesdale. Id. ---- crassus. Marble Rock at Middleton Burn. Id. ----- multifidus. On Beamish Moor; and rocks in Teesdale. Rev. J. Harriman. On stones in the Rivulet near the source of the Roman Aqueduct, which leads to the Station at Lanchester. Mr. Winch. ---- carnosus. Teesdale. Rev. J. Harriman. fulvus. Rocks between Shields and Sunderland. Mr. Winch. _____ gloliferus Gateshead Fell. Mr. Winch. Cronkley Fell. Rev. J. Harriman. ----- fragilis. Near Egleston. Id. tristis. Rocks near Blackton. Id. paschalis. Teesdale Moor. Rev. J. Harriman. Gateshead Fell. Mr. Winch. papillaria. Gateshead Fell; and heaths near the Roman Aqueduct, west cf Woodlands. Id. muscicola. Moors in Teesdale. Rev. J. Harriman. Near Muggleswick.

Mr. Thornhill.

LICHEN jubatus. Gateshead Fell. Mr. Thorn-
bill. Moors near Muggleswick. Mr. Winch.
exilis. Moors in Teesdale. Rev. J. Harriman.
plicatus. Waskerley Burn, near Wol-
singham. Mr. R. Wilson.
prunastri. Found in fruit in Gibside
and Ravensworth Woods by Mr. Thornhill.
pinastri. Firs near Gainford. Rev. J. Harriman.
glaucus. Teesdale. Id.
Islandicus. Cronkley Fell. Rev. J. Har-
riman. Gateshead Fell. Mr. Winch.
pulmonarius. Trees near Wycliffe. Mr.
Thornhill. Sheeplea Wood, Egleston.
Rev. J. Harriman.
glomuliferus. Near Egleston. Id.
lætè-virens.
scrobiculatus. Sheeplea Wood, Egleston. Id.
— lætè-virens. — scrobiculatus. — plumbeus. SheepleaWood, Egleston. Id.
scrobiculatus. Sheeplea Wood, Egleston. Id.
scrobiculatus. Sheeplea Wood, Egleston. Id. plumbeus. limbatus. affinis. Sharnberry, near Egleston. Id.
limbatus. affinis. } Sharnberry, near Egleston. Id.
limbatus. affinis. } Sharnberry, near Egleston. Id.
limbatus. affinis. } Sharnberry, near Egleston. Id. torrefactus. } Moors in Teesdale. Id.
— limbatus. affinis. torrefactus. polyphyllus. Sharnberry, near Egleston. Id.
— limbatus. affinis. Sharnberry, near Egleston. Id. — torrefactus. polyphyllus. polyphyllus. Rocks on the summit of
— limbatus. Sharnberry, near Egleston. Id. affinis. Sharnberry, near Egleston. Id. torrefactus. Moors in Teesdale. Id. polyphyllus. Rocks on the summit of Id. proboscidcus. Cronkley. Id.

266

Ń

DURHAM.

Winch. In a Burn on the North side of the Derwent opposite Gibside. Mr. Thornhill. LICHEN saccatus. Marble Rock, Middleton Burn. Rev. J. Harriman. ____ aphthosus. Teesdale, where it bears fruit. Id. resupinatus. Sheeplea wood, near Egleston. Rev. 7. Harriman. Stones in Cawsey Wood; and Walls behind Bea-Mr. Winch. mish Hall. borizontalis. Hag Crag, near high Force. Rev. J. Harriman. sylvaticus. Cawsey Wood. Mr. Winch. ----- granulatus. Trees near Hyshope, and Horsley Hope. Id. ----- fascicularis. Teesdale. Rev. J. Harriman. * ULVA laciniata. Sea Coast between Shields Mr. Thornhill. and Sunderland. ---- * defracta. ----- * elminthoides. ----- fistulosa. Sunderland Beach. Mr. Weighell. ----- purpurascens.

* I have taken the liberty of prefixing an Asterisk to some of the Fuci and Ulva in this list, where I have reason to suspect either the plant or its habitat to have been mistaken. Most of them rest upon the authority of the Herbarium of the late Mr. Weighell, now in the possession of my friend, Mr. Winch; and it is most probable that they were sent either by Mr. Wigg, or some other Botanist at a distance, but mixed by him with what he had himself collected at Sunderland, where he resided. D. T.

DURHAM.

ULVA plumosa. Whitburn. Mr. Weighell.
Fucus esculentus. Shores of this County. Mr. Winch.
membranaceus. Shields Beach. Id.
dentatus. On the shore, but rare. Id.
corneus.
* cartilagineus.
aculeatus.
plumosus. Shores of the County. Mr. Wincb.
* tomentosus,
clavellosus.
CONFERVA fluviatilis. Rivulet in Cawsey Wood.
Mr. Winch.
Mr. Winch.

CHARA flexilis. Gravel pits on Epping and Hen-
hault Forests. Mr. E. Forster, jun.
VALERIANA dentata. Cornfields near South End,
and Rickling. Id.
CROCUS sativus. Fields about Saffron Walden.
Ray. Not cultivated there in 1796, nor
could I find any traces of it. D. T.
IRIS fælidissima. Black Notley. Ray. In a hedge
between Chapel End and Higham Hill,
Walthamstow; and in the wood at
Purfleet. Mr. E. Forster. jun.
Scirpus pauciflorus. Bog on Epping Forest, be-
tween Walthamstow and Wanstead. Id.
acicularis. On the edge of the Pond,
called the Bason, in Wanstead Park, not
in Epping Forest, as mentioned by Smith. Id.

Scirpus sylvaticus. By the river Blackwater below Bocking. Robson. Banks of the Rhodon near the Red Bridge, Wanstead. Mr. T. F. Forster, jun. Near Loughton. M. J. Woods, jun. PHLEUM crinitum. Ditches on the Coast : at Purfleet, over against the Mill towards Rainham on the other side the great ditch. Ray. ACROSTIS Spica venti. Cornfields about Walthamstow, abundantly. Mr. E. Forster, jun. - littoralis. On the Essex coast. Mr. Dickson. Pox procumbens. By the road side near Grays Mr. E. Forster, jun. Thurrock. DACTYLIS stricta. At Crixcy Ferry. Merrett. In the marshes on the river Wallfleet. near Fambridge Ferry in Dengey Hundred. Ray. FESTUCA bromoides. About Walthamstow, com-Mr. E. Forster, jun. mon. uniglumis. On the Shore near South End. Mr. E. Forster. jun. Mersey Island, near Colchester. Dale. .---- loliacea. Marshes near Walthamstow, Mr. E. Forster, jun. abundant. AVENA fatua. On the Shore near South End, and Harwich. Id. pubescens. In a Marsh near Walthamstow, where there is no chalk within sixteen miles. Id.

ARUNDO Epigejos. On Epping Forest : near Salter's buildings, Walthamstow ; and on the borders of a wood, called the Larks, near Chingford. Mr. E. Forster, jun. HORDEUM maritimum. Common on the Coast. Id. TRITICUM loliaceum. On the Shore near South End. Mr. E. Forster, jun. Sea shore at Harwich. Dale. DIPSACUS pilosus. Hedges between Wanstead and Barking Side. Mr. T. F. Forster, jun. Between Chingford and Waltham Abbey ; and between Harlow, and Parndon. Mr. E. Forster, jun. Near Friday Hill house in the lane leading to Chingford Church. Warner. GALIUM tricorne. Cornfields about Newport, abundantly. Mr. E. Forster, jun. CENTUNCULUS minimus. Bog on Epping Forest near Highbeech. Is RUPPIA maritima. In a ditch by the road side, between Heybridge and Goldhanger, near Maldon. Id. ANCHUSA sempervirens. By the road side at Great Yeldham. Id. CYNOGLOSSUM sylvaticum. At Purfleet, Mr. T. F. Forster, jun. Hedges at Walthamstow, and Chingford, and on Hen-

hault Forest. Mr. E. Forster, jun.
On the London Road between Kelvedon
and Witham; but more plentifully about
Braxted by the way side. Ray.
Asperugo procumbens. Is introduced as an Essex
plant by Warner, but from his reference
to Ray it is clear that he meant Lycopsis
arvensis. Mr. E. Forster, jun.
MENYANTHES nymphaoides. In the river Rhodon
almost opposite the eight mile stone in
the road to Woodford Bridge near the
bridge; and between that and Lux-
borough House. Warner.
CAMPANULA hederacea. Bog on Epping Forest
near Highbeech. Mr. E. Forster, jun.
VIOLA hirta. South End: near Yeldham, and
elsewherc. Mr. E. Forster, jun. On
Inks Green near Hale End, Waltham-
stow. Miss Woods.
RHAMNUS catharticus. Hedges about Waltham-
stow. Mr. E. Forster, jun.
VINCA major. Near Colchester. Dr. Richardson.
CHENOPODIUM hybridum. Church Yard at Sible
Hedingham. D. T.
olidum. By way sides near Wal-
thamstow. Mr. E. Forster. jun.
SALSOLA fruticosa. In Wallasea Island. Mr.

Dickson. On the Western end of the Marsh Bank at Harwich, plentifully. Dale. CUSCUTA europæa. On nettles at Castle Hedingham. D. T.BUFLEURUM rotundifolium. Cornfields about Ugly, and Newport. Mr. E. Forster, jun. Near Purfleet. Mr. T. F. Forster, jun. ----- tenuissimum. In Marshes near Maldon : on a Common between Warley and Harestreet, at a considerable distance from any salt-marshes. Mr. E. Forster, jun. South End. Mr. J. Jansen. By the road side near Thorndon. Blackstone. PEUCEDANUM officinale. At Walton, near Harwich. Ray. SIUM latifolium. In the Rhodon near Wanstead. Mr. J. Woods, jun. Common in ditches. Mr. E. Forster, jun. ----- repens. On Hatfield Forest; and in the road from Heybridge to Maldon. Id. SISON segetum. About Walthamstow : Chingford : Gray's Thurrock : South End, and elsewhere. Mr. E. Forster, jun. On a Bank in Hagger Lane at the end of Hale-end Lane, Walthamstow. L. W. D. OINANTHE pimpinelloides. Marsh ditches near Purfleet. Mr. E. Forster, jun. In Bulvan Fen. Blackstone. VOL. I.

Т

CORIANDRUM sativum. About Coggleshall:	
Tolesbury, and other places. Ray. Un-	
der the Cliff, below South End.	
Mr. E. Forster, jun	0
SCANDIX Gerefolium. Formerly found in Clay	
Street, Walthamstow, but now destroyed	
by the Bank being cut away.	
Mr. T. F. Forster, jun	8
PIMPINELLA magna. In Woods about Henham.	
Mr. E. Forster, jun. In Widdington	
Wood. Mr. T. F. Forster, jun	•
SAMBUCUS Ebulus. In a Lane leading to Upton.	
Curtis. Ditch in the lane opposite Ham	
Hall. L. W. D. In the Castle Ditch	
at Pleshy: but no longer to be found in	
Warner's Habitat near Waltham Abbey.	
Mr. E. Forster, jun	P
LINUM usitatissimum. Occasionally at Waltham-	
stow, but perhaps not indigenous : on the	
shore near South End. Id.	
MYOSURUS minimus. Cornfields about Waltham-	
, stow. Id.	
ALLIUM oleraceum. At Black Notley. Ray	•
Acorus Calamus. In a Pond at Deux Hall,	
Lamburn: lane between Chigwell and	
Henhault Forest. It also grows in a	
pond, opposite where a House, called	
Harts, the residence of Warner, author of	

27-1

Plantæ Woodfordienses, formerly stood ; and in the River Rhodon, near Woodford bridge road ; but in these two places certainly planted by Warner. Mr. E. Forster, jun. JUNCUS Forsteri. On Henhault Forest, between Hoghill and Collyer row. Id. FRANKENIA lævis. On the shore near Wakering, below South End. Id. ALISMA Damasonium. In pools and gravel pits on Epping Forest, near Walthamstow, not uncommon, and in a Pond near Rumford, as mentioned by Ray. Mr. E. Forster, jun. Plentiful in the Pools about Woodford. L. W. D. EPILOBIUM roseum. By the road side at Moreton near Ongar; and by a rivulet near Great Parndon Church. Mr. E. Forster. jun. CHLORA perfoliata. Borders of Fields about Rickling. Mr. E. Forster, jun. In the chalk pit and cornfields near Newport, and in fields near the Broom, Henham. Mr. T. F. Forster. jun. POLYGONUM minus. Bogs on Epping Forest. Mr. E. Forster, jun, - Bistorta. Field behind Henham Vi-carage. Id. PARIS quadrifolia. Woods about Henham and Quendon. Mr. E. Forster, jun. Crack's

Wood, Henham; and Widdington Wood. Mr. T. F. Forster, jun. DIANTHUS caryophyllus. Tower on the wall at East Ham. L. W. D. SAPONARIA officinalis. In the wood at Purfleet; and in a hedge at Laindon hills. Mr. E. Forster, jun. SILENE anglica. Cornfields about Colchester. Blackstone. ----- noctiflora. In a cornfield between Rickling and Wicken; and in fields opposite Wicken Church, plentifully. Mr. T. F. Forster, jun. SEDUM album. Walls at Westham Abbey, Plaistow, and Barking. Mr. E. Forster. jun. EUPHORBIA platyphylla. Black Notley. Ray. In a cornfield, near Woodford. Mr. E. Forster, jun. β. In a cornfield near Rickling. Id. PRUNUS Padus. Lane leading from Angel Lane to Temple Mills near Low Layton. Warner, PYRUS torminalis. Plentiful in hedges between Blackmore and Ingatestone. Rev. J. Davies. On Epping Forest, and elsewhere in woods and hedges. Mr. E. Forster, jun. Rosa tomentosa. Hedge in a lane between Upmin-Id. ster and Great Warley. TORMENTILLA reptans. In Braintree parish. Ray.

ACTEA spicata. In a thick wood two miles from Thorndon. Blackstone. RANUNCULUS parviflorus. By the road at Leigh; and in a cornfield above South End. Mr. E. Forster, jun. In the Lee Bridge road, and elsewhere about Walthamstow. Mr. B. M. Fors'er. In a lane leading from the house, once the habitation of the learned and celebrated Mr. Ray, at Black Notley. Sir T. G. Cullum. ----- * paroulus. Near South End. Mr. E. Forster, jun. HELLEBORUS fatidus. In the hedge opposite High Laver. Mr. T. F. Forster, jun. Ajuga Chamapitys. In Cornfields near Purfleet. Mr. E. Forster, jun. MENTHA sylvestris. River side between Harlow and Sabridgeworth. Id. rotundifolia. Roadside at Wendon, near Saffron Walden. Mr. E. Forster, jun. At Walthamstow. Mr. T. F.

* Mr. Forster first cleared up the error attending this plant, and shewed that it must be expunged from the British Flora, by some specimens from South End, which, upon comparison were found to agree exactly with Mr. Dyer's from Bristol, and with what is preserved in the Linnzan Herbarium, but which, as he remarked, were evidently only starved procumbent plants of *Ranunculus birsutus*. *D. T.*

Forster, jun. In a meadow going to Faulkburn Hall. Ray. MENTHA viridis. Side of a pond in a field two miles from Thorndon. Blackstone. $----- \beta$. River side at Bocking near the Fulling Mill, in two or three places. Ray. side of the way going from the Fulling Mill to Harris' Mill. Id. ------ S. Meadow on the right hand of Marwood Bridge, leading from Mersey Island towards Colchester. Id. - piperita. Near Clay Street and Whip's Cross, Walthamstow. Mr. E. Forster, jun. - hirsuta. d. By Bocking river side a little above the Fulling Mill, and in a ditch near the Water Mill. Dale. of Chalk's Farm, going from Braintree to Leez House. Dillenius. - rubra. Mark House Lane, Waltham-Mr. E. Forster, jun. stow. -gracilis. At the hop ground at Bocking, plentifully. Dale. By the side of the Lea between Higham Hill and Chingford. Mr. B. M. Forster. gentilis. By the side of a mill-stream

near a house, called the Grove, at West-Mr. T. F. Forster, jun. ham. MENTHA Pulegium. Near Ponds and gravel pits on Epping Forest. Mr. E. Forster, jun. GALEOPSIS versicolor. In a cornfield at the bottom of Ribton Lane, near Woodford. Warner. BALLOTA nigra. B. In a wood, called the Hawk, L. W. D. near Chingford. THYMUS Nepeta. Way sides about Henham; Stanstead, Mount Fitchet; Chelmsford; Billericay, and elsewhere in many parts of the County. Mr. E. Forster, jun. ANTIRRHINUM Cymbalaria. On an old wall at Stubbers. Mr. E. Forster, jun. West-Mr. T. F. Forster, jun. ham. ------ spurium. Cornfields near Henham. Mr. E. Forster, jun. SCROPHULARIA vernalis. In the hedges at Hempsted, in great abundance. Sir T. G. Cullum. Moist hedges by the road side between Sampford and Hempsted, plentifully. Mr. E. Forster, jun. ALYSSUM sativum. By the Horse and Groom near Lea Bridge, but scarcely wild. Id. LEPIDIUM latifolium. At Heybridge near Maldon. Mr. E. Forster, jun. Near the Hythe at Colchester; and by Fulbridge

near Maldon Town's End. Ray. Marshes near Grays. Dr. Watson. LEFIDIUM ruderale. Near South Benfleet. Blackstone. On the shore at St. Osyth, Harwich; and near the river at Maldon. Mr. E. Forster, jun. THLASPI arvense. Near a house, called the Grove, at Westham. Mr. T. F. Forster, jun. Fields about Wormingford, plentifully, as also at St. Osyth. Ray. About Ongar and Chelmsford. Blackstone. COCHLEARIA anglica. On the Banks of the Thames ; and near the river at Maldon. Mr. E. Forster, jun. Salt Marshes near Harwich. Dale. _____ danica. On a Bank by the sea side near Manningtree. Id. IBERIS nudicaulis. In an old gravel pit on Epping Forest, not far from Ilford, near a gate opposite Eastham Lane. Mr. B. M. Forster. CRAMBE maritima. On the sea shore between the Town of Harwich and the Cliff. Dale. CARDAMINE amara. Bog on Bentley Common near South Weald. Mr. E. Forster, jun. About Braintree. Ray. TURRITIS glabra. Hedge near Mistley. Mr. E. Forster, jun. Banks by the high way side

as you go up the hill from Lexden to Co'ch ster. Ray. BRASSICA orientalis. Cornfields and Cliffs on the sea side near Harwich. Id. ---- campestris. At Harwich. Rev. Dr. Goodenough. ERODIUM m. schatum. At Dedham. Mr. T. F. Forster, jun. LATHYRUS Aphaca. Cornfields near the chalk pits, Newport. Mr. T. F. Forster, jun. Corn near Ongar and South End. Mr. E. Forster, jun. --- Nissolia. In Higham hill field, and near Hagger lane, Walthamstow, and elsewhere. Mr. E. Forster, jun. Near Milkwell Field between the end of Horn Lane and the Woodford Bridge road; and in a field near Chingford Church. Warner. - birsutus. In several parts of Rochford Hundred; and at Hadley Castle two miles from Leigh. Ray. Border of a field between the road and Mundon Church, and among bushes within the gate leading to Laydon Hall, both in Dengy Hundred. Blackstone. Near the Shore above South End. Mr. E. Forster, jun. Bor-

ders of Cornfields at Rawreth and Hockley. Rev. R. B. Francis. Astragatus glycyphyllos. Field near the

Thames at Purfleet: roadside between Gray's Thurrock and Stifford ; and near Bulmer Hill in the road to Sudbury. Mr. E. Forster, jun. TRIFOLIUM ornithopodioides. On Epping Forest between Woodford and Woodford Row. Mr. E. Forster, jun. Sandy Banks by the sea side at Tolesbury. Ray. ochroleucum. Fields and roadsides in the Northern part of the County, not Mr. E. Forster, jun. uncommon. ---- scabrum. Near South End. Mr. E. Forster. jun. Meadows near Saffron Walden. Blackstone. ---- maritimum. By the water side at Leigh, and at Little Holland. Ray. About Tilbury Fort. Petiver. On the shore in Canvey Island, plentifully. Blackstone. In a meadow near Hadley Castle; and on Epping Forest near Woodford, with T. ornithopodioides, about forty miles from the sea, and eight even from the Thames. Mr. E. Forster, jun. MEDICAGO polymorpha. B. Sandy shore at Harwich. Dalco

HYPERICUM androsamum. On Epping Forest; and in a wood near Thorndon. Mr. E.

Forster, jun. Near Woodford. Warner.

.

Ditch in a lane on the left hand side of
the road from Brentwood to Ongar, about
half a mile from Ongar. Rev. J. Davies.
TRAGOPOGON porrifolius. Marshes near Purfleet.
Mr. E. Forster, jun.
Sonchus palustris. Marshes near the Thames
below Plaistow. 'Id.
CREPIS fætida. Chalkpits at Purfleet. Id.
<i>biennis.</i> Chalkpits at Purfleet : among
bushes near South End, and elsewhere. Id.
HYOSERIS minima. In the common fields at Wal-
thamstow. Id.
CARDUUS tenuiflorus. Roadsides about Waltham-
flow. Id.
eriophorus. Roadsides about Saffron
Walden; and between Yeldham and
Clare Id.
pratensis. Epping Forest near Hale
End, and Highbeech. Id.
SANTOLINA maritima. At Landguard Fort.
Mr. T. F. Forster, jun. GNAPHALIUM margaritaceum. Plentiful in a
meadow near Bocking, but scarcely wild. Ray.
rectum. On different parts of Ep-
ping Forest, but sparingly. Mr. E. Forster, jun.
gallicum. Among corn in sandy
grounds about Castle Hedingham. Dale.
I have repeatedly sought for it in vain. D. T.

-

INULA Helenium. Hedge by the roadside at Loughton: lane from the Bald-faced Stag on Epping Forest to Chigwell church : near a Farm House at Muncombe near Woodford. Mr. E. Forster, jun. DORONICUM Pardalianches. Hedge in a lane leading from Widdington to a wood, called the Jock. Mr. T. F. Forster. By the roadside near Saling Hall. Mr. Walford. ANTHEMIS nobilis. On Epping Forest. Mr. E. Forster, jun. SATYRIUM viride. Meadow at Henham. Mr. B. M. Forster. Meadow at Ugly, near a wood called Ugly Park. Mr. E. Forster, jun. OPHRYS Nidus avis. On Epping Forest, not far from the gate of a wood, called the Sale, near Hale End, and on a hill called the Hawk. Mr. B. M. Forster. Wood on Laindon hills. Mr. E. Forster, jun. monorchis. Near Newport. Miss Howard. ------ muscifera. Chalk pit near Newport. Mr. E. Forster, jun. ------ apifera. Chalk pits at Purfleet: near Newport; and in the foot-way from Henham to Chickney. Mr. T. F. Forster, jun. ARISTOLOCHIA Clematitis. In a wood two miles from Thorndon. Rlackstone. TYPHA angustifolia. Bog on Epping Forest near

Salter's Buildings, Walthamstow. Mr. E. Forster, jun. Marsh ditches by the right hand side of the Lea Bridge road. L. W. D. CAREX axillaris. Side of a ditch near Ugly Green; and near Rickling Green. Mr. E. Forster, jun. divisa. Near the Hythe at Colchester. Mr. Newton. Banks of the Thames. Mr. J. Woods, jun. ---- pendula. Roadsides about Thaxted, and elsewhere, but not in ditches in Walthamstow Marsh, by Lea Bridge, as stated in Gough's Camden. Mr. E. Forster, jun. About Braintree. Ray. ----- strigosa. In a lane at Black Notley. Dale. ----- Pseudo-cyperus. By a marsh ditch in the road from Walthamstow to Tottenham: in Hoghill Pond on Epping Forest, and clsewhere. Mr. E. Forster, jun. extensa. Near Harwich. Rev. Dr. Goodenough. ----- binervis. Epping Forest, and elsewhere. Mr. E. Forster, jun. ----- distans. Side of a ditch near South End. 14. ----- lavigata. Boggy Thicket near Warley Common. T.I. vesicaria. Banks of the Rhodon; but certainly not by Lea Bridge, as stated in Gough's Camden. Ia.

URTICA pilulifera. On the North side of Harwich Church near the Steeple; and in a meadow on the West side of the gate, plentifully. Dale. In several parts of Canvey Ifland. Blackstone.

SALIX Russelliana. Hedges, common.

Mr. E. Forst	er, jun.
fusca. Epping Forest.	Id.
cinerea. Near Ugly.	Id.
oleifolia. Hedges, common.	Id.
slipularis. Banks of the Lea near Higham	
Hill, Walthamstow; and near Lea Bridge.	Id.
mollissima. Near the Lea Bridge road;	
and in a footpath from Marsh-street,	
Walthamstow, to Lea Bridge.	Id.
lanceolata. Banks of the Lea, near Higham	
Hill, Walthamstow.	Id.
HIPPOPHAE rhamnoides. Near Canvey Island.	.A. L 5 @
	ckstone.
ATRIPLEX laciniata. Near Maldon. Mr. IV.	
Allen. Sandy shore between the town	
of Harwich and the Cliff. Dale. Sea	
shore near Little Holland, and in the	
Isle of Mersey, not far from Colchester,	
plentifully.	Ray.
EQUISITUM sylvaticum. On Epping Forest, near	raje
Highbeech. Mr. B. M. Forster. Boggy	
thicket near, Warley Common.	
, they contract the	

286

Mr. E. Forster, jun.

EQUISETUM fluviatile. Moist places and woods. (Warner mistook this for the preceding Mr. E. Forster, jun. species.) OSMUNDA regalis. Boggy thicket near Warley Common. Not to be found in Warner's habitat. Id. ASPIDIUM Oreopteris. In an Alder Grove not far from Fox hatch, near South Weald. Mr. E. Forster, jun. Epping Forest near Mr. B. M. Forster. Highbeech. ---- lobatum. In hedges, more common than A. aculeatum. Mr. E. Forster, jun. spinulosum. At Walthamstow, and elsewhere in the County. Mr. T. F. Forster, jun. CYATHEA regia. On a wall at Leyton near Walthamstow. Id. PHASCUM axillare. Damp and shady places in Epping Forest, very common. Mr. E. Forster, jun. serratum. On the ground, very common. Id. ----- alternifolium. Epping Forest near Walthamstow. Id. _____ stoloniferum. Epping Forest near a great bog, not far from Salter's buildings, Walthamstow. Id. GYMNOSTOMUM intermedium.] Epping Forestnear fasciculare. S Walthamstow. Id. GRIMMIA Forsteri. On a felled tree near Chapel End, Walthamstow. Mr. T. F. Forster, jun.

¢

GRIMMIA Dicksoni. Rails of paling in Ho	P
Street, Walthamstow. Mr. E. For	
DICRANUM purpureum. B. Gravelly soils in	<i>sici , juli</i> .
Epping Forest.	
	Id.
cerviculatum. Epping Forest, nea	Id.
Woodford.	
ORTHOTRICHUM diaphanum. On trees, commo	Id.
NECKERA beieromalla. On a tree at Loughton.	n. Id.
Bevum hieder On Funine F.	
BRYUM bicolor. On Epping Forest.	Id.
HYPNUM medium. Epping Forest, near Chingford	
hatch.	Id.
inundatum. At the root of a willow	·
near Walthamstow.	Id.
nigro-viride. Trees in Epping Forest,	
	W.D.
lucens. Boggy thicket near Warley	
Common. Mr. E. Forst	er, jun.
crenulatum? Plashet. L. W. D.	
Wood near Hale End, Walthamstow.	
Mr. E. Forst	er, jun.
tenellum. On Chingford Church.	Id.
stellatum. Boggs on Epping Forest, but	
I have not seen fruit.	Id.
JUNGERMANNIA pusilla. On Epping Forest, bo-	
	7.1
tween Woodford Row, and Hale End.	Id.
tween Woodford Row, and Hale End. angulosa. Woody parts of Ep-	Id.
tween Woodford Row, and Hale End. <i>angulosa</i> . Woody parts of Ep- ping Forest.	
angulosa. Woody parts of Ep-	Id. Id. Id.

ANTHOCEROS punctatus. In the common fiel	d
near Walthamstow Church.	
. Mr. E. For.	ster. jun.
LICHEN pinicola. Ach. Elms at Higham Lodge	
Walthamstow.	D. T.
sophodes. Ach. On trees at Rickling.	
Mr. E. For	ster, jun.
tigillaris. Ach. On paling and barn	S
at Walthamstow and Loughton.	
lynceus. Old Oaks in Hainault Forest.	
	Sozverby.
graniformis. Barn at Walthamstow.	
pezizoides. On Epping Forest.	
Mr. E. Fors.	ter, jun.
sphæroides. Trees on Epping Forest, be-	
tween Loughton and Epping.	Id.
inquinans. Posts and rails at Walthamstow	. Id.
tessellatus. On a wall at Walthamstow.	Id.
palmatus. Epping Forest, near Wal-	
thamstow, but no fructification.	Id.
granulatus. On tombstones in church-	
yards.	Id.
subtilis. Epping Forest, near Hale End,	
Walthamstow.	Id.
trapeziformis. Epping Forest, near Wal-	
thamstow.	Id.
pulmonarius. Epping Forest, but no fructi-	
fication.	Id.
VOL. I. U	

-

LICHEN * vermicularis. Among Moss on trees in	
Epping Forest. Don S. de Roxas Cl.	emente.
ULVA pisiformis. Ditches in Walthamstow Marshes	
near Higham Lodge. L.	W. D.
crispa. Shady places under walls.	
Mr. E. Forste	r, jun.
filiformis. fistulosa. In the sea at South End.	Id.
dichotoma.	
Fucus vesiculosus. ζ. 7 In the Blackwater at Mal-	
Fucus vesiculosus. ζ. } In the Blackwater at Mal- don.	Id.
amphibius. Shore of the Blackwater at	
Maldon.	Id.
CONFERVA reticulata. Rivulet near Low hall,	
Walthamstow.	Id.
fracta. At Maldon.	Id.

* This may certainly be considered an extraordinary station for a plant, which, I believe, has never been found before in England, except on the Cumberland Mountains, but I am certain there is no mistake, as Mr. Clemente shewed me abundance of specimens gathered at this place. D. T.

FLINTSHIRE.

CHARA bispida. Stagnated places of water about	
Rhyd Marsh, not uncommon. Mr. Grij	fith.
VERONICA spicata. Above the Lead Mines near	
Prestatyn. Mr. Griffith. On the Hill	١
of Diserth Castle. Bing	ley.
montana. In the upper wood at Tower	
near Mold, and other woods about Mold.	
Mr. Griff	ith.
SCHOENUS compressus. Marsh about a mile West	
of Prestatyn, on the Sea coast. Id.	
albus. Boggy fields between Plâsne-	
wydd in Rhyl and Rhyd Marsh near	
Rhyddlan. Id.	
ERIOPHORUM vaginatum. Common on the	
mountains. Id.	
MILIUM lendigerum. About Trellewelin Farm in	
the Parish of Rhyddlan. Id.	
IT 2	

FLINTSHIRE.

FESTUCA vivipara. In all mountainous pastures.	
	Griffith.
bromoides. Sea coast at Rhyl.	Id.
BROMUS arvensis. Not uncommon in Wales.	Id.
LOLIUM arvense. Very common in cornfields in	
the Parish of Rhyddlan, particularly in	
Rhyl.	Id.
TRITICUM loliaceum. Sea coast between Rhyl	
and Prestatyn.	Id.
DIPSACUS pilosus. Road side from Mold to Wrex-	
ham, opposite the Gate leading to Tyd-	
dyn; also in the road hedge between	
Pentrobin and Mr. Eyton's, Leeswood,	
near Mold.	Id.
SANGUISORBA officinalis. In the Meadow below	
the old Bridge at Pentre, between Mold	
and Chester.	Id.
POTAMOGETON gramineum. Ditches about Rhyd	
Marsh.	Id.
* ANCHUSA semper-virens. Among the ruins of	
Basingwerk Abbey near Holywell. Mr.	
Griffith. Parish of Whiteford. P	ennant.
PRIMULA farinosa. Marshes near the mouth of	
the river Dee. Mr. O	. Sims.

* The Note relative to this plant in Bingley's Tour, was, though unacknowledged, communicated by Mr. Griffith, who examined Mr. Waring's specimer of *Pulmonaria angustifolia*, and therefore knows with certainty that it was merely *Anchusa sempervirens*.

CAMPANULA latifolia. Whiteford. Pennant. By the road side between Holywell and Ches. tcr, about two miles from the former. Rev. H. Davies. Shady places about Lceswood, and elsewhere in Mold's Dale. Waring. CHIRONIA pulchella. Sea coast near Prestatyn. Mr. Griffith. VINCA major. On the common near Rhyd y cilgwyn Bridge between Denbigh and Ruthin. Id. GENTIANA campestris. By the road leading from Holywell to Rhyddlan, about three miles from Holywell, growing with G. amarella. Bingley. SCANDIX odorata. On the Banks of the river Terrig between Leeswood and Nerguis, Parish of Mold. Mr. Griffuth. Plentiful for the length of 300 yards on both sides of the Brook Terrig, and in a field, called Y Packi, near Leeswood; and sparingly by the same brook a considerable distance below, in a field called Maes Madoc: in the Church Yard at Kilken. Waring. ORNITHOGALUM umbellatum. Meadows adjoining Basingwerk Abbey. D. T. JUNCUS acutus. Rhyddlan Marsh. Mr. Griffith. RUMEX maritimus. Rhyd Marsh near Pres-ALISMA ranunculoides. S tatyn, in abundance. Id. COLCHICUM autumnale. Under a Beech tree

FLINTSHIRE.

between upper Leeswood House and the river near Mold. Mr. Griffith. CHLORA perfoliata. Fields at the sea coast near Rhyl. Mr. Griffith. On the side of the hill by the road side leading from St. Winifred's well at Holywell to the sea coast. Bingley. Old pastures near Downing. Rev. H. Davies. VACCINIUM Oxycoccos. Bogs in Wales, very common. Mr. Griffith. POLYCONUM Bistorta. Moist meadows in the Front of Plas On, in the Parish of Mold. Id. PARIS quadrifolia. Sparingly in a wood in Treithin; and in one at Broncoed; both near Leeswood. Waring. CHRYSOSPLENIUM alternifolium. Moist places in the upper wood at Tower, and elsewhere about Mold, very common. Mr. Griffith. SAXIFRAGA stellaris. About most mountainous rills. Id. SAPONARIA officinalis. Among the ruins and rubbish of Basingwerk Abbey. Bingley. SILENE nutans. Rocks above the mine works at Dalea Gôch. Mr. Griffith. STELLARIA Nemorum. In a hedge close to the river about 100 yards above the Ford at Rhyd y Ddae Dwfr, betwixt St. Asaph and Rhyddlan, and on the Rhyddlan side of the River. Bingley.

ARENARIA verna. Between Holywell and St.
Asaph, on the Rubbish of the mines.
Mr. Griffith. Hilly parts of Whiteford
Parish. Pennant.
PRUNUS Padus. Between Mold and Nerguis.
Mr. Griffith. Woods and hedges about
Mold, very common. Bingley. Fre-
quent in the County. Waring.
GEUM rivale. Frequent about Mold. Id.
PAPAVER hybridum. Cornfields near the Store
House upon Rhyl Marsh. Mr. Griffith.
CISTUS marifolius. Rocks above the Lead Works
near Prestatyn. Mr. Griffith. On the
Hill of Diserth Castle, near Rhyddlan,
abundantly. Bingley.
THALICTRUM minus. On Diserth Castle Hill. Id.
RANUNCULUS Lingua. Great Pool at upper Lees-
wood near Mold. Mr. Griffith.
MENTHA sylvestris. In the road as you go from
the river to upper Leeswood House near
Mold. Id.
gentilis. Pools and brooks between Mold
and Llanrwst. Bingley.
LEONURUS Cardiaca. Hedge on the right of the
road leading from Hawarden to Holywell,
about two miles and a half from Hawar-
den. Id.

ANTIRRHINUM Cymbalaria. Walls about upper
Leeswood near Mold. Mr. Griffith.
LIMOSELLA aquatica. Rhyd Marsh near Presta-
tyn. Id.
LEPIDIUM latifolium. Hedge Banks near the
Store House between the village of Rhyd-
dlan and the Sea. Id.
CHEIRANTHUS sinuatus. Sea Beach near Pres-
tatyn. Id.
ERODIUM maritimum. Plentiful about Flint Castle. Id.
GERANIUM phaum. Wood above Mr. Pennant's
garden at Downing near Holywell. Bingley.
pratense. About Rhyd y Mwyn in
the Parish of Mold. Mr. Griffith. In
the field which is next above the Bridge,
called Pont Llong, and between the
Rivers Alen and Terrig in the Town-
ship of Leeswood. Waring.
sanguineum. Near Diserth Castle.
Bingley. Rocks above the Lead works
near Prestatyn. Mr. Griffith.
VICIA sylvalica. Wood below Pont yr Allt Gôch
near St. Asaph. Id.
bithynica. By the road side about two miles
from St. Asaph on the way to Chester.
Rev. H. Davies.
HYPERICUM dubium. Near Downing, plentifully. Id.
TRAGOPOGON porrifolius. Moist marches in the
Parish of Whiteford near Holywell, Bingley.

FLINTSHIRE.

INULA Helenium. Near the old Chapel at Wygfair near St. Asaph. Mr. Griffith. SATYRIUM viride. Meadows between the House of Frôn and the upper wood in the Parish of Mold. Id. SERAPIAS longifolia. Bog near the South East corner of Mr. Lloyd's Garden at Wŷgfair near St. Asaph. Id. CAREX binervis. On all the Moors of Wales. Id. LITTORELLA lacustris. Shores of all the Alpine lakes. Id. QUERCUS sessiliflora. Very common in Wales. Id. EMPETRUM nigrum. On the driest part only of the Mountain, called Gwern-to. Waring. EQUISETUM byemale. Near Maesmynnan. Rev. H. Davies. LYCOPODIUM alpinum. Moors in Wales, very common. Mr. Griffith. ASPIDIUM Orcopteris. Very common in all mountainous pastures. Id. DICRANUM varium. Abundantly among the Mine Pits. Rev. H. Davies. RICCIA fruticulosa. Trees near Holywell on the Road to Denbigh. D.T.LICHEN concentricus. Rocks of Whin-stone in the Parish of Whiteford. Rev. II. Davies.

GLAMORGANSHIRE.

SALICORNIA fruticosa. Salt House Point, near Swansea. Dr. Turton. Cromlyn Bog, near Swansea. CHARA bispida. ----- flexilis. Mr. J. Woods, jun. SALVIA pratensis. Limestone meadows about Port Eynon. Dr. Turton. SCHENUS Mariscus. Cromlyn Bog, near Swansea. L. W. D. ERIOPHORUM vaginatum. Frequent throughout the County. Id. ALOPECURUS bulbosus. In Cardiff marshes. Rev. J. Lightfoot. HORDEUM maritimum. Not unfrequent on the L. W. D. shore about Swansea. TRITICUM loliaceum. Sandy meadows about Penrice, twelve miles West of Swansea. Dr. Turton.

POLYCARPON tetraphyllum. Sandy Wastes be-	
tween Pyle Inn and the Sea. Dr. 7	urton.
SANGUISORBA officinalis. Common in boggy Pas-	
tures about Swansea. L. I	W. D.
ANCHUSA semper-virens. At Bagland, near	
Neath. Mr. J. Wood	ls, jun.
PULMONARIA officinalis. Woods between Neath	
and Pyle. Dr. 2	Turton.
CAMPANULA hederacea. Frequent in Neath Valley	
and other parts of the County. L. W.	
D. Plentiful in the moist places about	
Pont y Pridd. Sir J. 6	Cullum.
LOBELIA dortmanza. Lakes near Pont Nedd	
Vachn, and Aberpergam. L.	W.D.
VIOLA hirta. Peninsula of Gower, not unfrequent.	Id.
DATURA Stramonium. Not uncommon on Dung-	
hills about Swansea.	Id.
CHIRONIA pulchella. On Cromlyn Burrows, near	
Swansea, abundant. L. W. D. Lime-	
stone Hills about the Wormshead. Dr. 7	Turton.
RHAMNUS Catharticus. Woods at Penrice; and	
on the flat Holmes.	Id.
VINCA major. Hedges about Parkmill, between	
Swansea and Penrice.	Id,
CENANTHE pimpinelloides. Landore Marshes near	
Swansea. I. I	V. D.
LINUM angustifolium. Frequent about Swansea.	Id.
DROSERA longifolia. Cromlyn Bog, near Swansea.	
Dr. 7	urton.

GLAMORGANSHIRE.

SCILLA verna. Wormshead, plentifully. Mr. J. Woods, jun.
Asparagus officinalis. Meadows between Cow-
bridge and the Sea; and about Cardiff.
Dr. Turton.
JUNCUS acutus. Common on the sand hills about
Swansea. L. W. D.
ALISMA ranunculoides. Sketty Bog near Swansea. Id.
CHLORA perfoliata. Frequent in the Peninsula of
Gower, and other parts of this County. Id.
VACCINIUM Oxycoccos. Mountains near Pont
Nedd Vachn. Id.
ERICA vagans. Near Newton. Evans. Mr.
Milne told me he had seen specimens ga-
thered on a heath near Portardylais,
about ten miles from Swansea; and Dr.
Turton once told me he had a wild
specimen of E. mediterranea brought him,
which was gathered somewhere in the
neighbourhood. L. W. D.
SAXIFRAGA hypnoides. Mountain East of Pont
Nedd Vechan. M. J. Woods, jun.
SAPONARIA officinalis. Burrows about Marino,
near Swansea; and plentifully between
Cardiff and Llandaff. Dr. Turton.
DIANTHUS caryophyllus. Walls of Cardiff Castle. Id.
SILENE anglica. Fields about Marino, near Swan-
sea. L. W. D.

GLAMORGANSHIRE.

CERASTIUM tetrandrum. Sand hills near Swan-Mr. J. Woods, jun. sea. EUPHORBIA Portlandica. About Caswell Bay; and various other parts of the Coast. L. W. D. PRUNUS Padus. Hedges about Merthyr Tidvil, and Pont Nedd Vachn. Id. PYRUS torminalis. Sides of Neath Valley. L. W. D. Woods about Penrice. Dr. Turton. ----- Aria. Woods about Neath Valley; Merthyr Tidvil, &c. not unfrequent. L. W. D. POTENTILLA verna. Above the Cliffs between Port Eynon and the Worms-head. Mr. J. Woods, jun. GEUM rivale. Usgoed Eynon Garn. Mr. J. Woods, jun. About Pont Nedd Vachn. L. W. D. PAPAVER cambricum. About Pont Nedd Vachn, Aberdylais, &c. plentiful. Id. CISTUS marifolius. On the Worm's-head, plentiful. Mr. J. Woods, jun. * RANUNCULUS parvulus. Marshy field between Swansea and St. Helen's. L. W. D. TROLLIUS europæus. Between Pont Nedd Vechan and Usgoed Eynon Garn. Mr. J. Woods, jun. HELLEBORUS fatidus. Near Park Mill towards Pennard Castle by Swansea. Dr. Turton.

* See Note p. 277.

MENTHA piperita. Town Ditch near Rutland
Place, Swansea. Dr. Turtor.
ANTIRRHINUM repens. Ballast Banks, Britton
Ferry. Mr. W. Clayfield.
SIBTHORPIA europea. Under a damp shady wall
on the left about 200 yards before you
come to Pont y Pridd from Cardiff.
Sir J. Cullum.
DRABA aizoides. Walls and inaccessible rocks
about Pennard Castle eight miles west of
Swansea. Dr. Turton. Frequent also
on the rocks about the Worms-head. L. W. D.
LEPIDIUM petræum. Walls of Pennard Castle. 1d.
ruderale. Ballast Banks about Swansea. Id.
THLASPI alpestre. About Pont Nedd Vachn and
Aberpergam. Id.
COCHLEARIA officinalis. Banks of the Tawey, be-
tween Swansea and Landore, plentifully.
L. IV. D.
anglica. Banks of the Tawey a little
above Swansea. Mr. J. Woods, jun.
danica. Abundant on the rocks
about the Mumbles Light House, and in
similar situations on the Coast. L. W. D.
BERIS nudicaulis. Not unfrequent about Swansea. Id.
CRAMBE maritima. Rocks about Port Eynon. Dr. Turton.
SIEYMBRIUM murale. Plentiful about Swansea.
L.W.D.

CHEIRANTHUS sinuatus. Frequent about Swan-
sea, and on the shores of this County. L. W. D.
TURRITIS kirsuta. Walls of Oystermouth Castle. Id.
GERANIUM pratense. Very frequent about Pont
Nedd Vachn, and Aberpergam. Id.
sanguineum. Burrows about Penrice
and Port Eynon, plentiful. Dr. Turton.
LATHYRUS sylvestris. In Caswell Bay, near
Swansea. L. W. D.
HYPERICUM androsamum. Frequent about Car-
diff, Neath, Penrice, and throughout the
County. Id.
dubium. Plentiful about Penrice. Id.
CARDUUS heterophyllus. Road side between Llan-
daff and Cowbridge. Id.
pratensis. Cromlyn Bog, and other si-
milar situations about Swansea. Id.
GNAPHALIUM dioicum. About Pont Nedd Vachn. Id.
INULA Helenium. Pastures at Port Eynon. Dr. Turton.
crithmoides. Plentiful about Port Eynon.
Dr. Turton. Less plentiful in several
places between Port Eynon and Swansea.
L. W. D.
CINERARIA palustris. Sandy Meadows on the
Sea side near Aberavon. Ray.
PYRETHRUM maritimum. Salt House Point, and
other places near Swansea. L. W. D.
OPHRYS Nidus avis. Margam Woods; and woods

GLAMORGANSHIRE.

about Penrice. Dr. Turton. Woodsnear Pont Nedd Vachn. Mr. J. Woods, jun. OPHRYS apifera. About Penrice and Margam. Dr. Turton. CAREX dioica. Boggy places near Aberdylais Water-fall. L. W. D. 7 Cromlyn and Sketty Bogs ----- curta. ----- ampullacea. S near Swansea. Mr. J. Woods, jun. OSMUNDA regalis. Frequent in this County. L. W. D. POLYPODIUM vulgare, S. (P. cambricum. L.) On a rock in a wood near Dennis Powys Castle, not far from Cardiff. Ray. ----- Dryopteris. Rocks about Pont Nedd Vachn. L. W. D. Woods in the deep Glens of this County. Mr. J. Woods, jun. ASPIDIUM Lonchitis. Valley going up Pen Craigy, from Pont Nedd Vechan. Id. ASPLENIUM marinum. About Oystermouth, plentiful. Id. ADIANTUM Capillus Veneris. In Barry Island; and at Port Kirig. Lbruyd. SCOLOPENDRIUM Celerach. Common about Swan-L. W. D. sea. HYMENOPHYLLUM tunbrigense. Abundant on the moist rocks about Pont Nedd Vechan. Id. PILULARIA globulifera. Pen Craigy Llyn Vach, a mountain Pool near Pont Nedd Vechan. M. J. Woods, jun.

ANDREA alpina. About Llyn Vach near Aber	-
pergam. Mr. G. S	owerby.
TORTULA humilis. In a lane by St. Helen's near	
	. W. D.
DICRANUM polyphyllum. Common in this County	. Id.
cerviculatum. Burrows opposite St.	
Helen's near Swansea.	Id.
POLYTRICHUM urnigerum. Plentiful about the	
Waterfalls in the neighbourhood of Pont	
AT 11 WY 1	W.D.
HYPNUM implexum. Wood near Bryn Mill by	
Swansea.	Id.
JUNGERMANNIA emarginata. purpurea. About Llyn Vach near Aberper- gam. Mr. G. Sa	
JUNGERMANNIA emarginata. near Aberper-	
gam. Mr. G. So	werby.
ULVA dichotoma. filiformis. plumosa. Fucus dasyphyllus. Rock at the Mumbles. pear	
filiformis. Frequent on the rocks	
plumosa.) about Swansea. L.	W.D.
Fucus dasyphyllus. Rock at the Mumbles, near	
Swansea.	Id.
ligulatus. Shore at Oystermouth, and Cas-	
well Bay.	Id.
mervegicus. Rocks by the Mumbles Light-	
house near Swansea.	Id.
pygmaus. Abundant along the Coast.	Id.
kaliformis. }	
CONFERVA Plumula. 5 Shore near Swansea.	Id.
VOL. I.	

GLAMORGANSHIRE.

GLOUCESTERSHIRE.

In our list of this County we have particularly to beg the indulgence of our readers for a kind of inaccuracy, which in a work, like the present, it is impossible wholly to avoid. By far the greater part of the information we have been able to obtain respecting the Botany of Gloucestershire has related to those plants which grow on St. Vincent's rocks, or elsewhere in the immediate vicinity of Bristol, and whether these ought to be arranged under the prefent County or Somersetshire is what we have in many instances had no means of determining. Some of them we therefore fear are referred to wrong Counties, and others are, probably with equal impropriety, placed in both; a circumstance we thought it right to mention, that we might not mislead, more than could be avoided, our friends who may visit this part of the Kingdom, and who, we flatter ourselves, will hold us acquitted of mistakes, which, we are sorry to say, we have found inseparable from the nature of our undertaking.

VERONICA montana. Plentifully at Shortwood, Pucklechurch. Mr. Swayne. SALVIA pratensis. Wick Cliffs. Id. SCIRPUS acicularis. Near Baptist Mills, Bristol. Mr. Dyer. ALOPECURUS bulbosus. Meadows below St. Philip's, Bristol. Id. PHLEUM paniculatum. Bardminston Park. Herb. Banks. POA procumbens. Between Bristol Hot Wells and Clifton. Mr. E. Forster, jun. FESTUCA bromoides. Brandon Hill, Bristol. Mr. Dyer. BROMUS diandrus. St. Vincent's Rocks. Id. ----- pinnatus. On Broadway Hill. Mr. Rudge. arvensis. Cornfields about Bristol. Mr. Dyer. AVENA pubescens. On Wick Cliffs. Mr. Swayne Near Badminton. Mr. Dyer. ------ fatua. Cornfields about Bristol. Id. HORDEUM maritimum. Banks of the Avon. Id. DIPSACUS pilosus. Near Stapleton Bridge. Id. GALIUM tricorne. Cornfields at Pucklechurch. Mr. Swayne. RUBIA peregrina. On St. Vincent's Rocks. Mr. E. Forster, jun. RIBES alpinum. Near Bristol Hot Wells. Waring. RHAMNUS catharticus. Woods in this County. Mr. Dyer. PIMPINELLA dioica. St. Vincent's Rocks. Mr. E. Forster, jun. SISON segetum. Cornfields about Bristol. Mr. Dyer.

GLOUCESTERSHIRE. 309

SAMBUCUS Ebulus. Dursley. Mr. Dyer. GALANTHUS nivalis. In this County, I think wild. Id. ORNITHOGALUM umbellatum. On the top of a Hill, three miles on this side of Bristol. Merrett. SCILLA autumnalis. Clifton, on a point of land not far from an old Windmill. Mr. Dyer. ASPARAGUS officinalis. Marshes below Bristol. Merrett. Marsh near Thornbury. Mr. Dyer. CONVALLARIA multiflora. Wood at Dursley. Id. RUMEX digynus. On walls at Tortworth. Mr. Baker. MONOTROPA hypopithys. Woods near Uley. Id. PYROLA rotundifolia. Woods in this County. Mr. Dyer. minor. ARENARIA tenuifolia. St. Vincent's Rocks. Id. SEDUM dasyphyllum. Walls about Bristol. Id. ----- album. Wick Cliffs. Id. ----- rupestre. St. Vincent's Rocks. Mr. E. Forster, jun. On Wick Cliffs. Mr. PYRUS Aria. Swayne. On St. Vin-POTENTILLA verna. cent's Rocks. Mr. E. Forster, jun. ANEMONE pulsatilla. Common above Sir J. Paul's, Rodborough. Mr. Baker. HELLEBORUS fatidus. Woods in Tortworth Park. Id. MELAMPYRUM sylvaticum. On Wick Cliffs. Mr. Swayne. ANTIRRHINUM minus. Under St. Vincent's Rocks. Mr. E. Forster, jun.

ANTIRRHINUM Cymbalaria. Walls about Clifton. Mr. Dyer. LEPIDIUM ruderale. St. Vincent's Rocks: at Clifton under a wall, and almost every where about Bristol. Id. HESPERIS inodora. On Coteswould Ridge, by the road side between Frogmill and Cheltenham. Earl of Dartmouth. TURRITIS hirsuta. On St. Vincent's Rocks. Mr. T. F. Forster, jun. On Wick Cliffs. Mr. Savayne. ERODIUM moschatum. By the foot path under St. Vincent's Rocks. Mr. E. Forster, jun. maritimum. Banks of the Avon. Mr. Dyer. GERANIUM rotundifolium. St. Vincent's Rocks. Mr. J. Woods, jun. _____ sanguineum. On the left hand side the lower road from Cheltenham to Gloucester, two miles from the former place. Withering. TRIFOLIUM scabrum. On Wick Cliffs. Mr. Savayne. HYPERICUM androsamum. Wood at Wick Cliffs. Id. INULA Helenium. In a pasture ground near Wick Cliffs. Id. CENTAUREA solstitialis. By the hedges not far from Cirencester. Bobart. ORCHISustulata. On Wick Cliffs. Mr. Szwayne. CYPRIPEDIUM Calceolus. A Gentleman, who is a Botanist, resident at Woodchester, near

Stroud, informed me, when I was there, that he had found this plant in a wood very near his house, and described it very Mr. 7. F. Forster, jun. accurately. SERAPIAS grandiflora. Woods near Ulcy. Mr. Baker. ---- rubra. On a steep stony Bank, sloping to the South, on Hampton Common. Mrs. Smith. CAREX pendula. Near Frocester. Rev. W. Wood. ampullacea. In a Pool in the Lakewood near Pucklechurch. Mr. Szwayne. fliformis. Shortwood near Pucklechurch. Id. Buxus sempervirens. At Boxwell in Coteswold. Rav. POLYPODIUM Dryopteris. In woods North East of the road up Frocester Hill. Mr. Baker. SCOLOPENDRIUM Ceterach. Walls about Bristol. Mr. Clay field. NECKERA beteromalla. At Badminton. Mr. W. Sherard. LICHEN resupinatus. 7 barbatus. } In the Forest of Dean. Dillenius. deustus, St. Vincent's Rocks. Hudson.

PINGUICULA lusitanica. Common in this county. Hudson. In bogs near Stoncham, and in places a little North of Southampton.

Sir J. Banks. UTRICULARIA minor. Forest of Bere. Pulteney. IRIS fatidissima. Bitterne. Pulteney. Near the shore between St. Lawrence and Sheephill, in the 1ske of Wight. Mr. E. Forster, ium. Common in the Lie Ture i

jun. Common in the Isle of Wight.

Mr. J. Woods, jun.

PANICUM Crus Galli. By the rivulet side near Petersfield. Ray.

PHLEUM crinitum. Old Salt Pans at Drayton, about two miles from Portsmouth. Pulteney.

MILIUM lendigerum. Near Ride, I. W. Mr. J. Woods, jun. AGROSTIS setacea. In the higher parts of the New Forest, plentifully. Withering. Near Lymington. Earl of Gainsborough. DACTYLIS stricta. On the banks of the Southampton River by high water mark, in great plenty. Sir 7. Banks. FESTUEA ovina. B (F. rubra. With.) About Stoney Cross in the New Forest, plentifully. Withering. BROMUS arvensis. Fields near Southampton. Sherard. ELYMUS europæus. In the woods a mile West of Petersfield. Merrett. Chawton Park, Mr. J. Woods, jun. near Alton. TRITICUM loliaceum. At Yarmouth, I. W. Mr. E. Forster, jun. DIPSACUS pilosus. Watery lanes between Hambledon and Wickham. Pulteney. In the short and long Leigh at Selborne. White. GALIUM tricorne. Liphoeck. Martyn. RUBIA peregrina. About Ride, I. W. and very plentiful in the under-cliff between Luccomb and Bonchurch. Mr. J. Woods, june RUPPIA maritima. By the Ferry over the river Itching, near Southampton. Mr. T. F. Forster, jun. PULMONARIA officinalis. In a wood by Holbury

House, in the New Forest. Ray. Near Kinswood. Merrett. Exbury Wood. Mr. Rudge. PULMONARIA virginica. In the ruins of an old castle near Netley Abbey, far from any house, and apparently wild. Rev. Norton Nicholls. In a wood, through which the road passes, about two miles and a half from Newport, I. W. to Ride, as common as Scilla nutans in our woods. Mr. Griffith'. CAMPANULA Rapunculus. Pastures and hedges on the North West side of the Moor, not far from the great bog, near Petersfield. Merrett. PHYTEUMA orbiculare. Chalky hills by Maple Durham. Merrett. On the Downs of this County. Withering. Meonstoke. Pulteney. ---- var. flore albo. About Buriton. Id. DATURA Stramonium. At Ride, I. W. Mr. S. Woods. CHIRONIA pulchella. THESIUM linophyllum. Flower Down, near Winchester. Pulteney. Basingstoke. Martyn. Brading Downs, I. W. Mr. J. Woods, jun. Chalk cliffs on the South side of the Isle of Wight. Dr. Stokes. VINCA major. To the South of Yarmouth, I. W. plentifully. Withering.

HERNIARIA glabra. Sandy shores at Portsmouth. Martyn. CUSCUTA europæa. Near Lake, I. W. Mr. J. Woods, jun. CAUCALIS latifolia. Cornfields about Crook's Hudson. Easton. SIUM latifolium. Near Fordingbridge. Dr. Maton, Sison segetum. About Maple Durham. Goodyer. OENANTHE pimpinelloides. Near Ride, I. W. Mr. S. Woods. peucedanifolia. At East How, in the Patish of Subborton, seven miles from Petersfield. Merrets, SAMBUCUS Ebulus. Between Luccomb and Bonchurch, I. W. Mr. S. Woods. Among the rubbish and ruined foundations of the Priory of Selborne. White. Near Carisbrook Castle, I. W. and near Housborn. Mr. E. Forster, jun. TAMARIX Gallica. On the Beach near Hurst Castle, and Freshwater. Pulteney. LINUM usitatissimum. In the Isle of Wight. Id. ----- angustifolium. Near Ride, I. W. Pulteney. About Shanklin. Mr. S. Woods. DROSERA longifolia. In the bogs of Bin's pond, Selborne. White. Bogs in the New Forest. Mr. E. Forster, jun. anglica. Forest of Bere. Pulteney. SCILLA verna. Near Newport, I. W. Id. autumnalis, Priory, I. W. Mr. J. Woods, jun.

ASPARAGUS officinalis. At Christ Church, and
Freshwater, I. W. Pultenev.
CONVALLARIA Polygonatum. Chawton Park, near
Alton. Mr. J. Woods, jun.
multiflora. Woods about Bram-
dean. Withering. Under the walls of
Silchester. Mr. Fardon. Woods about
Appleshaw. Mr. Borrer.
COLCHICUM autumnale. In a small wood at
Appleshaw. Id.
ALISMA ranunculoides. Plentifully about the river
Sopley. Pulteney.
EPILOBIUM angustifolium. About two miles be-
fore you come to Alton from Ashton, by
the side of a copfe. Ray. In the road
from Chichester to Arundel.
Mr. E. Forster, jun.
CHLORA perfoliata. Bordean Hill; and in the
Isle of Wight. Pulteney. Banks in the
King's Field, Selborne. White. Road-
side between St. Lawrence and Shanklin,
I. W. Mr. E. Forster, jun. Near
Carisbrook Castle, I. W. Dr. Maton.
VACCINIUM Oxycoccos. In the bogs of Bin's
Pond, Selborne. White,
ERICA vulgaris. var. fl. albis. Forest of Bere:
and in the Isle
var. fl. albis. J of Wight. Pulteney.

DAPHNE Mezereum. In Selborne Hanger among the shrubs at the South East end above the Cottages. White. Woods near Miller. Andover, plentifully. MONOTROPA hypopithys. Holt Wood. Pulleney. North West end of Selborne Hanger under the Beeches. White. By the foot way through the under-cliff from Luccombe to Bonchurch. Mr. J. Woods, jun. CHRYSOSPLENIUM alternifolium. Near Hedly. Merrett. SAPONARIA officinalis. Odiham. Pulteney. ----- var. fl. plenis. Road between Cheriton and Bramdean. Id. SILENE anglica. Near Newport, I. W. Curtis. Very common about Shanklin, I. W. Mr. J. Woods, jun. Alverstoke. Pulleney. In the Cornfields between Itching Ferry and Netley Abbey. Mr. T. F. Forster, jun. ----- no Eliflora. Alverstoke. Pulteney. EUPHORBIA platyphylla. In the Isle of Wight. Dr. Stokes. Matterly Farm, about four miles from Winchester towards Abresford. Dr. Sibthorpe. _____β. (E. stricta. E. B.) Selborne. Mr. Yaldon. Pyrus domestica. Frequent in the Isle of Wight. Dr. Wavell. - Aria. New Forest. Dr. Withering. Be-

tween Basingstoke and Popham Lane. Mr. E. Forster, jun. Near Brading, I. W. Mr. S. Woods. Rosa tomentosa. Isle of Wight. Id. GEUM rivale. Andover. Mr. Borrer. ADONIS autumnalis. Matterley Farm. Pulteney. RANUNCULUS parviflorus. At Lymington, plentifully. Withering. HELLEBORUS viridis. In the deep stony lane on the left hand just before the turning to Norton Farm, Selborne; and at the -top of Middle Dorton under the hedge. White. fætidus. All over the High wood and Coney-croft-hanger, Selborne. White. Near the 39th mile stone in the road to Basingstoke. Mr. E. Forster, jun. Ajuga chamepitys. Longwood warren. Pulteney. TEUCRIUM chamadrys. Carisbrook Castle, I. W. Id. MENTHA sylvestris. At Great Waltham. Mr. E. Forster, jun. In the Isle of Wight. Mr. S. Woods. --- rotundifolia. In the under-cliff between St. Lawrence and Niton, I. W. MIr. J. Woods, jun. Roadside near Alton. Mr. E. Forster, jun. piperita. Near Ride, I. W. Mr. 7. Woods, jun.

MENTHA gracilis. G. In the New Forest.	Sole.
MELITTIS Melissophyllum. In the New Forest	•
and near Netley Abbey.	Hudson.
MELAMPYRUM cristatum. By Netley Abbey.	Pulteney.
LATHRÆA squamaria. Longwood.	Id.
* ANTIRRHINUM repens. In the little lane from	n
Cowes Castle up the hill on the righ	ıt
hand. Mr. T. F. For	ster, jun.
minus. Cornfields between Itcl	1-
ing Ferry and Netley Abbey.	Id.
† OROBANCHE carulea. Near Alton. Ray	V +
Steephill, I. W.	Pulteney.
ramosa. B. Fl. Ang. In thi	is
County. H	udson.
LEPIDIUM ruderale. On the shore at Southamp)-
ton. Mr. T. F. For	ster, jun.
COCHLEARIA anglica. In the Isle of Wight. Da	r. Stokes.
CRAMBE marilima. Western Court.	Pulteney.
GERANIUM pyrenaicum. Widley.	Id.
LAVATERA arborea. At Hurst Castle, over agains	t
the Isle of Wight.	Ray.
PISUM maritimum. Sandown Beach, I. W.	Pulteney.

* Sir Joseph Banks found a variety of this plant, with flowers like the *Peloria*, on stubble fields near Southampton in the autumn.

LATHYRUS sylvestris. In great quantities under the cliff at Shanklin and Luccomb, I. W. INIr. J. Woods, jun. ----- latifolius. Sandown Beach, I. W. Pulieney. VICIA sylvatica. In a wood near the East Turnpike at Appleshaw. Mr. Borrer. ASTRAGALUS bypoglottis. Carisbrook Castle Hill. Mr. Griffith. HYPERICUM androsamum. Buriton Lane, New Forest : Bere Forest. Pulteney. Stoney hollow lanes at Selborne. White. Steephill; and between Luccomb and Bonchurch, I. W. Mr. J. Woods, jun. CARDUUS eriophorus. Bere Forest, and elsewhere in the County. Pulteney. Between Luccomb and Bonchurch, I. W. Mr. J. Woods, jun. Near Overton; Popham Lane; and Basingstoke. Mr. E. Forster, jun. pratensis. Boggy soil in the New Forest near Setley. Id. GNAPHALIUM rectum. Frequent in the South-West parts of the Isle of Wight. Mr. J. Woods, jun. Tussila Go farfara. var. foliis variegatis. Between Hambledon and Meonstoke. Pulteney. INULA Helenium. Waste places about Freshwater, and elsewhere in the Isle of Wight. Waring. Near Ashey, I. W. Mr. J. Woods, jun.

INULA crithmoides. Marsh near Hurst Castle, plentifully. Ray. CINERARIA integrifolia. Near Basingstoke, and Andover. Hudson. Belhan, I. W. and Flower Down near Winton. Pulteney. ANTHEMIS nobilis. In the Isle of Wight. Mr. J. Woods, jun. Near Lymington. Dr. Maton. ORCHIS ustulata. Flower Down near Winton. Pulteney. ----- conopsea. fl. albis. ---- pyramidalis. fl. albis. Bordean Hill. Id. SATYRIUM viride. Near Lodge, Bere Forest. Id. OPHRYS Nidus avis. Holt and Wickham Woods. Pulteney. Schorne, in the Long Lith under the shady beeches among the dead leaves, in Great Dorton among the bushes, and on the Hanger, plentifully. White. ---- Monorchis. Bordean Hill. Pulteney. ------ muscifera. Bordean Hill; and Hambledon. Id. ----- apifera. Winchester : Bordean Hill : Steephill, and Carisbrooke, I. W. Id. ----- * var. albida. Bordean. Id. MALAXIS paludosa. Bere Forest near Wickham. Id.

* Dr. Pultency in the Hampshire Repository, whence so large a proportion of the present catalogue is borrowed, gives a coloured figure of this exquisitely beautiful variety; and I have seen in the possession of my friend, D. E. Davy, Esq. a drawing of a specimen very similar, found by Mifs Uhthoff near Halesworth, Suffolk.

VOL. I.

 $D, \mathcal{T}.$

SERAPIAS longifolia. Between Shanklin and Godshill, I. W. Mr. J. Woods, jun. CAREX pendula. In the Isle of Wight. Mr. S. Woods. ----- lævigata. Between Alton and Chawton Park. Mr. J. Woods, jun. XANTHIUM strumarium. Three miles from Portsmouth towards London. Ray. (Mr. Woods has sought for it here in vain). MYRIOPHYLLUM verticillatum. Ditches communicating with the Avon near Sopley. Pulteney. SALIX fusca. Near Southampton. Id. Ruscus aculeatus. B. (R. lanus, Linn. Trans.) At Stoke, near Gosport, plentifully. Mr. G. Caley. TAXUS baccata. Apparently indigenous in this Pulteney. County. OSMUNDA regalis. Bere Forest; and Isle of Wight. Pulteney. In the New Forest near Setley. Mr. E. Forster, jun. ASPIDIUM Orcopteris. Near Southampton. Mr. Lambert. SPLACHNUM ampullaceum. About Hitchin Ferry. Mariyn. MARCHANTIA hemispharica. Near Petersfield. Robson. ANTHOCEROS punclatus. Near Nosted, a mile from Petersfield. Dillenius. LICHEN ferrugineus. On flints in the highest parts Withering. of the Isle of Wight. ----- calicaris. Huds. Trees in the New Forest. Mr. E. Forster, jun.

Lichen caperatus.	Bears fruit in the New I	Forest.
	Mr. E	. Forster, jun
pulmonaria	s. On oaks in the New F	orest.
		Withering.
fucoides.	Rocks and wood by the se	
at Gospon	t.	Dickson
vulpinus.	Four miles from Basing	
	id to Salisbury.	Dillenius.
plicatus.	Woodcot wood.	Id.
ULVA dichotoma.)	Shores of the Isle of Wi	
fistulosa. 3		Forster, jun.
filiformis.		
rubra.	Rocks near Chrift Chur	rch. Hudson.
Fucus hypoglossum. A	B. A common parasitical	plant
	ns and tips of other Fuci in	
Isle of Wi		Stackhouse.
ovalis. At	Christ Church.	Hudson.
obtusus.	7	
tenuissimus.	Isle of Wight.	Stackhouse.
discors.	>	
abrotanifolius.)	
fibrosus.	Rocks on the coast of	this
bifidus.	S County.	Hudson,
ceranoides. 1	Jear Christ Church. Dr. (Goodenough.
bifidus. S.		-
ciliatus. β.	Shores of the Isle of Wig	ght.
	Mr.	Maimburg.

Fucus edulis. At Freshwater Gate, I. W. Mr. E. Forster, jun. ----- nodosus. B. Portsmouth. Mr. Gilman. ----- cartilagineus. Freshwater Bay, I. W. Dr. Withering. coronopifolius. Shores of the Isle of Wight. Miss E. Everett. ----- pinastroides. At Freshwater Gate, I.W. Mr. E. Forster, jun. At Portsmouth. Mr. Gilman. ----- kaliformis. Near Christ Church. Hudson. CONFERVA equisetifolia. Isle of Wight. Mr. E. Forster, jun.

FROM Mr. Duncumb's History of Herefordshire, of which the first Volume has very recently made its appearance, we had hoped to have collected materials for a copious list of the plantæ rariores of this County. In this hope, however, we have found ourfelves disappointed, for the Catalogue given by that Gentleman is far from considerable; and, as he has particularized no stations of plants, but merely divided them into the productions of the Northern, Eastern, and centrical parts of the County, it is of no use towards the leading object of our work, that of directing Botanists in their researches. At the same time, as it serves to convey some idea of the vegetable productions of Herefordshire, we have extracted from it all the materials in our power, regretting most heartily that we have not had the means of procuring better information with respect to a part of the kingdom so favorable to the pursuit of Natural History.

VERONICA montana. Near the Malvern Hills. Du	ncumb
LITHOSPERMUM purpuro-caruleum, Northern	
part of the County.	Id.
CAMPANULA patula. Eastern, Northern, and cen-	
trical parts.	Id.
hederacea. Centrical part of the	
County.	Id.
VIOLA lutea.	
VERBASCUM virgatum. Northern part of the	
VIOLA lutea. VERBASCUM virgatum. Blattaria.	Id.
SELINUM palustre. About the Malvern Hills.	Id.
PIMPINELLA divica. Centrical part of the	
County.	Id.
MYOSURUS minimus. Northern parts.	Id.
COLCHICUM autumnale. About the North and	
Center of the County.	Id.
ALISMA natans. Northern parts of the County.	Id.
CHLORA perfoliata. Northern and Eastern parts.	Id.
VACCINIUM Vilis idea. POLYGONUM Bistoria. PARIS quadrifolia. Northern parts of the County.	
POLYGONUM Bistoria. Northern parts of the	
PARIS quadrifolia. County.	Id.
CHRYSOSPLENIUM alternifolium. About the Mal-	
vern Hills.	Id.
SAXIFRAGA hypnoides. 7	
DIANTHUS caryophyllus. Northern parts of the	
deltaides	Īđ.
ARENARIA verna. County.	100
SEDUM rupestre.	

About the Center of the Pyrus torminalis. County. Duncumb. POTENTILLA verna. 7 Northern parts of the S County. GEUM rivale. Id. MENTHA rotundifolia. By the River's side at Lydbrook near Ross. Ray. GALEOPSIS versicolor. Northern parts of MELAMPYRUM sylvaticum. 5 the County. Duncumb. ANTIRRHINUM minus. GERANIUM phaum. About Malvern Hills. Id. LATHYRUS Nissolia. sylvestris. Northern, Eastern, and centrical parts of the County. Id. HYPERICUM androsamum. Northern parts of dubium. I the County. Id. CARDUUS eriophorus. Centrical parts of the County. Id. GNAPHALIUM rectum. INULA Helenium. Northern parts of the ORCHIS ustulata. County. Id. SATYRIUM viride. OPHRYS muscifera. 7 Centrical parts of the Coun------ apifera, S ty. Id. SERAPIAS longifolia. Northern parts of the County. Id. CAREX capillaris. About the Malvern Hills. Id. ----- pendula. SALIX nigricans. Near Shobden Court. Mr. Dickson. EMPETRUM nigrum. Northern parts of the County. Duncumb.

TAXUS baccata. On the Mountains. Gough's Camden.
OSMUNDA lunaria. Northern parts of the
Polypodium Dryopteris. S County. Duncumb.
Aspidium Thelypteris. Centrical parts of the
County. Id.
TORTULA rigida. Near Wigmore. Mr. L. Brown.
DICRANUM strictum. Shobdon Marsh. Id.
FONTINALIS secunda. About the Malvern Hills.
Duncumb.
RICCIA crystallina. In muddy inundated places
near Shobdon Court. Dickson.
Troumy maine
sphæroides.] About the Malvern Hills.
pezizoides. Duncumb.
bæomyces.
bypnorum.
concentricus.
ferrugineus. Northern parts of the
geographicus. County. Id.
ventosus.
Oederi.
niger.
aleurites. Park Pales of Croft Castle. Dickson.
tenuissimus. About the Malvern Hills. Duncumb.
tunæformis. Rocks near Wigmore. Dickson.
marginalis. Near Lucton. Dillenius.
sinuatus. Rocks at Wigmore. Mr. Littl. Brown.

LICHEN omphalodes.	7
LICHEN omphalodes.	Northern parts of the
polyrrhizos.	S County. Duncumb.
furfuraceus.	Pales of the park of Croft Castle. Mr. Littl. Brown.
glaucus.	S Castle. Mr. Littl. Brown.
globiferus.	
fragilis.	Northern parts of the
bispidus.	County. Duncumb.
venosus.	
sylvaticus.	Lucton Vallet. Dillenius.
articulatus.	Gattley Park. Mr. Littl. Brozun.

VERONICA montana. Old chalk pits near Hertford. Mr. J. Woods, jun. IRIS fatidissima. Between St. Alban's and Dunstatle. Blackstone. Near Hertford. L. W. D. Scirpus, pauciflorus. On Cheshunt Common. Mr. J. Woods, jun. _____ sylvaticus. Near Totteridge. Id. AGROSTIS Spica venti. At Cheshunt. Id. MELICA nutans. Puckeridge. Martyn. FESTUCA loliacea. Common about Hertford. Mr. J. Woods, jun. ELYMUS europaus. Near Berkhamstead. Withering. DIPSACUS pilosus. Common about Hertford. Mr. J. Woods, jun. LYSIMACHIA thyrsiflora. By King's Langley. Ray.

CAMPANULA Rapunculus. At Effaton, a mile from Wigmore. Merrett. VIOLA birta. Abundant about Hertford. L. W. D. On the chalk hills of this County. Mr. J. Woods, jun. * CHIRONIA pulchella. Between Hertford and Bayford. Id. HERACLEUM sphon /ylium. B. In a hedge in the road opposite Flamstead. Mr. L. Forster, jun. PIMPINELLA magna. Akeridge. Mr. Woodward On the Bank of a Wood between Flamstead and Redburn. Mr. E. Forster. jun. Near Hertford, not uncommon. Mr. J. Woods, jun. MYOSURUS minimus. Plentiful on the Hyde near Edmonton fields by the Rib near Hertford. Id. ALISMA Damaschium. Totteridge Green. Id. EPILOBIUM angusijolium. Near Cheshunt Church. Mr. J. Woods, jun. Plentifully near Berkhamstead. Mr. Woodward. roscum. By the road side between Hockeril and Sabridgeworth, opposite

* The Chironia here designed by Mr. Woods is of taller and more slender growth than that which is found near Yarmouth, and figured in E. B VII. t. 458. Its habit also is dissimilar, but I can detect no essential marks of difference between them. It approaches very nearly to a fpecimen sent me from Germany, and marked "Chironia ramosissima." D. T.

Wallbury. Mr. E. Forster, jun. At Totteridge. Mr. J. Woods, jun. CHLORA perfoliata. Common in the chalky parts of the County. Id. POLYGONUM Bistorta. By the river-side at Rickmansworth. Mariyn. In a marshy meadow near Redburn. Mr. E. Forster, jun. MONOTROPA hypopithys. Near Tring. Ray. SEDUM dasyphyllum. Market Eit. Martyn. PYRUS torminalis. Near Broxted Park. Mr. E. Forster. TORMENTILLA reptans. About Berkhamstead. Mr. Woodward. ANEMONE ranunculoides. Near King's Langley. Hudson. apennina. Near Berkhamstead. Mr. Goodal. RANUNCULUS parviflorus. Near Hertford. Mr. S. Woods. AJUGA Chamapitys. Between Hertford and Bayford ; and common in the chalky parts of the County. Mr. J. Woods, jun. MENTHA piperita. Wet places in this County. Robson. ANTIRRHINUM spurium. Between Hertford and Bayford; and in the chalky parts of the County, frequent. Mr. J. Woods, jun. minus. About Hertford. Mr. J. Woods, jun. Near Berkhamstead. Mr. Woodzuard. SCROPHULARIA scorodonia. We are authorized

by our friend, Mr. E. Forster, jun. who

supplied the greater part of the botanical information for Gough's Camden, to say that the plant introduced under this name in that work is only a variety of S. nodosa.

GERANIUM nodosum. In a hilly situation between Hatfield and Welwyn. Rev. Dr. Abbot. LATHYRUS Nissolia. About Wormsleybury. Miss Hume. Between Hertford and . Bayford. Mr. J. Woods, jun. sylvestris. At Seale Wood, and other places near Hertford. Id. ASTRAGALUS glycyphyllos. On Royston Heath. Ray. TRIFOLIUM ochroleucum. Between Hertford and Bayford. Mr. J. Woods, jun. MEDICAGO falcata. In the foot path between Walford and Bushy Hill. Doody. HYPERICUM androsamum. Duke of Bridgewater's woods at Askeridge; and near Berkamstead. Mr. Woodward. ____ dubium. At North Mimms. Mr. Sabine. OPHRYS muscifera. B. Found by Dr. Eales near Welling. Ray. apifera. Chalkpits on Northall Common, and towards Rushgreen near Hertford. Mr. J. Woods, jun. MALAXIS paludosa. On the low wet grounds between Hatfield and St. Albans. Ray.

HUNTINGDONSHIRE.

FROM the immediate vicinity of this County to Cambridgeshire, we had hoped to have been able to procure at least a respectable list of its vegetable rarities, but how entirely these hopes have been disappointed is sufficiently evident from the two following pages. Mr. Relhan is indeed almost the only Botanist whom we know to have bestowed any considerable attention upon its Natural History; and this Gentleman some years since furnished Mr. Hutchinson, who then intended to publish a History of Huntingdonshire, with a list of about 800 indigenous plants, describing the habitats of the more rare ones. This list we have unfortunately not been able to procure, nor has Mr. Relhan preserved any copy of it; so that we can only hope some future Botanist will fill up the blank we here find ourselves obliged to leave.

HUNTINGDONSHIRE.

ARUNDO calamagrostis. Ripton wood. Mr. Woodward. ELYMUS europæus. Ripton wood. Mr. Woodward. In a wood on the right hand side of the road leading towards the North beyond the Wheat Sheaf Inn, Alconbury Hill. Row. Richard Relbon. SANGUISORBA officinalis. About Ripton. Mr. Woodward; MENYANTHES nymphæoides. In the backwater at Hemmingford. Rev. Richard Relban. BUPLEURUM tenuissimum. On a bank by the Northern road a little beyond Huntingdon. Ray. * SELINUM palustre. On the banks of Whittlesea Mere. Sir J. Banks. PIMPINELLA magna. Ripton wood. Mr. Woodward. SAMBUCUS Ebulus. Warboys. Id. LINUM usitatissimum. Cornfields at Ripton. Id. PARIS guadrifolia. In Ripton wood. Id. SEDUM dasyphyllum. On walls at Bugden. Hadson. LYTHRUM hyssopifolium. In a field on the left near the road side, nearly opposite the Inn at Alconbury Hill. Sir T. G. Cullum. EUPHORBIA platyphylla. Cornfields immediately opposite Alconbury Inn. D.T. - ---- B. Near Kipton. Mr. Woodward. MELAMPYRUM cristatum. In Ripton wood. Id.

* Sir Joseph Banks has found on this plant the larva of Papilie Miachaon. HUNTINGDONSHIRE.

337

GERANIUM pratense. Ripton. Mr. Woodward. I.ATHYRUS Nissolia. ______ sylvestris. ? Ripton wood. Id. TRAGOPOGON porrifolius. } Ripton. Id. CARDUUS eriophorus. INULA Helenium. About Ripton and Warboys. Mr. Woodward. About Bugden. Hudson. CINERARIA palustris. Near Ramsey. Mr. Woodward. SERAPIAS rubra. I have a specimen marked, "Riston, Huntingdonshire," exactly corresponding with Dr. Smith's description. Id. AMARANTHUS Blitum. At Ripton. Id.

KENT.

SALICORNIA fruticosa. In the Isle of Grain, Dillenius. Near the Isle of Shepey. Sir Hans Sloane. CHARA hispida. B. Near Chiselhurst. Ray. VERONICA montana. In Charlton wood. Martyn. In most of the woods about Dillwyn. Dover. SALVIA pratensis. Near Cobham. Ray. Near Mr. Dickson. Feversham. VALERIANA dentata. Near Rowling Wingham. Mr. Hatton. IRIS fatidissima. Common in Charlton wood; and all about Sandgate Castle. Sherard. Dillavyn. Frequent near Dover. SCHOENUS Mariscus. In Ham Ponds near Eastry. Id. compressus. About Chiselhurst. Ray. In boggy ground at Cockshill, and about Id. Ham Ponds.

Scikpus pauciflorus. About Ham Ponds. Dillwyn. _____ sylvaticus. Charlton wood. Dr. Smith. PRLEUM crinitum. In a Salt Marsh by Erith. Merrett. ALOPECURUS bulbosus. In the first field next the road before you go into Northfleet. Hudson. MILIUM lendigerum. Plentiful in the Isle of Shepey. Id. AGROSTIS littoralis. Near the Powder Magazine, about four miles from Woolwich. Mr. Jackson. MELICA nutans. Charlton wood. Martyn. Pos bulbosa. Among the Sandhills between Deal and Sandwich. Dillwyn. DACTYLIS stricta. Near the mouth of Feversham Creek. -Col. Velley. CYNOSURUS echinatus. Near Sandwich. Hudson. I have repeatedly looked for it in vain. L. W. D. BROMUS arvensis. Gravesend. Martyn. About Bushy Rough, near Dover. Mr. J. Woods, jun. pinnatus. Between Radigund's Abbey and Dover; and on Narbourne Downs. Id. AVENA pubescens. Plentiful about Dartford. Id. ARUNDO colorata. y. In a field in Romney Marsh near Dimchurch. Dillauyn. ELYMUS geniculatus. Near Gravesend. Mr. Dickson. HORDEUM maritimum. Salt marshes at Seabrook, near Hithe. Mr. J. Woods. jun. TRITICUM loliaceum. About Sandwich salt pans. Dillauyn, DIPSACUS pilosus. Beckenham church yard, 22

Mandan T.C.1. 1.1. C.1. M
Martyn. Left hand side of the Turn-
pike Road leading from Deptford to
Lewisham, not far from the latter. Gurtis.
GALIUM tricorne. Isle of Thanet. Hudson.
anglicum. Sandy ground between Dart-
ford and Northfleet; and on a wall at
Farmingham. Hudson.
RUBIA peregrina. Cliffs at Dover, East of the
Caves. Dillwyn. About Langdon Bay,
and Lydden Spout near Dover.
Mr. 7. Woods, jun.
CENTUNCULUS minimus. On Ashford Common. Curtis,
Ротамосстоя gramineum. Near Deptford. Dillenius.
RUPPIA maritima. Plentiful in the marsh ditches
at Shepey. Ray.
LITHOSPERMUM purpuro-caruleum. Near Green-
LITHOSPERMUM purpuro-caruleum. Near Green- hithe. Dr. Latham.
LITHOSPERMUM purpuro-caruleum. Near Green- hithe. Dr. Lathum. ANCHUSA sempervirens. On Boughton Hill, and
LITHOSPERMUM purpuro-caruleum. Near Green- hithe. Dr. Lathum. ANCHUSA sempervirens. On Boughton Hill, and in a Church yard near Feversham. L.W.D.
LITHOSPERMUM purpuro-caruleum. Near Green- hithe. Dr. Latham. ANCHUSA sempervirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about
LITHOSPERMUM purpuro-caruleum. Near Green- hithe. Dr. Latham. ANCHUSA sempervirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about Tunbridge Wells. Mr. T. F. Forster,
LITHOSPERMUM purpuro-caruleum. Near Green- hithe. Dr. Latham. ANCHUSA sempervirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about
LITHOSPERMUM purpuro-caruleum. Near Green- hithe. Dr. Latham. ANCHUSA sempervirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about Tunbridge Wells. Mr. T. F. Forster,
LITHOSPERMUM purpuro-caruleum. Near Green- hithe. Dr. Latham. ANCHUSA sempervirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about Tunbridge Wells. Mr. T. F. Forster, jun. At South End, near Eltham, plen-
LITHOSPERMUM purpuro-cærulcum. Near Green- hithe. Dr. Latham. ANCHUSA sempervirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about Tunbridge Wells. Mr. T. F. Forster, jun. At South End, near Eltham, plen- fully. Sherard.
LITHOSPERMUM purpuro-cærulcum. Near Green- hithe. Dr. Latham. ANCHUSA semperwirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about Tunbridge Wells. Mr. T. F. Forster, jun. At South End, near Eltham, plen- fully. Sherard. SYMPHYTUM officinale. β. In abundance by the
LITHOSPERMUM purpuro-cærulcum. Near Green- hithe. Dr. Latham. ANCHUSA semperwirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about Tunbridge Wells. Mr. T. F. Forster, jun. At South End, near Eltham, plen- fully. Sherard. SYMPHYTUM officinale. β. In abundance by the Marsh by the Mill near Penshurst. Forster,
LITHOSPERMUM purpuro-cæruleum. Near Green- hithe. Dr. Latham. ANCHUSA semperwirens. On Boughton Hill, and in a Church yard near Feversham. L. W. D. CYNOGLOSSUM sylvaticum. Sides of lanes about Tunbridge Wells. Mr. T. F. Forster, jun. At South End, near Eltham, plen- fully. Sherard. SYMPHYTUM officinale. β. In abundance by the Marsh by the Mill near Penshurst. Forster, BORAGO officinalis. var. fl. albis. At the entrance

PRIMULA farinosa. Near Eltham. Martyn. (surely this is an error.) CAMPANULA Rapunculus. Hedges between Bexley and Dartford Heath. Mr. E. Forster, jun. PHYTEUMA orbiculare. Beacon Hill. Martyn. VIOLA hirta. Charlton Wood, and Woods near Dover. Mr. J. Woods, jun. Frequent throughout the County. L. W. D. Old chalk pits at Dartford. · D. T. VERBASCUM Lychnitis. Near Farnborough. Mr. T. F. Forster, jun. Near Lewisham; Farningham; and Bromley; Mr. E. Forster, jun. About Dartford, and most parts of the County, plentifully; just by the 17th mile stone on the road from London to Maidstone. L. W. D. Near Chiselhurst: on the North side of Madam's Court Hill. Mr. J. Woods. jun. - Blattaria. By the road side near the 14th mile stone from London to Dartford. Blackstone. Near Horn's place

ford. Blackstone. Near Horn's place by Rochester. Ray. Between Deptford and Greenwich. Merrett. On a bank about three miles from Rochester on the Medway. Mr. J. Rayer.

CHIRONIA pulchella. Near Sandwich; and plentifully above Sandgate. Mr. J. Woods, jun. Common near Tenterden. Mr. S. Woods.

RHAMNUS catharticus. Hedges near Shottenton Hill. Martyn. VINCA major. Between Knowlton and Deal; and near Roehill. Ray. In a field at Beckenham, certainly wild. Curtis. Lanes adjoining East Langdon church : near Eyethorne; and lanes at the back of Hythe. Dillwyn: CHENOPODIUM hybridum. Plentiful about Northfleet. Hudson. ----- olidum. At Woolwich: on the Parade at Dover; and near Sandwich. Mr. J. Woods, jun. GENTIANA campestris. In a very old chalk pit as you go from Dartford Heath to Stanhill. Marlyn. BUPLEURUM rotundifolium. Between Greenhithe and Stone, in a field by the road; and near Lewisham, Martyn. About Dartford. Mr. 7. Woods, jun. tenuissimum. Plentifully in the Isles of Shepey and Thanet. Hudson. CAUCALIS daucoides. About Dartford. Mr. J. Woods, jun. ESHINOPHORA spinosa. Between Feversham and Sea-Salter. Blackstone. Near Sandwich. Gerard. In both these places I have sought for it in vain, L. W. D.

Gerard also says it grows between Whitstaple and the Isle of Thanet. PEUCEDANUM officinale. About a quarter of a mile below Feversham by the river side. Sherard. SIUM latifolium. Northfleet. Martyn. SISON segetum. Near Northfleet. Blackstone. About half a mile from New Romney on Mr. J. Woods, jun. the road to Rye. OENANTHE pimpinelloides. About Sandgate. Mr. J. Woods, jun. Ditches in Sandwich and Romney Marshes. Dillauyn. CICUTA virosa. In running water at Canterbury and Ashford. Mr. W. Borrer, jun. PIMPINELLA magna. Between Canterbury and Sittingbourne; and between Maidstone and Farningham. Martyn. About Dover. L. W. D. Common in the County. Mr. W. Borrer, jun. _____β. Between Sittingbourne and Maidstone. Id. STAPHYLEA pinnata. About Ashford. Parkinson LINUM perenne. Shepey. Martyn. ----- angustifolium. On Beacon Hill. Ray. Near Minster in the Isle of Shepey; Deal, and elsewhere. Hudson. Between Frant and Eridge, near Tunbridge Wells. Mr. S. Woods. On Nourbourne Common: near Hastings; and Dover. Mr. J. Woods. jun.

MYOSURUS minimus. Plentiful in fields about Wingham and Canterbury. L. W. D. LEUCOJUM astivum. Between Greenwich and Woolwich, about half a mile below the former, close by the Thames side just above high water mark. Curtis. NARCISSUS poeticus. Back of Mount Pleasant, Tunbridge Wells. Mr. J. Woods, jun. Near Hougham. Rev. J. Lyon. On a rabbit warren at Shorne between Gravesend and Rochester. Mr. J. Rayer. ---- biflorus. With the preceding. Mr. J. Rayer. In the neighbourhood of Tunbridge Wells towards Penshurst. Mr. T. F. Forster, jun. ALLIUM carinatum. Near Ramsgate; and between Sandwich and Deal. Hudson. oleraceum. Lanes near Rochester. Mr. Winch. SCILLA autumnalis. By the highway to Eltham. Martyn. CONVALLARIA multiflora. Woods about Rochill. Withering. JUNCUS acutus. Sand hills near Deal. Mr. J. Woods. jun. ------ Forsteri. Between Bexley and Dartford Heath; and in many places about Dart-. ford. Mr. E. Forster, jun. FRANKENIA lavis. In the Isle of Shepey, abundantly. Rev. Dr. Goodenough. Salt marshes near Sandwich. Mr. J. Woods,

jun. On the Coast of the Isle of Thanet : and by Archeliff Fort, Dover. L. W. D. RUMEX sanguineus. About Maidstone. Hudson. On Mount Sion by Tunbridge Wells. Forster. ALISMA Damasonium. Bogs on Ashdown Forest. Mr. T. F. Forster, jun. In a pool under the Cliff between Folkstone and Sandgate, sparingly. Dillauyn. ranunculoides. Ditches between Ham Ponds and Eastry; and in the boggy grounds about Wingham. Id. ÉPILOBIUM angustifolium. Woods about half a mile West of River Poorhouse, by Dover. Dillavyn. Maize Hill beyond Greenwich. Curtis. CHLORA perfoliata. Extremely plentiful about Dover. Dillwyn. About Gravesend. Blackstone. VACCINIUM Vitis idæa. Golden Wood, Feversham, towards the North end, abundantly. Id. PARIS quadrifolia. About the middle of Byseing Wood, near Feversham. Blackstone. Woods at Waldershaw. Mr. J. Woods, jun. In the Orchard at the Elms, and in most of the woods about Dover, often with 5, 6, or 7 leaves. Dillauyn. MONOTROPA hypopithys. In a wood near Maidstone. Blackstone.

CHRYSOSPLENIUM alternifolium. In * Charlton Wood. Blackstone. SAPONARIA officinalis. By the road side at Linton. Mr. E. Forster, jun. Blackheath. Martyno _____flore pleno. On the road to Sittingbourne. Id. DIANTHUS caryophyllus. On Rochester and Deal Castles, but not on Sandown Castle. L. W. D. deltoides. On the banks of Beacon Hill near Feversham. Blackstone SILENE quinquevulnera. Sandy fields about Wrotham. Hudson. I have searched for it there in vain. L. W. D. + nutans. Plentiful on the Cliffs about Dover, and on Sandgate Castle. Id. conica. Common on the range of Sand Hills beginning at Deal and running Eastward of Sandwich; and on the Sand Hills direstly "opposite the Warren-house at New Romney," sparingly. Dillauyno

* I never heard of its being found there, not do I believe it : C. oppositifelium abounds there. L. W. D.

[†] Although Ray has described the Dover Plant as another species, he suspects it is the same as that which grows on Nottingham Castle Rock, and this is undoubtedly the case. S. paradoxa, 1 believe, does not grow about Dover. L. W. D.

STELLARIA glauca. Side of clear springs about Tunbridge Wells. Mr. T. F. Forster, jun. ARENARIA tenuifolia. In a Close on the left hand going down a Hill a mile from Deptford towards South Fleet. Ray, SEDUM album. Walls and stony places about Tunbridge Wells. Mr. T. F. Forster, jun. sexangulare. Near Northfleet : Sheerness ; and in the Isle of Shepey, plentifully. Hudson. On the South side of Greenwich Park wall near the Western Corner. Curtis. LYCHNIS dioica. floribus albis plenissimis. On the sides of the North end of the Old Haven at Sandwich. Dillwyn LYTHRUM hyssopifolium. Ditches near the Abbey Pond at Feversham. Jacobs EUPHORBIA Portlandica. Bogs by Charlton Wood. Martyn. (This certainly must be a mistake, but we have inserted it, because we are not aware what species could be intended.) ----- platyphylla. Near Northfleet. Hudson. Near Chiselhurst. Mr. J. Woods, jun. ----- biberna. Between Feversham and Sittingbourne. Hudson PYRUS Aria. On Wrotham Hills: Madam's Court Hill, near Seven Oaks, and elsewhere. Mr. E. Forster, jun. Old chalk pits at Dartford. D.T.

TORMENTILLA reptans. Neighbourhood of Tunbridge Wells towards the High Rocks. Mr. 7. F. Forster, jun. PAPAVER hybridum. At Darent near Dartford. Mr. Lewin. Edge of the cliff near Archcliff Fort, Dover. Dillruyno Argemone. S. (P. maritimum. With.) Near Sandown Castle, Id. ---- somniferum. Cornfields about Dartford. Mr. S. Woods. Near the Medway at Rochester. Mr. Winch. DELPHINIUM Consolida. Fields between Blackheath and Eltham. Dillenins. ANEMONE ranunculoides. Near Wrotham. Hudson ADONIS autumnalis. In the closes between Stonechurch and Queenhithe, in great plenty. In the cornfields about Dartford. but more abundantly in the Marshes by the side of the Thames between Dartford and Greenhithe. Mr. J. Woods, jun. RANUNCULUS Lingua. Between Rotherhithe and Deptford. Hudson. In the Old Haven at Sandwich; and in and about Ham Ponds. Dillwyn. ----- parviflorus. Near Green Street green, not far from Deptford. Hudson. HELLEBORUS viridis. On the right hand side of the walk by the Parsonage House in

Sir N. D'Aeth's woods at Knowlton, about seven miles North of Dover. Sherard. HELLEBORUS fatidus. Between Northfleet and Gravesend. Martyn. Ajuga reptans. fl. allis. Charlton Wood. Curtis. ---- Chamapitys. Barren fields on the South side of Roe Hill. Sherard. In several places about Rochester; Dartford, &c. Ray. About Tunbridge Wells. Mr. T. F. Forster, jun. At Purfleet. Mr. Jacob Rayer. MENTHA rotundifolia. At Sir F. Leigh's near Hally. Ray. In a ditch near Wrotham. Mr. E. Forster, jun. ---- acutifolia. By the Medway. Rand. Between Rochester and Chatham. Miller. Pulegium. On Chiselhurst Common, very plentiful. Mr. J. Woods, jun. BALLOTA nigra. B. (B. alba.) Near Speldhurst Church, in the Neighbourhood of Tunbridge Wells. Mr. T. F. Forster, jun. THYMUS Nepeta. About Dartford, and many other places in the County. Blackstone. In the old sand-pits at the back of Charlton Church. Curtis LATHRÆA squamaria. On a ditch bank at Bredgate near Sittingbourne. Ray. In the wood South from Chiselhurst bog, the

end towards the bog, on the left hand: also in the woods about Maidstone, plentifully. Sherard. ANTIRRHINUM spurium. In Shepey Island near the sea. Sowerby. Farnborough. Mr. E. Forster, jun. Fields near Ramsgate, on the road to Margate. Dillwyn. Common near Dover, and Dartford. Mr. J. Woods, jun. minus. About Dover. Id. repens. Field's near Gad's Hill. Martyn. OROBANCHE minor. In a field two miles beyond Rochester, on the left hand going towards Horn's Place. Ray. Sand pit at Charlton; and under the Cliffs East of Dover. Mr. J. Woods, jun. ---- ramosa. Isle of Shepey; and near Feversham; and Rochester. Hudson. Near Hougham. Rev. J. Lyon. In a field on the right hand side of the road from Rochester to Maidstone near the 32d mile stone. Mr. J. Rayer. LEPIDIUM latifolium. Isle of Thanet. Martyn. ----- ruderale. On the shore at Ramsgate, and elsewhere. Mr. T. F. Forster, jun. COCHLEARIA anglica. All along the Thames side below Greenwich. L. W. D. ISATIS tinctoria. By the Church at Long Reach. Martyn.

CRAMBE maritima. In St. Margaret's and Lang-
don Bays; and very plentifully on the
beach, about half way from Dover to
Folkstone. Dillwyn.
DENTARIA bulbifera. Sides of rivulets about Tun-
bridge Wells. Mr. T. F. Forster, jun.
Between Tunbridge Wells and Wood-
gate. Mr. J. Woods, jun.
CARDAMINE amara. Lewisham. Martyn. River
head near Seven Oaks. Mr. J. Woods,
jun. On the muddy shore of the Thames
below Greenwich. Mr. E. Forster, jun.
SISYMBRIUM murale. About Ramsgate, and other
parts of the Thanet. L. W. D.
TURRITIS glabra. Near Lewisham. Hudson.
In the lane which leads down by Charl-
ton Church, sparingly. Gurtis. Charl-
ton Sand-pits. Mr. J. Woods, jun.
Near Bromley. Mr. E. Forster, jun.
birsuta. On old walls at Colken's
near Boughton. L. W. D.
BRASSICA campestris. Dover Cliffs. Mr. J. Woods, jun.
oleracea. About the Caves, and Shaks-
peare's Cliff, Dover. Dillauyn.
ERODIUM maritimum. On the walls of Sandgate
Castle. Id.
GERANIUM phæum. At Tovell, in the old Valley
by the Fulling Mill. Ray,

GERANIUM rotundifolium. Near Dartford.
Rev. H. Davies.
MALVA rotundifolia. S. (M. pusilla. With.)
Near Hithe. Sherard.
moschata. floribus albis. Fields near
The set
ALTHÆA birsuta.* In a field near Cobham.
Mr. J. Rayer.
FUMARIA parviflora. In the fields about Wold-
ham near Rochester. Id.
PISUM maritimum. On the West side of Dunge-
ness, near Lyd. Ray. Abundantly on
the beach near Walmer Castle. L. W. D.
LATHYRUS Nissolia. Isle of Shepey. Curtis.
About the 65th mile-stone on the sides
of the London Road near Dover. Dillwyn.
sylvestris. Between Rochester and
North Fleet. Hudson. Side of the road
from Rochester to Canterbury in many
places; and in the dark lane South of
the entrance into Waldershare Park near
Dover. L. W. D.
latifolius. In a hollow lane at Fever-
sham. Jacob.
VICIA lathyroides. On the chalky banks near
Greenhithe. Sherard.
Sherara.

• We have admitted this plant, though unnoticed in the Flora Britannica, on the authority of Mr. Symmon's Synopsis,

ASTRAGALUS glycyphyllos. About Charlton, plentifully. Hudson. TRIFOLIUM ornithopodioid's. On Blackheath. Mr. E. Forster, jun. Between Dim. church and New Romney. Mr. J. Woods, jun. Not unfrequent in any part of the County. L, W, D---- maritimum. In Dartford Marshes. Merrett. Between Greenhithe, and Northfleet. Hudson. In Sandwich salt marshes. Dillwyn. About Sheerness. plentifully. Doody. ---- scabrum. By the Thames side between Northfleet and Gravesend. Dillenius. Between Dimchurch and New Romney. Dillwyn. About Chatham; and in the Isle of Shepey, plentifully. Hudson. ---- glomeratum. About Blackheath and Greenhithe. Doody. Isle of Shepey. Hudson. In Greenwich Park. Mr. E. Forster, jun. Between Dimchurch and New Romney. Mr. J. Woods, jun. Near Sandown Castle. Dillwin. MEDICAGO polymorpha. B. (M. minima. L.) near Sandown Castle. Id. HYPERICUM androsamum. In most of the woods about Dover, as in the wood on Lymne

VOL. I.

3

hill, plentifully : between Chiselhurst and Bromley. L. W. D. TRAGOPOGON porrifolius. Marshes near Long Reach, below Woolwich. Blackstone. Thorne Hop Garden, near Feversham. Jacob. SONCHUS palustris. On the banks of the Thames near Greenwich. Ray. LACTUCA scariola. In a sand pit in Charlton Wood. Mr. E. Forster, jun. ------ saligna. In Erith Marshes. Dale. CREPIS fatida. Charlton chalk-pits. Petiver. Northfleet chalk-pits. Blackstone. Near Greenhithe. Hudson. Near Rochester. Mr. Winch. ----- biennis. Plentifully on the road from Sittingbourn to Rochester. Sherard. About Northfleet, and elsewhere, frequent. Hudson. Chalk-pits at Erith. Mr. J. Woods, jun. Charlton. Mr. E. Forster, jun. Heights near the Horse Barracks, Dover. Dillwyn. Dartford. D. T. HYPOCHÆRIS glabra. Under Greenwich Park Wall on Blackheath. Curtis. At Woolwich Warren. Mr. Winch. CARDUUS tenuiflorus. Common about London. Curtis. pratensis. Meadows about Ore Mill pond near Feversham. Jacob.

CARDUUS eriophorus. About a farm, called Polton, near Radigund's Abbey; and about the ruins of Lymne Castle by Dover. Dillwyn. SANTOLINA maritima. Isle of Shepey. Hudson. GNAPHALIUM rectum. About Charlton. Id. SENEC10 viscosus. About the chalk pits at Dartford. Dr. Smith. INULA crithmoides. Isle of Shepey. Shererd. Sheerness. Martyn. * BELLIS perennis. I found a remarkable variety of this plant, in every respect, except the length of the scapus, more than twice as small as it generally grows, in chalky pastures about Dartford. D. T.ANTHEMIS nobilis. Common on heaths about London. Mr. J. Woods, jun. CENTAUREA solstitialis. Near Northfleet. Hudson. Near Margate. Mr. W. Allen. ORCHIS ustulata. In the foreland meadow, and on the hills about Dover, sparingly. Dillauyn. militaris. B. (O. fusca. Jacq.) Northfleet Sherard. On chalky banks abounding with Milkwort and Juniper near woods, and in the woods themselves in many

* The root transplanted into my garden is in one year become exactly similar to the common Daisy. D. T.

A 2 2

parts of Kent, especially about Rochester, in great abundance.

Curtis.

SATYRIUM bircinum. In the road which leads from Crayford to Dartford. From the street, named Lofield, in Dartford, is a place called Fleate Lane, and on the left hand are several plants of it : also beyond Dartford is a place named the Brent, and on the right hand a great highway going to a village called Grimsteed Green; a little way on the right hand you may also find it. Ray. In chalky pastures by the side of Darent wood, two miles from Dartford; and in Haley Wood Pits near Dartford Heath. Mr. J. Woods, jun. ---- viride. Ospringe Chalk-Pits. Jacob. OPHRYS Nidus avis. About Maidstone. Ray. On Roe Hill, plentifully. Merrett. Woods about Chariton. Hudson. Near the little brook by the High Rocks, Tunbridge, towards Rusthall Common. Forster. Wood between Alkham and Ewel Minnis. Dillwyn. Loeselii. Boggy ground about Ham Ponds near Eastry. Id. anthropephora. Bank Westward of Crabbe. Dillauyn. In Bocton Church

Yard. Jacob Near Northfleet: Greenhithe; and elsewhere in Kent, frequent. Hudson. Pastures at Dartford.

Mr. J. Woods, jun.

OPHRYS muscifera. Chalk-hills near Northfleet. Ray. Plentiful about Wrotham and Greenhithe. Hudson. Haley wood pits near Dartford, Mr. J. Woods, jun. Sunny bank at the side of Coombe Wood, by Dover. Dillavyn.

β. At Greenhithe; and in the old chalk-pit near the white house by North Gray, in great plenty. Sherard.
 apifera. In Haley wood-pits near Dartford. Mr. J. Woods, jun. Common about Dover, and throughout the County. L. W. D.
 aranifera. Frequent on the hills about Northfleet. Ray. On those about Dover, some years much more plentifully than others. Dillwyn.

MALAXIS paludosa. Divers places in Romney Marsh. Ray. At Hurst Hill: Tunbridge Wells. Doody.

SERAPIAS longifolia. Chiselhurst; and near Eltham. Ray. About Ham Ponds near Dover: marshes about Hacklinge, be-

tween Deal and Sandwich; and in Wing-
ham marshes. Dillwyn.
SERAFIAS grandiflora. Roe Hill, near Dartford. Merrett.
ARISTOLOCHIA Clematitis. Near Maidstone,
and in other parts of Kent. Hudson.
Турна angustifolia. In the middle of Woolwich
Common. Curtis. Common in Rom-
ney Marsh. Mr. J. Woods, jun.
CAREX divisa. Very abundantly in the Marshes
by the side of the Thames between
Greenhithe and Dartford. Mr. J.
Woods, jun. Near Ham Ponds, by Do-
ver; and in Sandwich Marshes. Dillroyn.
pendula. Wood on Lymne Hill, by Dover,
plentifully. Id.
depauperata. In a small wood at the back
of Charlton Church. Curtis.
Sandwich; and in Sandwich old Warren.
Dillwyn.
Ham Ponds. Id.
Buxus semper-virens. There are woods of this
plant at Boxley. Ray.
URTICA pilulifera. Parkinson says it has been
found growing of old at Lydde near Rom-
ney, and in the streets of Romney.
Ray. Extinct at Romney, but e little

South of Lyd Church Yard still plentiful. Dillourn. XANTHIUM strumarium. In the road at Dulwich, a little on this side the College, just by the stile, going the footway thence to London. Newton. I have sought for it there in vain. Mr. J. Woods, jun. MYRIOPHYLLUM verticillatum. Ditches between Deal and Sandwich. Dillwyn. Plentiful in Romney Marsh. Mr. J. Woods, jun. SALIX Lambertiana. About Ham Ponds near Sandwich. Id. ----- rubra. Just out of Kent Street, London, on the Dover road. Id. ----- amygdalina. Near Sandgate. Id. argentea. A little North of Sandown Castle, plentifully; and about Walmer Castle. Id. fusca. Common on Heaths about London. Mr. J. Woods, jun. ----- cinerea. In a wood by the road side near Bromley. Mr. E. Forster, jun. mollissima. Common in the South East of Kent. Mr. J. Woods, jun. HIPPOPHAE rhamnoides. In a salt marsh, two miles from Gravesend, Blackstone, Abundant between Folkestone and Sandgate, undercliff; and a little West of St. Margaret's Bay, by Dover. Dillauyn.

TAXUS baccata. Between Tunbridge Wells and the High Rocks. Mr. J. Woods, jun. Certainly indigenous in that neighbourhood. Forster. ATRIPLEX pedunculata. In the Isle of Thanct near the Ferry, in great plenty. Sherard. EQUISETUM sylvaticum. Woods North East of Tunbridge Wells. Mr. J. Woods, jun. ----- fluviatile. In the under-cliff between Folkstone and Sandgate. Mr. J. Woods, jun. In Folkstone Cherry Orchard. Dillwyn. OSMUNDA lunaria. In Scadbury Park; and on Chiselhurst Common. Sherard. LYCOPODIUM inundaium. On the Common before you come to Chiselhurst. Ray. ASPIDIUM lobatum. In Woods and Rocks about Tunbridge Wells ; as on the Rocks going to the Cold Bath Rocks from the High Rocks, near the rivulet. Mr. T. F. Forster, jun. ASPLENIUM septentrionale. On the North side of Bocton Church. Jacob. (Is it not probable that the Author of the Plants. Favershamienses mistook A. ruta muraria for A. septentionale ?) SCOLOFENDRIUM Ceterach. On Chiselhurst, Lenham and Stroud Churches. Martyn. On a Wall at Riverhead near Seven

Oaks. Mr. E. Forster, jun. Wall at the entrance into Ewell Church yard. Dillauyn. GRIMMIA calcarea. Chalk pits at Dartford, in great plenty. Mr. Sozverby-TORTULA cuneifolia. On an old wall, North East side of Dover Castle. L. W. D. POLYTRICHUM attenuatum. On Sydenham Com-Mr. Sozuerby. mon. PTEROGONIUM Smithii. Trees on Barham Downs. Dr. Smith. A little North of the half way house between Dover and Canterbury; and plentifully on the trees in Waldershare Park, by Dover. Dillwyn. NECKERA heteromalla. On trees West of Buckland Church. Id. --- crispa. Chalk Cliffs near Gravesend. Dillenius. BRYUM nutans. Mountainous Heaths about Woolwich. Id. HYPNUM undulatum. Shooter's Hill. Id. tenellum. On old walls at the Priory, and other walls about Dover. Divervin. JUNGERMANNIA pusilla. 7 Banks of Woolwich ----- reptans. Heath in moist ANTHOCEROS punctatus.] shady places. Dillenius. RICCIA minima. Blackheath near Greenwich. Id. MARCHANTIA hemisphærica. In a small water-VOL. I. вh

course near the Wells at Sydenham towards Norwood. Martyn. ULVA dichotoma. Sometimes found on the beach ----- rubra. at Dover. _____ plumosa. Dillauyn. pisiformis. Ditches between Greenwich and Woolwich. Hudson. Sometimes, but not fre-quently, found near Do-ver. Fucus hypoglossum dasyphyllus. Dillwyn. membranifolius. B. Rocks near the Caves, Dover, sparingly. Id. ----- barbatus. Shepey. Martyn. ----- fibrosus. Sometimes washed on the Do-_____ bulbosus. S ver shore after storms. Dillavyn. ----- Norvegicus. Not unfrequent among the rejectamenta of the sea at Dover; and at neap tide I found it on the rocks opposite Shakspear's Cliff near low-water mark. Id. edulis. Rocks near the Caves at Dover, plentifully. Id. _____ plumosus. Rocks about Dover, common. Id. ----- radiatus. articulatus. On the Rocks about Dover. Id. ----- Opuntia. ----- pinastroides. Deal. Martyn. ----- Wigghii. A single specimen found at Folkstone by Miss Eliz. Everitt.

CONFERVA vagabunda.	Salt ditches about Graves-
end.	Dillenius.
equisetifolia. 	On the Beach at Do-
byssoides.	yver. Dillavyn.
pennata. Bet	ween Dover and Margate.

END OF THE FIRST VOLUME.

•

London: printed by Phillips & Fardon, George Yard, Lombard Street, Hudson.





• • ,

,

•

.

Poler 1 & cambrain - How M. Joncesh . _ Poar heilders werten in 1 .

